

delivery of 10,000 yards of the new
fabric daily.

SOMETHING KILLS FISH

During the fire last Wednesday evening some acid or something escaped into Greenbrier river and killed a great number of fish. On Friday morning one of the town's best fishermen went down the river and came back with a string of nine pike. He had a lot of fun telling how he landed them, especially the big one, which weighed nine pounds, but soon he cut the fun and told the story of how he found the fish lying along the banks in great numbers, dead. It is not known whether it was something from the tannery, the excessive heat of the fire, or what it was that killed the fish—but they are dead, and now there is no use for any fisherman to go down the river with the expectation of making a big catch.—*Martinton Journal.*

WEST VIRGINIA GEOLOGICAL SURVEY



Greenbrier County

By

PAUL H. PRICE, State Geologist

E. T. HECK, Assistant Geologist

1939



PLATE I.—View from side of Greenbrier Mountain, featuring the grounds of the White Sulphur Springs, Inc. One of the golf courses is shown in the foreground, the Cashio and the Greenbrier Hotel in the center, and Kates Mountain with typical Devonian topography in the background. Photo. by Cummins.



WHEELING NEWS LITHOGRAPH COMPANY
WHEELING, W. VA.

1939

GEOLOGICAL SURVEY COMMISSION

HOMER A. HOLT.....*President*
GOVERNOR OF WEST VIRGINIA

R. E. TALBOTT.....*Vice-President*
TREASURER OF WEST VIRGINIA

J. B. McLAUGHLIN.....*Member*
COMMISSIONER OF AGRICULTURE

CHARLES E. LAWALL.....*Executive Officer*
ACTING PRESIDENT, WEST VIRGINIA UNIVERSITY

C. R. ORTON.....*Secretary*
DIRECTOR, AGRICULTURAL EXPERIMENT STATION

STATE BOARD OF CONTROL

WALTER R. THURMOND.....*President*

JOHN BAKER WHITE.....*Secretary*

MERRILL D. CARRICO.....*Treasurer*



PLATE LII.—Blue Sulphur Springs. The pavilion shown above is over one hundred years old, mute reminder when this spring was the site of a famous resort. The hotel and bath-houses were destroyed during the Civil War and never rebuilt.



New England Aerial "Aerograph"

PLATE LIII.—Airplane photograph of Rainelle and the lumber yards of the Meadow River Lumber Company.

to give mud-baths. The most probable source of the water is the Webster Springs or Edray Sandstone. Certain physical data and a chemical analysis are given below:

Elevation: 1670'.

Geological Horizon: Bluefield Group Shale.

Temperature: Date observed, 6-3-35, 52.7° F.; 10-3-35, 58.0° F.

Rate of flow: Date observed, 6-3-35, 6 gallons per minute; 10-3-35, 6 gallons per minute.

Owner: Buster Helrs, Address A. M. Buster, Blue Sulphur Springs, W. Va.

Analyst: Homer A. Hoskins.

Constituent.	Parts per Million.
Solids after evaporation.....	1652.0
Ignition loss.....	121.0
<hr/>	
Silica (SiO ₂).....	24.0
Iron (Fe).....	0.24
Calcium (Ca).....	299.0
Magnesium (Mg).....	49.0
Sodium (Na).....	119.0
Potassium (K).....	4.0
Bicarbonate (HCO ₃).....	190.0
Sulfate (SO ₄).....	815.0
Chloride (Cl).....	58.0
Nitrate (NO ₃).....	None
Manganese (Mn).....	0.02
Hydrogen sulfide gas (H ₂ S).....	7.2
<hr/>	
Total of determined constituents.....	1565.46

Alvon Springs Nos. 1 and 2.—These springs are located on the south side of Anthony Creek, 0.5 mile west of Alvon. They provide the main source of water-supply for the town of White Sulphur Springs. In times of large demand the supply of water is supplemented by the Alvon Spring No. 3 and rarely by Alvon Spring No. 4, which springs will be described on a subsequent page. All of these springs are carefully protected in every way. Plate LI shows the spring-house over Nos. 1 and 2, the pump-house, and bottling-house erected just west of Alvon by the White Sulphur Springs, Inc.

As shown by the analysis below, the water is exceptionally pure and it is reported that the quality of this water does not vary materially. The water emerges from near the base of the Helderberg and although the point of emergence of the water is enclosed in such a manner that it can not readily be examined in detail, it appears to be limestone. As a rule limestone water is comparatively high in mineral content and the

by the way of Blue Sulphur Springs, crosses the divide to Sinkias Creek and continues on in that direction toward Trout Valley.

"The limestone section produced excellent timber, free from insect injuries and defects of every kind. It was nearly all hardwood, such as white oak, red oak, poplar, black walnut, hickory, and some wild cherry."

The third district lies in the mountainous sections of the north and northwest, and is characterized by such species as spruce, hemlock, yellow birch, and others that thrive at high altitudes. Even here, however, hardwoods predominate below an altitude of 3,000 feet and sometimes higher up than this. Following is a list of trees and the number of each kind growing on 1,000 acres on the head of Cherry River in this county. Locusts, hickories, and black walnuts with a diameter over 10 inches, and all others over 18 inches were counted*.

White oak.....	132
Chestnut oak.....	889
Hickory	86
Chestnut	1,513
White maple.....	3,258
Sugar maple.....	7,291
Locust	4
Beech	1,965
Birches	1,120
Gum	104
Cherry	349
White walnut.....	1
Poplar	529
Linden	1,014
Cucumber	937
Ash	576
Hemlock	2,303
Yew pine (Spruce).....	34
Total	22,264

THE LUMBER INDUSTRY.

Most of the limestone area, where the best hardwoods grow, was settled and the timber destroyed in the process of clearing the land for cultivation before it could be sold for profit and in a day when timber was considered inexhaustible and of little value. A little of it was utilized for building and fencing purposes and for fuel.

Small water-power sawmills were located here and there in an early day. After those came the portable steam sawmills.

*"Resources of West Virginia"—Maury and Fontaine.

The latter were not common until the Chesapeake and Ohio Railroad was extended westward from White Sulphur Springs about the year 1873. After that time many of these mills were located near the line. The principal shipping points for lumber were White Sulphur Springs, Caldwell, and Roneeverte. When the Chesapeake and Ohio Railroad was built up the Greenbrier River a similar industry was begun all along the line. When available sites for the small mills became scarce near the railroad many of them moved back into the interior where they are still engaged in sawing for small owners and hauling the lumber wagons to the railroad.

The first large band-saw operation in the county was that of the St. Lawrence Boom and Manufacturing Company. This company came to Roneeverte in 1882 and erected a circular mill. In 1884 this was replaced by a double band mill which continued to operate until 1910. During 24 years the mill cut 433,000,000 feet of white pine from Greenbrier and Pocahontas Counties. After 1902 the white pine supply began to fall off and considerable hemlock and hardwood timber was sawed. This company erected a single band mill at Shryock on Anthony Creek in 1909 which it is now operating.

Some of the large operators that have completed their work were the Henderson Lumber Company, with a band mill at the mouth of Anthony Creek; the Clear Creek Lumber Company, and the Kittanning Lumber Company, both with large circular mills in the Greenbrier section.

Among the present extensive operators, some of which have cut over vast forest areas, may be mentioned the Cherry River Boom and Lumber Company located at Richwood in Nicholas County; the Fenwick Lumber Company at Fenwick, Nicholas County, and the Neola Lumber Company at Neola, all band mill operations. Donaldson Lumber Company and Kendall-Deter Lumber Company are operating large circular mills near Anthony on the Greenbrier River.

Much of the fine walnut timber was destroyed. That which remained until after the coming of the railroads was eagerly sought after and even the stumps throughout the Greenbrier Valley were bought and removed.

Chestnut oak timber once grew in abundance in the county and furnished material for an active tan-bark industry which has lasted through a long period of years.

PRESENT FOREST CONDITIONS.

Mr. A. B. Brooks mentions that in 1911 there was approximately 140,000 acres of virgin forest and 105,000 acres of cut-over land in Greenbrier County. At the present time the area of virgin forest remaining in the county has been reduced to a few scattered patches aggregating a few thousand acres. There is one stand of virgin forest on Beaver Lick Mountain about four miles north of Alvon and scattered areas in other parts of the county.

Much of the cut-over land is unfit for anything but forests and as will be described below steps are being taken by the Federal government toward replanting and protecting this land.

MONONGAHELA NATIONAL FOREST.

The purchase area of the Monongahela National Forest extends into Greenbrier County in two prongs, one in the northeast corner and one in the northwest part. The eastern boundary coincides with the State line from the Pocahontas County line, southward to the junction of the State line and the White Sulphur-Anthony Creek District line. The Forest boundary roughly follows the district line to the Greenbrier River just below Anthony. From this point the boundary of the purchase area extends northward along the Greenbrier River to the Pocahontas County line. The boundary of the northwestern prong enters Greenbrier from Pocahontas County at Boggs Run about one mile north of Beulah Church and extends in a straight line to Twin Sugars. The boundary extends northwest to Cold Knob and Grassy Knob. From this point it follows the Meadow Bluff-Williamsburg District line to the Nicholas County line near Lile.

As outlined above the proposed area of the National Forest occupies 210,903 acres in Greenbrier County of which 93,981 acres have already been acquired by the United States Government. The following table, taken from a report of the Department of Agriculture, shows the proposed acreage

of the Monongahela National Forest and the amount acquired in each county. The figures are as of June 30, 1937:

County	Proposed extent, Aeres.	Approved for Purchase, Aeres.	Acquired, Aeres.
Grant	43,700	13,634	13,329
Greenbrier	210,903	96,571	93,981
Nicholas	45,939	23,428	20,286
Pendleton	149,500	58,198	56,321
Pocahontas	537,288	266,987	243,859
Preston	12,192	3,891	3,891
Randolph	361,299	164,692	152,443
Tucker	202,700	87,913	85,860
Webster	110,131	64,707	60,744
Total	1,673,652	780,021	730,713

Under the direction of Mr. Arthur A. Wood, Forest Supervisor, Elkins, W. Va., many improvements have been made on the land already acquired. Several fire trails have been built in Greenbrier County and fire towers have been erected. A beautiful recreation spot has been developed at Blue Bend on Anthony Creek about three miles east of Anthony. The location of this park is shown on Map II. The Forest Service has issued a very interesting pamphlet on the Blue Bend Park and copies of this pamphlet may be obtained at any of the district offices. The improvements made at the park consist of a large log Administration Building, picnic shelter, bath-houses, toilets, and facilities for camping.

LUMBER MILLS.

The following is a list of the larger lumber concerns operating at the present time in Greenbrier County:

Meadow River Lumber Company at Rainelle.

Roneeverte Lumber Company at Roneeverte.

Spring Creek Lumber Company at Spring Creek.

Wilderness Lumber Company at Nallen, Fayette County.

Cherry River Boom and Lumber Company at Richwood, Nicholas County.

In addition to the above concerns there are several small portable sawmills operating in the county.

SOILS OF GREENBRIER COUNTY*

by

Anton J. Vessel,

United States Department of Agriculture,

Bureau of Chemistry and Soils.

The soils of Greenbrier County belong to the Gray-Brown Podzolie group of United States soils. They have developed from various parent materials chiefly under a deciduous forest cover in a humid climate where the winters are not too cold and the summers are not too hot. The soils of the area do not contain much organic matter. In forested areas a thin layer of leaf-mold is mixed with the topmost layers of the surface soil. The soils are dominantly light in color and highly leached of bases and plant nutrients. All of the soils except those recently limed are acid throughout the profile. They respond well to fertilizer and lime treatments. The most important soils are those of the limestone valley. They dominate the agriculture of the county which is centered around livestock raising.

Greenbrier Valley is underlain with Greenbrier Limestone of great thickness. Various members comprising this series cause some variation in the types of the resulting soils. The purer members give rise to the Frederick soils which possess grayish-brown and brownish-yellow surface soils and reddish-yellow or light-red subsoils. Locally some shale is included with the Frederick soils. On steep slopes where great thicknesses of the Greenbrier Formation have been mapped as one type, some areas of Hagerstown soils are intermixed with the Frederick and are mapped as Frederick-Hagerstown stony silt

*The above brief summary of the soils of Greenbrier County shows the relationship of the soils to the geologic formations outcropping in the county. Mr. Vessel has recently completed the field work for a detailed soils map and report of the area. It is expected that the report will be published in the near future by the Bureau of Chemistry and Soils of the United States Department of Agriculture, Washington, D. C. Inquiries concerning the soils of the county should be addressed to the Bureau.

loam. The Hagerstown soils possess a browner surface soil and a darker red subsoil than the Frederick. They are developed from the residue that is left after the solution of limestone of great purity. Siliceous and platy limestone practically free from chert give rise to the Frankstown soils in this county. These differ from the Frederick in having more friable subsoils that are yellow or brownish-yellow in color. Cherty limestones as the Hillsdale member give rise to Frederick cherty silt loam which differs from the type in having a scattering of angular fragments on the surface and throughout the profile. Associated with Frankstown and Frederick soils, but on smoother relief is a small area of Pickaway silt loam that is mottled and slightly plastic in the subsoil. The surface soil is gray or grayish-yellow in color. The soils of the limestone valley dominate the agriculture of the county and are the most productive.

Directly overlying the Greenbrier Formation is the Mauch Chunk Series comprising in ascending stratigraphic order the Bluefield, Hinton, Princeton Conglomerate, and Bluestone Groups. (*) The Bluefield Group occurs as a wide belt running in a southwest-northeast direction throughout the central portion of the county. It is dominantly shaly, comprising yellow and greenish-gray shales together with some interbedded impure limestone. Generally the tops of ridges are capped with sandstone. At the junction of the Greenbrier Series with the Bluefield Group and including the shaly and limestone horizons, the material is very heterogeneous. The soil that has developed from this mass is the Westmoreland. It possesses a grayish-yellow or brownish-yellow surface soil and a yellow-brown friable subsoil. A smooth phase has been recognized in addition to the type soil. Westmoreland silt loam possesses a characteristic surface relief. In many places it is limited in profile development due to the hilly to steep relief.

(*) For a description of these members as well as other geological formations the reader is referred to the West Virginia Geological Survey report and map of Greenbrier County by Paul H. Price and E. T. Heck; West Virginia Geological Survey, Morgantown, West Virginia.

plateau section, and has some fairly level summit areas. However, the land is not suitable for farming because the surface is strewn with boulders.

Closely related to and resembling the Dekalb soils are the Clymer soils. They are developed from the same parent materials, but occur on smoother relief. The loam type occurs on the tops of ridges that are capped with sandstone. The silt loam type generally occurs at lower elevations, and is developed from sandstone and shale material. In the early soil survey mapping, such areas were included with the Dekalb soils. Recently they have been separated from this series because of the deeper profile development and better agricultural adaptation.

The Mareellus Series occurs in the eastern part of the county as a continuous narrow belt occupying the lower mountain slopes and low rounded hills immediately adjacent to the small streams. This series is composed for the most part of finely laminated shales from which is developed Berks shaly silt loam. This is a very shallow soil and is non-agricultural. The surface soil is light-yellow or brownish-yellow in color, and the subsoil is yellowish-brown or brownish-yellow tinged with red.

Below the Mareellus Series are the Oriskany and Helderberg Series. These outcrop on Coles and Beaver Lick Mountains. The Huntersville Chert member of the Oriskany Series contains a small amount of lime and together with the siliceous limestones of the Helderberg gives the Elliber soils. Where not too stony the Elliber soil makes better grass land than surrounding areas of Dekalb or Berks soils. Weathering of the parent material leaves the chert fragments strewn over the surface and throughout the profile. Virgin areas of Elliber soils have a mat of organic matter on the surface which may be 3 or 4 inches thick. The surface soil is gray or grayish-yellow, and the subsoil is brownish-yellow or yellowish-brown. The purer limestone members of the Helderberg Series give rise to Hagerstown soil. Such areas are inextensive and have been included in with the Frederick soils.

In the eastern part of the county rocks of Silurian age are exposed. These are unimportant as far as soils are concerned because the areas are too stony, and are mapped chiefly as Rough stony land.

The soils developed on terraces include the Elk, Holston, Monongahela, and Sequatchie. These soils differ from each other not only in the character of the parent material, but also in the stage of maturity to which they have developed. The youngest of the group is Sequatchie loam. It resembles the Pope soil which occurs on the flood-plain. Elk silt loam is not a fully mature soil. The parent material for this soil is old alluvium from limestone uplands and to a lesser degree from Upshur soils. Elk silt loam as mapped is light-brown in color throughout the profile. The parent materials of the Sequatchie, Monongahela, and Holston soils were washed from areas of Dekalb soils. Monongahela silt loam is a poorly drained soil. It resembles Philo silt loam which occurs on the flood-plain. Holston loam is the most mature soil of this group. It is highly leached and possesses a gray or grayish-yellow surface soil with a light-yellow subsoil.

The alluvial soils include the Moshannon, Pope, Philo, and Atkins. They occur on the flood-plain, and are subject to frequent overflow resulting in deposition of new material. They have not developed a profile because the parent material has not been in place sufficiently long to be altered by the normal soil-forming processes of the region. The Pope and Philo soils resemble each other in the color of the surface soil, being light-brown or brownish-yellow. Philo differs from Pope in that it becomes mottled with gray and rust-brown at depths of 14 to 16 inches. Atkins is a poorly drained soil and is gray throughout the profile. Much of it is marshy throughout the year. Moshannon silt loam is one of the most productive soils. The alluvium has been washed from Upshur and limestone underlain soils. The color of the Moshannon soil resembles the Upshur soils.

CHAPTER XIII.

MINERAL WATERS, WATER-POWER, IRON ORE, MANGANESE, AND PRECIOUS METALS.

MINERAL WATERS.

GENERAL STATEMENT.

In Greenbrier, as in most of the counties in West Virginia lying within the limits of the folded Allegheny Mountains, there are numerous springs, most of which carry sufficient minerals in solution to be classified as mineral springs. Some of the minerals, in several of the springs, precipitate out upon reaching the surface and discolor the spring basins. This fact has resulted in the application of many descriptive terms such as "Blue Sulphur," "Black Sulphur," etc.

Some of these waters have long been used for medicinal purposes. The waters of the White Sulphur Springs are being used on a large scale for this purpose at the present time.

As reported by Price and others¹ some of the springs of the county are reported to be high in salt and salt was manufactured from water obtained from shallow wells in the early part of the nineteenth century. The salt was manufactured at two localities along the Greenbrier River, one being on the east side of the river three miles below Spring Creek and four miles upstream from Anthony Station. The other locality was one mile upstream from Renick P. O., on an island at Burr Ford. Both localities produced the brine from rocks of the Pocono Series. At the present time there are several known salt licks in the county.

¹Price, Paul H., Hare, C. E., McCue, J. B., and Hoskins, Homer A., Salt Brines of West Virginia, W. Va. Geol. Sur., Vol. VIII, pp. 31-2; 1937.

MINERAL SPRINGS.

Much of the data presented in this section has been published in an earlier publication of the Survey². A description of individual springs with chemical analyses of some of the waters tested in the Survey laboratories follows:

Black Sulphur Spring.—This is one of the springs on the grounds of the famous Greenbrier Hotel at White Sulphur Springs. Over this spring has been built a beautiful pavilion that has been the subject of many poems and essays. (See Plate L.) The spring emerges from the Marellus Shales but it appears likely that the actual aquifer is the Oriskany which would be 300 to 500 feet below the surface at this point.

Certain physical data and a chemical analysis of water from this spring follow:

Elevation: 1850'.

Geological Horizon: Marellus Shale.

Temperature: Date observed, 6-3-35, 62.5° F.; 9-25-35, 63.0° F.

Rate of flow: Date observed, 6-3-35, 25 gallons per minute.

Owner: White Sulphur Springs, Inc., White Sulphur Springs, W. Va.

Analyst: Homer A. Hoskins.

Constituent.	Parts per Million.
Solids after evaporation.....	2218.0
Ignition loss.....	155.0
<hr/>	
Silica (SiO ₂).....	17.0
Iron (Fe).....	1.1
Calcium (Ca).....	439.0
Magnesium (Mg).....	125.0
Sodium (Na).....	22.0
Potassium (K).....	1.2
Bicarbonate (HCO ₃).....	205.0
Sulfate (SO ₄).....	1416.0
Chloride (Cl).....	17.0
Nitrate (NO ₃).....	None
Hydrogen sulfide gas (H ₂ S).....	12.5
<hr/>	
Total of determined constituents.....	2255.8

White Sulphur Spring.—This is the spring that gives the famous resort its name. It is located about 100 feet south of the Black Sulphur Spring described above and is very similar to it in every respect. Certain physical data and a chemical analysis of the water from this spring follow:

²Prlee, Paul H., McCue, J. B., and Hoskins, Homer A., Springs of West Virginia, W. Va. Geol. Sur. Vol. VI; 1936.

Elevation: 1850'.

Geological Horizon: Marcellus Shale.

Temperature: Date observed, 9-25-35, 64° F.

Rate of flow: About 30 gallons per minute.

Owner: White Sulphur Springs, Inc., White Sulphur Springs, W. Va.

Analyst: Homer A. Hoskins.

Constituent.	Parts per Million.
Solids after evaporation.....	2057.0
Ignition loss.....	338.0
Silica (SiO ₂).....	17.0
Ferric oxide and Alumina (Fe, Al) ₂ O ₃	2.0
Calcium (Ca).....	362.0
Magnesium (Mg).....	84.0
Sodium (Na) and Potassium (K).....	66.0
Bicarbonate (HCO ₃).....	236.0
Sulfate (SO ₄).....	1355.0
Chloride (Cl).....	16.0
Manganese (Mn).....	0.01
Hydrogen sulfide gas (H ₂ S).....	9.4

Total of determined constituents..... 2147.41

Probably no springs in the State have such a world-wide favorable reputation as do the ones described above. The flow from these springs is reputedly constant, with a constant temperature the year around. The fact that they are warmer than nearly all of the other surface springs of the county supports the thesis that the main source bed lies at some depth similar to the supposed position of the Oriskany at that locality.

White Sulphur Chalybeate Spring.—This is another of the famous springs on the grounds of the Greenbrier Hotel. The small flow, varying temperature, and nature of the water, all suggest that this water is ground water derived from the shale itself and not from the underlying Oriskany. Certain physical data and analyses of the water are given below:

Elevation: 1850'.

Geological Horizon: Marcellus Shale.

Temperature: Date observed, 6-3-35, 59° F.; 9-25-35, 64° F.

Rate of flow: Date observed, 6-3-35, 0.5 gallon per minute.

Owner, White Sulphur Springs, Inc., White Sulphur Springs, W. Va.

Analyst: Homer A. Hoskins.

Constituent.	Parts per Million.
Solids after evaporation.....	88.0
Ignition loss.....	16.0
Silica (SiO ₂).....	4.0
Iron (Fe).....	4.5
Calcium (Ca).....	7.0
Magnesium (Mg).....	1.9
Sodium (Na).....	2.4

Potassium (K).....	1.6
Bicarbonate (HCO_3).....	Acid
Sulfate (SO_4).....	43.0
Chloride (Cl).....	Trace
Nitrate (NO_3).....	None
Hydrogen sulfide gas (H_2S).....	None
<hr/>	
Total of determined constituents.....	64.4

Remarks: This water is acid.

Comments: Of all the samples analyzed by the Survey, this is the only one found to be acid. The water is bottled and sold by the owners and considerable use is made of it on the premises. Much mention of this spring is found in the literature, but the following analysis by Froehling & Robertson, Richmond, Va., is the only one observed.

Constituent.	Parts per Million.
Silica (SiO_2).....	2.3
Alumina (Al_2O_3).....	2.0
Iron (Fe).....	8.39
Calcium (Ca).....	18.46
Magnesium (Mg).....	1.54
Sodium (Na).....	3.64
Potassium (K).....	0.49
Bicarbonate (HCO_3).....	21.0
Sulfate (SO_4).....	52.89
Chloride (Cl).....	3.55
Phosphate (PO_4).....	0.57
Manganese (Mn).....	0.9
Strontium (Sr).....	0.715
Iodine (I).....	0.007
<hr/>	
Total of determined constituents.....	116.452

Remarks: Recalculated to free radicals by B. R. Drake from folder issued by the owners.

Blue Sulphur Springs.—This spring is situated in a meadow beside the town of the same name, nine miles north of Alderson. It is reported to have once been a natural fountain that spouted vertically from the ground and a famous buffalo-lick. An early owner caused gravel to be dumped into the spring until it became a placid pool. In the early part of the nineteenth century many improvements were made around the spring, including a hotel and fifteen or twenty bath-houses. These buildings were burned during the Civil War and never rebuilt. In addition to the other improvements a pavilion was built over the spring and this still remains to-day. (See Plate LII). It is thought by some that this is the first place in the United States



PLATE L.—The Black Sulphur Spring at White Sulphur Springs. This lovely pavillon is situated just in the rear of the Greenbrier Hotel. The spring rises in the center, and the waters from the other springs are brought to the dispensing fountain where all who wish may drink of them freely. This is said to be the oldest spring pavillon in the United States, being erected about 1818. Photo. by Cummins.



PLATE LI.—Alvon Springs, spring-house, pump-house, and bottling-house owned by White Sulphur Springs, Inc. This is the source of the water used in the town of White Sulphur Springs. View looking northeast.

is sometimes oolitic. It sometimes contains scattered nodules of dark chert. Several quarries have been opened in this member.

H. B. and H. N. Fullen Quarry No. 1—No. 7 on Map II.

On west side of U. S. Route 219, 0.25 mile south of Lewisburg; mild dip to northwest; Sinks Grove; elevation, 2200'.

	Thickness. Total.	
	Feet.	Feet.
1. Limestone, dark, hard.....	10	10
2. Limestone, dark, hard.....	28	38

Two samples (Nos. 187-PH and 186-PH) were taken from Nos. 1 and 2 of the above section and the results of chemical analysis are published in the Table of Limestone Analyses on page 632. There is a small crushing and screening plant at the quarry.

H. B. and H. N. Fullen Limestone Quarry No. 2—No. 8 on Map II.

On Wade farm, 0.2 mile east of U. S. Route 219, 2.4 miles northeast of Lewisburg and 0.95 mile west of Edgewood School; dip, 5° west northwest; Sinks Grove; elevation, 2200'±.

	Feet
Limestone, dark-gray, brittle.....	15

A sample (No. 164-PH) was collected from the above quarry and the results of the chemical analysis are published in the Table of Limestone Analyses on page 632.

The State Department of Mines reports a limestone quarry operated by Mr. S. O. Collison of Lewisburg. In reply to an inquiry, Mr. Collison reports that his quarry is located just west of the city limits of Lewisburg, on the north side of the Midland Trail (U. S. Route 60). The quarry is probably in the Patton or Sinks Grove.

HILLSDALE LIMESTONE.

The Hillsdale Limestone, previously described on page 279, is of minor economic importance in Greenbrier County. As a rule it contains too much chert to be used for most purposes and the nodules are hard on crushing machinery. Occasionally, however, the chert may be largely absent and the limestone appears to be of high quality.

One abandoned quarry (**No. 12 on Map II**) was noted in this member. It is located along the Midland Trail (U. S. Route 60) 0.8 mile west of Alta. The crushed limestone was probably used for road material.

LIMESTONES OF THE DEVONIAN AND SILURIAN PERIODS.

GENERAL STATEMENT.

The older limestones of Greenbrier County, including the Lower Devonian and Upper Silurian limestones, are of much less extent and commercial value than are those of the Mississippian. Their outcrops are confined to the Coles-Beaver Lick Mountain area as shown on Figures 14 and 15.

In the Devonian, some of the limestone of the Helderberg, particularly the Keyser Member, is fairly high in lime content. In the Silurian there are a few beds in the Salina and Niagara Series that also seem to be fairly pure. Because of the general inaccessibility of these beds and because of the vast supply of limestone from the Greenbrier Series, it is doubtful if they will ever be of more than local value.

HELDERBERG LIMESTONE.

The limestone of possible commercial value in the Helderberg in Greenbrier County is confined to the Beeraft and Keyser Members. The limestone in the Beeraft is high in silica due to sand, silicified fossils, and chert, so that it will probably not be used for anything but local use. Some of the beds of the Keyser appear to be fairly pure but the better limestones of the Greenbrier Series and Silurian minimize the importance of this source of lime. The description of these limestones is published on pages 323-5, and a few analyses are published in the Table of Limestone Analyses, page 632.

SALINA SERIES.

The Salina Series, composed of the Bossardville and Rondont Groups, was described on pages 331-2, where it was pointed out that their outcrop is generally inaccessible except for the region just west and north of Alvon. A few

of the beds appear to be fairly pure, one of the chemical analyses showing 93.4 per cent. calcium carbonate. Three analyses from rocks of this series are published in the Table of Limestone Analyses, page 632.

NIAGARA SERIES.

The Niagara Limestone, previously described on pages 332-4, is the only one of these lower limestones that is being used commercially at the present time. The C. C. C. workers have opened a quarry in the Anthony Creek gorge just west of Alvon and are using the limestone for bridge abutments and construction work. As shown in the Table of Limestone Analyses, one sample was collected from this limestone. As seen from the analysis the limestone is fairly pure but due to its small area of outcrop and its general inaccessibility, it will not be of more than local importance.

TABLE OF LIMESTONE ANALYSES.

The following table gives a summary of the results of the chemical tests made on the limestones of Greenbrier County. The samples were collected by Price and others and the analyses were made in the Survey Laboratory by Mr. Homer A. Hoskins, Chemist. No attempt was made to collect complete sets of samples except at a few of the quarries but samples were obtained from the various calcareous members that appear to have commercial value. Following the table are brief references to the location and portion of the formations sampled. The sample numbers are the same as those mentioned in the foregoing text:

Table of Limestone Analyses, Greenbrier County.

Sample No.	Name of Limestone	Silica (SiO ₂)	Ferrie Iron (Fe ₂ O ₃)	Alumina (Al ₂ O ₃)	Manganese Dioxide (MnO ₂)	Calcium Oxide (CaO)	* Calcium Carbonate (CaCO ₃)	Magnesium Oxide (MgO)	* Magnesium Carbonate (MgCO ₃)	Potash (K ₂ O)	Soda (Na ₂ O)	Titanium Oxide (TiO ₂)	Phosphoric Acid (P ₂ O ₅)	Moisture	Not Determined	* Loss on Ignition	Total
185-PH	Avila	14.72	2.73	3.07	Tr.	39.25	70.00	2.22	4.64	0.024	0.23	3.10	83.70	100.00
169-PH	Girray	9.00	(1.81)	Tr.	47.35	84.51	3.69	7.72	0.06	0.13	39.32	101.36
184-PH	Alderson	9.25	1.40	2.35	Tr.	46.72	83.38	1.66	3.84	0.03	0.24	0.15	38.20	100.00
183-PH	Union	2.35	0.51	0.57	Tr.	52.38	93.48	1.22	2.57	0.01	0.28	0.37	42.31	100.00
182-PH	Union	1.30	0.04	0.48	54.30	96.91	0.90	1.87	0.006	0.15	43.19	100.37
181-PH	Union	3.25	0.47	0.81	Tr.	50.23	89.65	2.74	5.74	0.02	0.20	0.39	41.80	100.00
180-PH	Union	16.10	1.44	3.92	Tr.	34.20	61.04	8.22	17.18	0.02	6.71	36.51	101.12
179-PH	Union	5.75	0.41	1.26	Tr.	49.15	87.72	2.21	4.62	0.02	0.16	0.22	40.82	100.00
178-PH	Union	4.40	0.51	0.99	Tr.	50.50	90.13	2.01	4.21	0.01	0.07	0.20	41.31	100.00
177-PH	Union	15.09	2.05	2.42	Tr.	37.94	67.71	6.13	12.87	0.03	0.14	0.36	35.84	100.00
176-PH	Union	8.62	1.90	1.54	Tr.	45.93	81.97	2.03	4.60	0.04	0.00	0.41	38.54	100.00
175-PH	Union	7.90	2.03	0.09	Tr.	46.38	82.77	2.69	5.63	0.04	0.19	0.87	30.03	100.00
168-PH	Union	7.12	(2.46)	44.87	80.00	4.45	9.30	Tr.	0.04	0.18	1.28	39.60	100.00
170-PH	Union	3.50	(1.10)	Tr.	50.78	90.63	2.53	5.29	0.04	0.11	0.13	41.81	100.00
Average	Union	6.85	(2.35)	Tr.	46.42	82.85	3.11	6.43	0.26	0.10	0.68	30.44	100.20
174-PH	Pickaway	5.52	1.08	0.93	Tr.	50.08	89.38	1.24	2.59	0.008	0.18	0.74	40.22	100.00
173-PH	Pickaway	32.40	2.31	0.67	Some	25.90	46.38	4.04	10.32	Some	0.20	2.25	25.24	100.00
172-PH	Pickaway	17.08	2.34	0.11	Some	37.06	66.14	2.78	5.82	Some	0.26	1.82	32.55	100.00
171-PH	Pickaway	31.23	2.10	5.28	Some	29.53	52.69	3.08	6.44	Some	0.29	1.60	26.80	100.00
167-PH	Pickaway	26.83	(8.41)	31.90	57.11	2.22	4.64	Tr.	Tr.	0.07	0.26	2.79	27.43	100.00
Average	Pickaway	22.61	(3.01)	34.93	62.34	2.85	5.06	0.016	0.24	1.86	30.45	100.00
00P-H	Taggard	12.54	1.27	2.12	Tr.	45.10	80.65	1.83	3.81	Tr.	Tr.	Tr.	Tr.	37.29	100.24
165-PH	Taggard	0.45	(1.71)	47.81	85.50	1.40	2.94	0.30	0.07	0.51	38.75	100.00
Average	Taggard	21.09	(1.70)	46.50	83.08	1.61	3.38	0.15	0.07	0.51	38.02	100.12

Table of Limestone Analyses, Greenbrier County—(Continued).

Sample No.	Name of Limestone	Silica (SiO ₂)	Ferrie Iron (Fe ₂ O ₃)	Alumina (Al ₂ O ₃)	Manganese Dioxide (MnO ₂)	Calcium Oxide (CaO)	Calcium Carbonate (CaCO ₃)	Magnesium Oxide (MgO)	Magnesium Carbonate (MgCO ₃)	Potash (K ₂ O)	Soda (Na ₂ O)	Titanium Oxide (TiO ₂)	Phosphoric Acid (P ₂ O ₅)	Moisture	Not Determined	** Loss on Ignition	Total
Gr. 8-9	Patton	33.47	2.28	5.46	0.015	20.15	52.02	1.06	4.69	Tr.	Tr.	Tr.	None	0.12	3.15	24.40	100.00
Gr. 8-8	Patton	3.00	0.04	0.15	0.008	53.02	94.02	1.30	2.71	Tr.	Tr.	Tr.	None	0.08	0.30	42.35	100.47
Gr. 3-7	Patton	5.72	0.73	1.16	0.01	60.32	89.80	0.92	1.02	Tr.	Tr.	Tr.	None	0.03	0.07	40.07	100.00
Gr. 3-6	Patton	8.17	0.03	1.53	0.01	47.40	85.75	2.38	4.98	Tr.	Tr.	Tr.	None	0.02	0.70	39.34	100.00
Gr. 8-5	Patton	36.54	0.81	2.50	0.01	30.50	54.44	2.39	4.99	Tr.	Tr.	Tr.	None	0.04	0.08	26.44	100.00
Gr. 3-4	Patton	6.11	0.65	1.38	0.01	47.05	86.57	3.15	6.60	Tr.	Tr.	Tr.	None	0.04	0.08	40.63	100.00
Gr. 8-3	Patton	5.14	0.72	1.56	0.01	38.40	88.50	1.89	24.38	Tr.	Tr.	Tr.	None	0.03	0.08	42.70	100.51
Gr. 8-2	Patton	3.02	0.33	0.46	0.02	51.81	92.50	1.87	3.91	Tr.	Tr.	Tr.	None	0.03	0.08	42.56	100.07
Gr. 3-1	Patton	2.00	0.37	0.40	Tr.	52.07	94.00	1.21	2.52	Tr.	Tr.	Tr.	None	0.03	0.08	42.38	100.11
Average Patton		11.57	0.83	1.63	0.010	44.59	70.80	3.60	0.20	Tr.	Tr.	Tr.	0.03	0.058	0.89	37.05	100.13
187-PH Sinks Grove		4.03	0.25	0.66	Pr.	53.35	95.21	0.85	1.51	Tr.	Tr.	Tr.	0.03	0.18	2.36	42.48	102.73
136-PH Sinks Grove		2.48	0.61	0.84	Pr.	51.22	91.41	1.25	2.66	Tr.	Tr.	Tr.	0.01	0.13	0.42	41.05	100.00
166-PH Sinks Grove		6.10	(2.10)	(1.00)	Tr.	40.74	88.77	1.27	2.66	Tr.	Tr.	Tr.	0.03	0.08	0.18	30.89	100.00
164-PH Sinks Grove		13.67	(1.00)	(2.73)	Pr.	45.20	80.07	1.03	4.04	Tr.	Tr.	Tr.	0.09	0.14	0.08	40.15	100.68
Average Sinks Grove		0.81	0.34	0.37	Tr.	49.88	80.01	1.07	2.70	Tr.	Tr.	Tr.	0.04	0.146	0.08	20.72	100.43
101-PH Becraft		51.70	0.82	0.82	Tr.	26.78	46.02	0.45	0.93	Tr.	Tr.	Tr.	Tr.	0.08	0.08	28.57	101.33
103-PH Keyser		32.01	0.82	2.32	Tr.	34.60	61.86	1.26	2.63	0.55	0.45	Tr.	0.09	0.08	0.08	42.46	100.51
109-PH Keyser		4.55	0.90	0.60	Tr.	40.05	80.15	1.00	3.91	Tr.	Tr.	Tr.	Tr.	0.13	0.13	30.97	100.46
116-PH Keyser		6.44	0.00	0.02	Tr.	45.20	80.82	4.75	9.04	0.36	0.63	Tr.	0.07	0.10	0.08	37.00	100.60
Average Keyser		14.53	0.87	1.28	Tr.	43.30	77.27	2.63	5.49	Tr.	Tr.	Tr.	0.05	0.10	0.08	42.21	102.05
111-PH Bossardville?		8.54	0.90	0.08	Tr.	37.81	97.48	11.49	24.63	Tr.	Tr.	Tr.	0.02	0.05	0.05	43.63	100.08
112-PH Rondout		1.92	0.45	0.22	Tr.	52.35	93.43	2.34	4.89	Tr.	Tr.	Tr.	0.02	0.12	0.08	42.07	100.01
113-PH Rondout		2.79	0.35	0.50	Tr.	40.66	88.63	2.50	5.23	Tr.	Tr.	Tr.	0.03	0.08	0.08	45.37	103.11
114-PH Niagara		4.22	0.53	0.71	Tr.	47.51	87.70	4.68	0.77	Tr.	Tr.	Tr.	0.02	0.08	0.08	45.37	103.11

*Not included in total.

**Mainly CO₂

- 185-PH. Avis Limestone, collected from Quarry No. 1 on Map II, 2 miles southeast of Rupert.
- 169-PH. Glenray Limestone, outcrop sample collected along Midland Trail (U. S. Route 60), 2 miles west of Alta, elevation, 2250' B.
- 184-PH. Alderson Limestone, Acme Limestone Company Quarry—No. 5 on Map II, one mile west of Fort Spring; see section, page 622.
- 183-PH. Union Limestone, Acme Quarry; see 184-PH.
- 182-PH. Union Limestone, Acme Quarry; see 184-PH.
- 181-PH. Union Limestone, Acme Quarry; see 184-PH.
- 180-PH. Union Limestone, Acme Quarry; see 184-PH.
- 179-PH. Union Limestone, Acme Quarry; see 184-PH.
- 178-PH. Union Limestone, Acme Quarry; see 184-PH.
- 177-PH. Union Limestone, Acme Quarry; see 184-PH.
- 176-PH. Union Limestone, Acme Quarry; see 184-PH.
- 175-PH. Union Limestone, Acme Quarry; see 184-PH.
- 168-PH. Union Limestone, outcrop sample, along Midland Trail (U. S. Route 60), at Richlands.
- 170-PH. Union Limestone, outcrop sample, along Midland Trail, 2 miles west of Alta.
- 174-PH. Pickaway Limestone, Acme Quarry; see 184-PH.
- 173-PH. Pickaway Limestone, Acme Quarry; see 184-PH.
- 172-PH. Pickaway Limestone, Acme Quarry; see 184-PH.
- 171-PH. Pickaway Limestone, Acme Quarry; see 184-PH.
- 167-PH. Pickaway Limestone, outcrop sample of jointed member, 5 feet thick, along Midland Trail (U. S. Route 60) 0.7 mile northwest of city limits of Lewisburg, elevation, 2140' B.
- 99-PH. Taggard Limestone, outcrop sample, 5 feet thick, 0.7 mile northwest of Renick, elevation, 2160' B.
- 165-PH. Taggard Limestone, outcrop sample, along Seneca Trail (U. S. Route 219) at bridge crossing Spring Creek.
- Gr 8-9. Patton Limestone, Renick Stone Company Quarry—No. 11 on Map II, one mile east of Renick P. O.; see section, page 627.
- Gr 8-8. Patton Limestone, Renick Quarry; see Gr 8-9.
- Gr 8-7. Patton Limestone, Renick Quarry; see Gr 8-9.
- Gr 8-6. Patton Limestone, Renick Quarry; see Gr 8-9.
- Gr 8-5. Patton Limestone, Renick Quarry; see Gr 8-9.
- Gr 8-4. Patton Limestone, Renick Quarry; see Gr 8-9.
- Gr 8-3. Patton Limestone, Renick Quarry; see Gr 8-9.
- Gr 8-2. Patton Limestone, Renick Quarry; see Gr 8-9.
- Gr 8-1. Patton Limestone, Renick Quarry; see Gr 8-9.
- 187-PH. Sinks Grove Limestone, H. B. and H. N. Fullen Quarry No. 1—No. 7 on Map II, 0.25 mile south of city limits of Lewisburg, see section, page 628.
- 186-PH. Sinks Grove Limestone, Fullen Quarry, see 187-PH.
- 166-PH. Sinks Grove Limestone, dark-gray, hard, upper 50 feet sampled, outcrop sample, along Midland Trail at eastern city limits of Lewisburg.

- 164-PH. Sinks Grove Limestone, H. B. and H. N. Fullen Quarry No. 2—No. 8 on Map II, 2.45 miles northeast of Lewisburg; dark brittle limestone 10-15 feet thick.
- 101-PH. Beeraft Member, outcrop sample, on north side of Anthony Creek, 0.5 mile west of Alvon, thickness represented, 15 feet, at top of member.
- 108-PH. Keyser Member, outcrop sample, on north side of Anthony Creek, 0.5 mile west of Alvon, thickness represented, 20 feet; 130 feet below 101-PH.
- 109-PH. Keyser Member, outcrop sample, on north side of Anthony Creek, 0.5 mile west of Alvon, thickness represented, 40 feet; just below 108-PH.
- 110-PH. Keyser Member, outcrop sample, immediately below 109-PH; thickness sampled, 20 feet.
- 111-PH. Bossardville Group (?), outcrop sample; 210 feet below 110-PH.; thickness represented, 90 feet.
- 112-PH. Rondout Group, outcrop sample; just below 111-PH; thickness represented, 55 feet.
- 113-PH. Rondout Group, outcrop sample; just below 112-PH; thickness represented, 20 feet.
- 114-PH. Niagara Series, outcrop sample; just below 113-PH; thickness represented, 80 feet.

ROAD MATERIAL.

Limestone.—Probably the best local material for road building is limestone. As already pointed out vast deposits of limestone are available and it is often found outcropping along the roads, so that almost any amount needed can be secured close at hand or with very little distance of transportation. The distribution and suitability of the various limestones have been discussed in the preceding section of this chapter.

Chert.—For material to improve secondary roads in that part of the county east of the Greenbrier River and north of White Sulphur Springs, the value of the limestones is overshadowed by the presence of large deposits of chert fragments. The Huntersville Chert weathers in such a manner that it can be used on the roads without further treatment and large deposits are present that can be worked by steam shovels. These deposits usually contain enough fine material to serve as a natural binder under the weight of traffic. The surface of such a road may be kept smooth by periodic scraping.

River and Creek Gravel.—Many of the rivers and larger creeks contain large amounts of gravel and afford a cheap supply of material for road improvement. This gravel may be

used particularly to improve muddy roads of secondary importance, where paved roads would be too expensive to maintain. Usually a good grade of gravel can be secured for aggregate for concrete paving, bridge abutments, and concrete in general.

Sand.—Sand, which is an important item in road building both for masonry and concrete, can generally be found along the rivers and creeks, being derived from the weathering of the various sandstones. Sand of better quality can be secured by crushing it from the sandstones but it is usually more expensive. Some of the sandstones, particularly those of the Pottsville and Maneh Chunk, are so situated at their outcrops that weathering has produced large quantities of loose sand.

In addition to these materials there are numerous sandstones as well as arenaceous shales that may often be used advantageously on local roads to improve their condition.

BUILDING STONE.

The sandstones of the county, as described in Part II of this report, vary from thin flaggy and shaly beds that are of no value as building stone to massive ledges 50 to 75 feet in thickness that can be worked into any desired shape. In the Pottsville Series there are several coarse, gray to white sandstones that can be used locally for dimension stone as the needs arise. In the Maneh Chunk Series many of the sandstones are often shaly and lenticular, while others are of massive and durable character with a pleasing texture. In the Greenbrier Series there are no sandstones suitable for building stone but some of the limestones might be successfully used for such purposes. The Maccrady Series offers no stone durable enough for construction material, but the Broad Ford Sandstone member of the underlying Pocono often attains a character suitable for dimension stone. As previously noted it has been quarried quite extensively at many points along the Greenbrier River for use in bridge abutments, building foundations and steps, where durability and abrasive resistance are important.

In the Devonian Period, the Chemung and Portage Series contain sandstones that are generally flaggy but often attain beds of considerable thickness. These beds weather out, break-

ing along the joint-planes into rectangular shapes of various sizes and with very smooth faces, so that further shaping is unnecessary. The colors vary from gray to brown to green and buff. That a market could be found for these flags is quite likely since structures built from them are not only pleasing in appearance but very durable. Universities of central New York have constructed some of their finest buildings from stone of similar character.

The Genesee, Hamilton, and Marellus Series are quite devoid of any rocks suitable for building stone in this county. The Oriskany is often massive and persistent but in this area it is generally unfit for masonry.

In the Silurian Period there are heavy sandstones in the Clinton Series, two of which are quartzitic and very durable but of such a character as to be very difficult to work, while a third, or "Iron Sandstone", is of a red color, very durable and often weathers into rectangular blocks so that further shaping is seldom necessary. Where these beds are not already broken by weathering, it is very difficult to shape them. The White Medina Sandstone is massive and generally quartzitic, like those of the Clinton, and it is very difficult to work into desirable shapes. In the Red Medina the sandstones are generally too shaly and irregular to be of any value.

CLAY.

GENERAL STATEMENT.

Clay, according to Ries¹, is an earthy substance of fine texture containing a mixture of hydrous aluminum silicates, with fragments of other minerals such as silicates, oxides, carbonates, etc., and colloidal material which may be of either organic or mineral character. The mass possesses plasticity (usually) when wet and becomes rock-hard when fired to at least a temperature of redness. The two most important classes of clays are residual and transported.

¹Ries, H., Economic Geology, 5th edition, p. 170; 1925.

AVAILABLE CLAY AND SHALE.

RESIDUAL CLAY.

Residual clay is a type which was derived from the decomposition of the parent rock and which now remains where it was formed. Furthermore the most important deposits are formed from crystalline rocks although similar clay may be formed from stratified beds. So far as known no crystalline rocks occur in Greenbrier County and hence there are no clays from such an origin but occasional clay beds are found in this region at localities where decomposition of the stratified rocks has been sufficient to produce a clay which is residual and which has not been carried off by erosion. As a matter of fact all the rocks contain a certain amount of clay but in most cases it is only a thin veneer and is now better suited for soils than for ceramic use. The limestones, however, often leave a residual clay of varying thickness composed of the insoluble argillaceous impurities of the original formation. Such deposits can be found along the present outcrops of the limestone series where the topography is such that the decomposed product is not readily carried away by surface drainage.

In using a residual clay formed from decomposed limestone it is well to keep in mind that fragments of the limestone are quite injurious if not removed because when burned the limestone tends to slake and form a cavity of weakness and a white blotch on the finished product.

TRANSPORTED CLAY AND CONSOLIDATED CLAY OR SHALE.

Along the river valleys there are many points that retain considerable deposits of river clay which were derived from the decomposition of the rocks over which these streams flowed. These clays are suitable for the manufacture of brick or drainage tile, although the product might not compare favorably with the results from the original material as the sorting is often less complete. These deposits are included under Alluvium and are noted on Map II.

The consolidated clays or shales, composed principally of silica and alumina, with varying quantities of ferric iron and other minor impurities and having sufficient plasticity for molding, occur in large quantities over the county. Throughout

the Mauch Chunk Series, described in detail in Chapter VII on stratigraphy of the series and shown by outcrop on Map II, there are vast quantities of red shale suitable for building brick or drainage tile. Because of the generally high ferric iron content the finished product would have a pleasing red color without the need of adding a flux.

Subsequent to the completion of the field work for this report, a sample of shale was collected from the Mauch Chunk. The test results, as reported by Mr. John P. Nolting, Jr., are as follows:

Report on East Rainelle Brick & Tile Co. Sample.

This sample, composed chiefly of red shale, but including a small amount of yellowish shale, was collected from the Blue-stone Group of the Mauch Chunk Series, about one-fourth mile east of East Rainelle, W. Va., along Route W. Va.-U. S. 60.

For test purposes, this shale was ground to pass through a 40-mesh sieve, mixed with water and passed through a pug mill a number of times. It was finally formed into bars about 1 inch in cross section. Part of these bars were cut into briquettes about 2 inches in length, and part into test bars about 8 inches in length.

The briquettes were then fired, part to cone 015 (770° C.), part to cone 05 (1030° C.) and the remainder to cone 5 (1180° C.). After firing, various tests were run on them, the results of which are shown on the accompanying sheet of "Average Characteristics".

All of the bars were fired to cone 5 (1180° C.). They were then measured for shrinkage and tested to determine the Modulus of Rupture, the results being shown on the accompanying sheet of "Average Characteristics".

As a result of these tests, it is apparent that the clay should be suitable for drain tile when fired to cone 016; for building brick when fired to cone 05 to 02; and for paving brick when fired to cone 5 to 6.

On the basis of the Modulus of Rupture, the test bars would be classified as grade "A" building brick.

The accompanying briquettes show the physical character of the material when fired to different temperatures:

Briquette Number	Conc Number	Degrees Centigrade	Degrees Fahrenheit
7	015	770	1418
16	.05	1030	1886
23	5	1180	2156

Average Characteristics.

Data from Briquettes:

Water of Plasticity (based on 3 samples).....	23.31%
Shrinkage Water (based on 3 samples).....	8.95
Pore Water (based on 3 samples).....	14.36
Volume Drying Shrinkage (based on 3 samples).....	17.24
Linear Drying Shrinkage (based on 3 samples).....	6.12
Apparent Porosity of Fired Piece:	
Fired to 770° C. (8 samples).....	28.38%
Fired to 1030° C. (8 samples).....	17.64
Fired to 1180° C. (8 Samples).....	1.06
Volume Firing Shrinkage:	
Fired to 770° C. (7 samples).....	+1.73%
Fired to 1030° C. (8 samples).....	-1.18
Fired to 1180° C. (8 samples).....	-1.91
Apparent Sp. Grav. of Fired Piece:	
Fired to 770° C. (8 samples).....	2.55
Fired to 1030° C. (8 samples).....	2.54
Fired to 1180° C. (8 samples).....	2.33
Bulk Sp. Grav. of Fired Piece:	
Fired to 770° C. (8 samples).....	1.83
Fired to 1030° C. (8 samples).....	2.09
Fired to 1180° C. (8 samples).....	2.30
Absorption of Fired Piece:	
Fired to 770° C. (8 samples).....	15.55%
Fired to 1030° C. (8 samples).....	8.44
Fired to 1180° C. (8 samples).....	0.46

Data from Bars:

Linear Drying Shrinkage (7 samples).....	5.71%
Linear Firing Shrinkage (7 samples).....	7.29
Modulus of Rupture (7 samples).....	3,144 lbs. per sq. in.

The shales of the Maerady Series are similar to those of the Mauch Chunk and are favorably located along the railroad. As already pointed out, the shales of the Mauch Chunk and Maerady Series could be used with the limestones of the Greenbrier Series to make a mixture suitable for Portland cement or rock-wool.

In the Pocono Series in general the shales are too closely associated with sandstone to offer much inducement to the ceramic industry.

In the Devonian Period the shales of the Catskill Series correspond favorably with those of the Mauch Chunk and are located in most cases along the Chesapeake and Ohio Railway so that they are easily available. The shales of the Chemung and Portage Series are interbedded with flaggy sandstones so that they offer little inducement, while the black Genesee and Marcellus Shales, lower down, contain so much organic matter that their shrinkage would be too great.

In the Silurian Period shales occur in the Clinton and Red Medina Series. In some cases the former by careful selection might be successfully used for building brick or tile, but their exposures are generally inaccessible so that the better located deposits would naturally outrank them in importance.

FIRE CLAY.

The true fire clays that have a quality of resisting high furnace temperatures are not known to occur in the county. It is possible that in the western portion of the county some of these clays may be associated with the coals but all clays associated with the coals are not fire clays, so that only further investigation will definitely determine their presence.

GLASS-SAND.

No development of glass-sand has been attempted in Greenbrier County, although there are one or more deposits that deserve detailed investigation. Since silica is the major ingredient of glass-sand, it influences the character of the ware. Sands with impurities, unless they can be easily removed, and especially if they are to be used for the higher grades of glassware, should be avoided. Chemical analyses of most sands show at least traces of iron oxide, alumina, titanium oxide, lime, magnesia, and organic matter, but these are often included in mineral grains separate from the quartz and may be easily removed.

Along with a good sand two other factors are important, one being a favorable quarry site and the other, access to good transportation. These various factors were considered in sampling sandstones for analysis in Greenbrier County. Among the numerous sandstones available only two offer glass-

sand possibilities, these being the Droop and Healing Springs Sandstones. The Oriskany Sandstone, which is quarried extensively in Berkeley County, is generally quite impure in Greenbrier.

The Droop Sandstone that covers several hundred acres on Muddy Creek Mountain meets the general requirements of a glass-sand unless it should be too fine. Unfortunately no screen tests were made but the sand is in general quite fine and might not all be retained in the 120-mesh which is usually demanded.

The Healing Springs Sandstone appears to be sufficiently pure to be considered as a possible glass-sand. Three analyses of samples collected from this horizon are given in the table below. The chief objections to this source of sand are its general inaccessibility and lack of good quarry sites.

As noted in the table one sample of the Keefer Sandstone was analyzed as a possible glass-sand. This rock is probably too quartzitic and its outcrop too inaccessible for economical use.

Table of Sandstone Analyses.

Sample No.	Sandstone	Silica (SiO ₂)	Ferric Iron (Fe ₂ O ₃)	Alumina (Al ₂ O ₃)	Manganese (MnO ₂)	Lime (CaO)	Magnesia (MgO)	Titanium (TiO ₂)	Phosphoric Acid (P ₂ O ₅)	Moisture	Loss on Ignition	Total
104-PH	Healing Springs.....	98.43	0.34	0.23	0.29	0.00	0.20	Trace	0.03	0.47	100.74
107-PH	Healing Springs.....	98.59	0.30	0.38	Trace	0.40	Trace	Trace	0.03	0.31	100.01
127-PH	Healing Springs.....	98.24	0.14	0.13	Trace	0.70	0.02	Trace	Trace	0.02	0.04	99.80
115-PH	Keefer	95.45	0.50	0.80	Trace	1.37	0.75	Trace	Trace	0.08	1.00	100.91

FORESTS.

In Volume V, pages 146-150 of the Survey Reports (1911), Mr. A. B. Brooks, former State Forester, has described briefly the forests and lumber industry of Greenbrier County. The descriptions of present conditions and lumber mills are now out of date but certain items are of much interest and are reprinted, in part, here.

ORIGINAL FOREST CONDITIONS.

The county may be divided into three districts according to the kinds of timber which each produced in greatest abundance. First, in the mountainous section on the east of Greenbrier River, white pine was the most valuable species. It grew in this county most abundantly on Anthony Creek and its tributaries. The following description of the white pine growing in Greenbrier and Pocahontas is given by Mr. Cecil Clay, former president of the St. Lawrence Boom and Manufacturing Company of Ronceverte:

"There are several hundred million feet of good white pine lumber in this district. The white pine growing as it does here at an altitude of 2,000 to 2,500 feet, has a climate about like that of lower Pennsylvania and much likeness to Susquehanna pine. Where the white pine grows it takes the ground to itself, and but little of other timber is found with it. It grows in several localities through the valley (Greenbrier). On Deer and Sitlington Creeks are 100,000,000 feet; on Knapp Creek and branches another 100,000,000 feet; and Splee, Laurel, and Davy Runs, with Anthony Creek, and some outlying patches, would yield a third 100,000,000 feet. This pine timber is perhaps a little heavier than the Pennsylvania pine, but is soft and smooth to work. It is generally a sound, red-knot timber, with remarkably thin sapwood, often averaging not over half an inch in a lot of 1,000 logs. As much as 40,000 feet can sometimes be cut on an acre."*

The timber of the limestone plateau, before referred to, was distinct from that on the east and north. Mr. W. A. Mastin, of White Sulphur Springs, describes the limestone area and its timber as follows:

"The eastern boundary line of the principal limestone area is, of course, the Greenbrier River as far down as Caldwell. Here the river turns more to the west passing out through the limestone and leaving an area of considerable size on its east and south. The western boundary line of the area begins at Alderson, passes up Muddy Creek,

*"Resources of West Virginia"—Maurv and Fontaine.

limestone for use in the chemical processing of wood for paper. In addition to the uses mentioned above, chemical analyses indicate that the limestone of the county is suitable for the manufacture of rock-wool, Portland cement and some of it is sufficiently pure to be used in making glass, steel, etc. Some of the limestone has a pleasing color and a texture suitable for building stone.

The commercially important limestones are those of the Greenbrier Series but the limestones of the overlying Mauch Chunk and the much lower, Lower Devonian and upper Silurian have some commercial possibilities. In quantity the supply of limestone in Greenbrier County might be roughly spoken of as unlimited. Located, as it is, with ready access to railroads and highways, it is probable that industries based upon this resource will be greatly expanded in the future. The outcrop of the rocks of the Greenbrier Series, which are the most important, is shown on Figure 10, page 269, and on Map II in greater detail.

LIMESTONES OF THE MISSISSIPPIAN PERIOD.

In the Mississippian Period there are numerous limestones of a variety to suit every purpose. In the Mauch Chunk Series the Avis, Reynolds, and Glenray Limestones are generally too impure for many uses but they are suitable for road material or manufacture of rock-wool and cement. In the Greenbrier Series, limestone of almost any degree of purity can be found. The Macerady and Pocono Series as herein delimited are devoid of true limestone deposits although some of the shales and sandstones are calcareous.

LIMESTONES OF THE MAUCH CHUNK SERIES. AVIS LIMESTONE.

The Avis Limestone of the Hinton Group, previously described on page 261, is the youngest and therefore the highest calcareous formation with economic possibilities in Greenbrier County. It is generally composed of fairly good limestone that is steel-gray in color and is sometimes separated into two benches with a thin calcareous shale between. It varies in thickness from 10 to 30 feet.

This limestone has been quarried 2.15 miles southeast of Rupert along the Midland Trail (U. S. Route 60). The limestone from this quarry (No. 1 on Map II) was crushed and used in paving the road mentioned. The quarry is abandoned but a sample (No. 185-PH) was collected from the least weathered face. The results of the chemical analysis are shown in the Table of Limestone Analyses, page 631. Judging from this analysis the rock would be well suited to the manufacture of rock-wool or Portland cement but from the way the rock slakes on weathering it is doubtful if it would be suitable for concrete aggregate. The limestone is too high in impurities to be used as a source for lime.

REYNOLDS LIMESTONE.

The Reynolds Limestone of the Bluefield Group, described on page 264 as a blue or yellowish-blue limestone, is generally too high in impurities for most uses. It varies from 15 to 40 feet in thickness but it is doubtful if this rock could be used to a good advantage. Limestone of a similar or better character can generally be found exposed to a better advantage in the underlying Greenbrier Series.

GLENRAY LIMESTONE.

The Glenray Limestone of the Bluefield Group, described on pages 264-5 as being a gray limestone, varies in thickness from 10 to 60 feet. Like the Reynolds, this limestone will probably not be exploited due to the fact that better limestone can usually be found in the near-by outcrops of rocks of the Greenbrier Series. A sample was collected from this horizon and its analysis is published under number 169-PH in the Table of Limestone Analyses, page 631.

LIMESTONES OF THE GREENBRIER SERIES.

The Greenbrier Series, varying in thickness from 475 to 700 feet and composed almost entirely of limestone, offers numerous opportunities for commercial exploitation. As shown by Figure 10 and in more detail by Map II there is a vast area of this series exposed in Greenbrier County. In Chapter VII attention has been called to the difference in physical

features of the respective members of this series, and to the fact that it is often possible to recognize them at widely scattered points by means of their lithology.

In chemical composition the limestones of this series vary from 1.3 to 36 per cent. in silica, from 29 to 97 per cent. in calcium carbonate, and the magnesium carbonate is generally low. It can therefore be seen that these limestones might be used for numerous purposes where calcareous material is desired. Many of the analyses tabulated fall within the range required for the manufacture of Portland cement where a calcium carbonate content of approximately 75 per cent. and a magnesium carbonate content of less than five per cent. is required. From an examination of the Table of Limestone Analyses it can be seen that there are some localities where certain members of the Greenbrier Series are suited for Portland cement without the admixture of other material. At other points it would be necessary to add certain quantities of shale to lessen the lime content, but this material is readily available just above the limestone in the basal Mauch Chunk Series. Many analyses shown in the table indicate that some of the limestone would be suitable for the manufacture of rock-wool. With more and more emphasis being placed on air-conditioning, the manufacture of this ideal insulating material may become a major industry in the near future.

The Chesapeake and Ohio Railway roughly parallels the outcrop of these deposits and there is ample water and labor supply, while vast quantities of coal are available in nearby areas. With such favorable factors it would appear that the further development of these deposits will not long be overlooked.

ALDERSON LIMESTONE.

The Alderson Limestone, coming at the top of the Greenbrier Series and already described on page 271, is a dark-gray, siliceous or shaly limestone, with a thickness ranging between 50 and 150 feet. This member is generally too impure for the many uses that require a high lime content but it would appear that it might be suitable for rock-wool or Portland cement. The more massive beds could also be used where

features of the respective members of this series, and to the fact that it is often possible to recognize them at widely scattered points by means of their lithology.

In chemical composition the limestones of this series vary from 1.3 to 36 per cent. in silica, from 29 to 97 per cent. in calcium carbonate, and the magnesium carbonate is generally low. It can therefore be seen that these limestones might be used for numerous purposes where calcareous material is desired. Many of the analyses tabulated fall within the range required for the manufacture of Portland cement where a calcium carbonate content of approximately 75 per cent. and a magnesium carbonate content of less than five per cent. is required. From an examination of the Table of Limestone Analyses it can be seen that there are some localities where certain members of the Greenbrier Series are suited for Portland cement without the admixture of other material. At other points it would be necessary to add certain quantities of shale to lessen the lime content, but this material is readily available just above the limestone in the basal Manah Chunk Series. Many analyses shown in the table indicate that some of the limestone would be suitable for the manufacture of rock-wool. With more and more emphasis being placed on air conditioning, the manufacture of this ideal insulating material may become a major industry in the near future.

The Chesapeake and Ohio Railway roughly parallels the outcrop of these deposits and there is ample water and labor supply, while vast quantities of coal are available in near-by areas. With such favorable factors it would appear that the further development of these deposits will not long be overlooked.

ALDERSON LIMESTONE.

The Alderson Limestone, coming at the top of the Greenbrier Series and already described on page 271, is a dark-gray, siliceous or shaly limestone, with a thickness ranging between 50 and 150 feet. This member is generally too impure for the many uses that require a high lime content but it would appear that it might be suitable for rock-wool or Portland cement. The more massive beds could also be used where

Brand Limestone Identification Marks Ledges Nos. 8 to 12 incl.
Source Acme Limestone Company, Alderson, W. Va.
Sampled from quarry Quantity Represented unlimited

Test Results

Chemical Analysis	%
Silica and Silicates (Insoluble in HCl).....	12.94
Iron Oxide.....	1.28
Aluminum Oxide.....	0.51
Calcium Carbonate.....	78.13
Magnesium Carbonate.....	6.48

Freezing and Thawing—25 cycles.....O.K.
Toughness (ledge No. 12)..... 8
Abrasion—Per cent. Wear—Deval..... 3.6
Abrasion—Per cent. Wear—Los Angeles..... 11.0

Report on Sample of Rock

Laboratory No. 78236 February 19, 1936
Road (Bluestone Dam) County Summers
Submitted by U. S. Engineer's Office, Huntington, W. Va.
Received February 7, 1936
Brand Calcareous sandstone Identification marks ledge No. 13
Source Acme Limestone Company, Alderson, W. Va.
Sampled from quarry Quantity Represented unlimited

Test Results

Chemical Analysis	%
Silica and Silicates (Insoluble in HCl).....	53.19
Iron Oxide.....	2.24
Aluminum Oxide.....	1.05
Calcium Carbonate.....	34.32
Magnesium Carbonate.....	8.86

Freezing and Thawing—25 cycles.....O.K.
Toughness 16.5
Abrasion—Per cent. Wear—Deval..... 1.9
Abrasion—Per cent. Wear—Los Angeles..... 6.2

Respectfully submitted,

FRED A. DAVIS,
Materials Engineer."

Subsequent to the completion of the field work for this report, a quarry was opened in the upper part of the Union Member, located approximately 3.5 miles west of Lewisburg. The reported location of this quarry was received just before the completion of the drafting work on the geologic map and is shown as quarry No. 5A on Map II. Unfortunately the reported location is in error, the actual location being 0.8 mile northwest of that shown on Map II.

A kiln has been built for use in burning lime from the quarry. Lime is produced for both chemical and agricultural use. It is reported that the Cherry River Paper Company of



PLATE XLVI.—Crushing and screening plant of the Aeme Limestone Company west of Fort Spring. Photo. by courtesy of Aeme Limestone Company.



PLATE XLVII.—Putting off a large shot at the Acme Limestone Company Quarry (No. 5 on Map II) 0.9 mile west of Fort Spring. Photo. by courtesy of Acme Limestone Company.



PLATE XLVIII.—Acme Limestone Company Quarry face (No. 5 on Map II). Note drilling machines and stripping the limestone of soil by hydraulic "gun."



PLATE II.—Small shot in the Frazier Limestone Company Quarry (No. 4 on Map II), 1.4 miles southwest of Fort Spring.

Richwood, Nicholas County, is the largest purchaser of lime from this quarry at the present time. The quarry was visited and sampled in connection with the preparation of a report on the limestone resources of the State.

Lewisburg Limestone Products Company Quarry— No. 5A on Map II.

On Frank Tuckwiller land, just east of Muddy Creek Mountain, 3.5 miles west of Lewisburg, and 0.8 mile northwest of location shown on Map II; face 30 feet by 150 feet and growing west; beds, flat; home office, Richwood, W. Va.; upper Union Member; measured and sampled by John B. Lucke.

Section rewritten in descending order:

	Thickness. Total.	
	Feet.	Feet.
Limestone, impure, very shaly, greenish-gray, not used in quarry operation, sample 7.....	10+	10
Limestone, varies upward from black, coarsely crystalline irregular fracture, very fossiliferous to dark bluish-gray, finely crystalline, less fossiliferous except top foot which is a rich crinoid bed, samples 5 and 6.....	9	19
Limestone, single bed, but varies upward from medium light to medium dark gray, largely oolitic, samples 3 and 4.....	10	29
Limestone, single bed, pure, nearly white, oolitic bed, samples 1 and 2.....	11	40

The chemical analyses of the samples are as follows:

Sample	Lab. No.	CO ₂	SiO ₂	Fe ₂ O ₃	CaCO ₃	MgCO ₃	Total
*1	1211	43.8
*2	1212	43.8
3	1213	43.8
4	1214	43.4
5	1215	40.3
*6	1216	43.6
7	1217	24.9
*Composite analysis	2.23	0.31	95.71	2.15	100.40

Abandoned Limestone Quarry—No. 6 on Map II.

On west side of U. S. Route 219, 2.25 miles northeast of Falling Springs, and 1.8 miles west of Julia; Union; elevation, 2300'

Limestone, thickness undetermined.....

Abandoned Limestone Quarry—No. 10 on Map II.

On west side of U. S. Route 219, 0.9 mile north of Frankford; dip, 10 degrees W. N. W.; **Union**; elevation, 2260'.
Limestone, thickness undetermined.....

PICKAWAY LIMESTONE.

The Pickaway Limestone, already described on pages 272-7, is a blue to yellowish-gray limestone that is high in silica. The peculiar jointing in one of the ledges in this member was described on the pages cited where it was pointed out that the joint filling is considerably higher in impurities than is the rest of the limestone. The more massive beds of this member including the unweathered jointed ledge, would be suitable for road material, concrete aggregate, railroad ballast, etc. The upper part of this member is exposed in the east end of the **Acme Limestone Company Quarry (No. 5 on Map II)**. From the chemical analyses given on page 631, it would appear that this member would be ideal for the manufacture of rock-wool.

TAGGARD LIMESTONE.

The Taggard Limestone, previously described on pages 277-8, was sampled at two localities in the northern part of the County. As shown in the Table of Limestone Analyses, page 631, it is somewhat high in silica. Its characteristics can be duplicated or bettered in other beds of the Greenbrier Series so that its commercial possibilities are small.

PATTON LIMESTONE.

The Patton Limestone, previously described on page 278, is somewhat impure at the top but contains several ledges of very pure limestone. In appearance this member closely resembles the underlying Sinks Grove Limestone. The following quarries were noted as belonging in this member:

Abandoned Limestone Quarry—No. 9 on Map II.

On east side of U. S. Route 219, 1.0 mile south of Frankford; used for road material; **Patton**; elevation, top, 2290' B.
Limestone, dark-gray, hard, fossiliferous.....

Renick Stone Company Limestone Quarry—No. 11 on Map II.

Located 0.9 mile east of Renick P. O., on the Chesapeake and Ohio Railroad; crusher and screening plant; main part of output is used for railroad ballast; plant capacity, 5 to 7 railroad cars a day (cars of 50- to 55-yd. cap.); quarry floor, 100± feet above red Maerady shales; quarry face, 90-160 feet high by 500 feet long; R. B. Holt, lessee; address, Renick, W. Va.; Patton; elevation, 1940' B.

	Feet.
Limestone, dark bluish-gray.....	90 to 160

The following section, measured by John B. Lucke, has been rewritten in descending stratigraphic order. The numbers refer to chemical analyses published on page 632. Samples procured from the quarry floor under direction of the foreman:

	Thickness. Total.	
	Feet.	Feet.
9. Sample of ledge about $\frac{2}{3}$ to top of quarry face. Ledge about 20 feet thick, underlying 2 similar ledges	20	20
Limestone, not sampled.....	30	50
8. Limestone, very hard, finely crystalline, fossiliferous, many silty or thin green shale breaks.....	7	57
7. Limestone, dark-gray, silty, very massive, dense, stylolite, fossiliferous in nodules or reefs.....	8	65
6. Limestone, very dark gray to black, finely crystalline, fossiliferous, very hard, brittle, stylolite.....	20	85
5. Limestone, massive, sandy, banded light to medium gray to brown, no fossils.....	5	90
4. Limestone, very fine-grained, light-gray, crypto-crystalline, similar to 3 but slightly darker, black stylolites.....	9	99
3. Limestone, very light gray, crypto-crystalline, dense, smooth, perfect conchoidal fracture, few black stylolites	3	102
2. Limestone, dark-gray, tough, crystalline, fossiliferous	6	108
1. Limestone, best exposed on west end of quarry, blue-gray, massive, finely crystalline, fine fossils, many black stylolites.....	12	120

From the analyses shown in the Table of Limestone Analyses, page 632, the limestone in the above quarry would be suitable for railroad ballast, concrete aggregate, road material and much of it is suitable for the many uses requiring a high lime content.

SINKS GROVE LIMESTONE.

The Sinks Grove Limestone, previously described on pages 278-9 has an appearance that is quite similar to the overlying Patton Limestone. It generally has a very high lime content and

No. 3 Pocahontas Coal, Meadow Bluff District.

The thickness and stratigraphic position of the No. 3 Pocahontas Coal are shown in the Goddard Mountain, Little Clear Creek, Sims Station, Sims School, Big Clear Creek Mountain, Little Sewell Mountain—West Side and Little Sewell Mountain—South End Sections, published in Chapter V. It is also reported in the records of borings Nos. 13, 14, and 15 published in preceding pages of this Chapter.

The description of the prospects and openings noted follows:

On public road between Sims Station and Boggs Knob, 2.9 miles south of Ranelle; used in Sims Station Section; No. 3 Pocahontas Coal?; elevation, 2960' B.

	Ft.	In.
Coal	0	4

W. H. Sims Mine—No. 485 on Map II.

Local mine, on southeast side of Sims Mountain, 2.25 miles south of Ranelle; used in Sims School Section; No. 3 Pocahontas Coal; elevation, 2910' B.

	Ft.	In.
1. Coal, blocky	0'	9"
2. Bone parting	0	2
3. Coal, blocky, laminated	2	0
4. Congealed	2	11

A sample (No. 160PH) was taken from Nos. 1 and 3 of the above section, the analysis of which is published under No. 485 in the Table of Coal Analyses at the end of this Chapter.

The same opening was visited by Ray V. Hennen who gives the following section under No. 574 on page 854 of the Fayette County Report:

	Ft.	In.
Coal, soft (root, sandstone, HARRY)	0'	9"
Bony slate, 0" to	0	2
Coal, soft (slate, black, pavement)	2	6
	3	5

Hennen correlated this seam with the No. 6 Pocahontas but later work apparently proves it to be No. 3 Pocahontas.

Cyrus Goddard Mine—No. 486 on Map II.

Local mine, on west side of Goddard Mountain, 1.9 miles south of East Rahnelle; used in Goddard Mountain Section—West Side; No. 3 Pocahontas Coal; elevation, 2890' B.

1.	Coal (shale roof).....	0'	2"
2.	Shale	0	1
3.	Coal	0	1
4.	Shale	0	5
5.	Coal	0	1½
6.	Shale	0	2
7.	Coal, clean, good (shale floor)	1	5½

A sample (No. 138PH) was taken from No. 7 of the above section, the analysis of which is published under No. 486 in the Table of Coal Analyses at the end of this Chapter.

V. F. Eagle Mine—No. 487 on Map II.

Local mine, on the east side of Goddard Mountain, 2.6 miles south of East Rahnelle and 0.75 mile northwest of Meadowvale School; No. 3 Pocahontas Coal; elevation, 3020' B.

Coal (sandstone roof)	0'	4"
Shale	1	6
Coal	1	8

Coal Opening—No. 488 on Map II.

On west side of Little Sewell Mountain, below public road, 0.9 mile southeast of East Rahnelle; used in Little Sewell Mountain Section—West Side; No. 3 Pocahontas Coal; elevation, 2780' B.

Coal, soft, good (shale roof).....	2	0
------------------------------------	---	---

Meadow River Coal and Land Company Prospect—

No. 489 on Map II.

On the west side of Little Sewell Mountain, 2.9 miles southeast of Rahnelle and 2.3 miles southwest of Rupert; No. 3 Pocahontas Coal; elevation, 2960' B.

In. Ft.

Sandstone, brown, medium-grained.....	0'	8"
Coal, laminated with fusulin	0	5
(mineral charcoal)	0	5
Coal, hard	1	1
Coal, banded	1	1

A sample (No. 137PH) was taken from the above section, the analysis of which is published under No. 489 in the Table of Coal Analyses at the end of this Chapter.

Meadow River Lumber Company (?) Prospect— No. 490 on Map II.

On west side of Little Sewell Mountain, 3 miles southeast of Rainelle and 2.3 miles southwest of Rupert; No. 3 Pocahontas Coal; elevation, 2985' B.

Coal (sandstone roof) reported 3' 0" to.....	3	
	Ft.	In.
		6

Coal Exposure—No. 491 on Map II.

On public road, on south end of Little Sewell Mountain, 2.5 miles south of Rupert; used in Little Sewell Mountain Section—South End; No. 3 Pocahontas Coal; elevation, 3190' B.

Coal, exposed	1	
	Ft.	In.
		0

Coal Exposure—No. 492 on Map II.

On upper side of State highway, 0.4 mile north of Charmco; used in Charmco Section; No. 3 Pocahontas Coal; elevation, 2580' B.

Coal, exposed	0'	1"
Fire clay	1	9
Coal, exposed	0	2
	Ft.	In.
		0

Coal Exposure—No. 493 on Map II.

On public road, on south end of Mill Creek Mountain, 2 miles northwest of Rupert; No. 3 Pocahontas Coal; elevation, 2745' B.

Coal, exposed, 1' 0" to	2	
	Ft.	In.
		0

Coal Exposure—No. 494 on Map II.

On public road, on east side of Big Clear Creek Mountain, 1 mile north of Rupert; used in Big Clear Creek Mountain Section; No. 3 Pocahontas Coal; elevation, 3033' B.

Coal, exposed	1	
	Ft.	In.
		0

Gauley Coal Land Company Prospect 599—No. 495 on Map II.

On west side of Big Clear Creek, 1.7 miles north of Rupert; No. 3 Pocahontas Coal; elevation, 3019' L.

Coal (sandstone roof)	0'	8"
Bone	0	3
Coal (fire clay floor).....	1	10

Gauley Coal Land Company Prospect 598—No. 496 on Map II.

On west side of Big Clear Creek, 1.55 miles north of Rupert; No. 3 Pocahontas Coal; elevation, 2989' L.

Coal (sandstone roof)	0'	6"
Sandstone	3	8
Coal	0	8
Slate	0	3½
Coal	1	9
Slate	0	0½
Coal (fire clay floor).....	0	5

Gauley Coal Land Company Prospect 597A—No. 497 on Map II.

On west side of Big Clear Creek, 3 miles north of Rupert; No. 3 Pocahontas Coal; elevation, 2930' L.

Coal (sandstone roof; fire clay floor).....	3	2
---	---	---

Coal Exposure—No. 498 on Map II.

Along road, on south side of Little Clear Creek Mountain, 1.2 miles south of Anjean; used in Little Clear Creek Section; No. 3 Pocahontas Coal; elevation, 3125' B.

Coal, badly weathered, exposed.....	4	0
-------------------------------------	---	---

No. 3 Pocahontas Coal, Williamsburg District.

The following is the only exposure noted in Williamsburg District:

Coal Exposure—No. 499 on Map II.

On Cold Knob road, 0.8 mile north of Cold Knob and 0.9 mile northeast of Grassy Knob; No. 3 Pocahontas Coal; elevation, 4055' B.

In. Ft.

Coal, thickness not determined.....

Quantity of No. 3 Pocahontas Coal Available.

The following table, giving the estimated tonnage of No. 3 Pocahontas Coal in Greenbrier County, has been computed from planimetric measurement of the outcrop as drawn on work sheets for the area indicated on Figure 23, page 602. A low figure for the average thickness was assumed so that the total would not be too great if local areas prove to be cut out or too thin for mining:

Probable Amount of No. 3 Pocahontas Coal.

District.	Thickness of Coal Assumed. Feet.	Square Miles.	Acres.	Cubic Feet of Coal.	Short Tons of Coal. (2000 lbs.)	Meadow Bluff	Williamsburg	Totals
						20.5	0.4	20.9
				13,120	256	1,143,014,400	22,302,720	1,165,317,120
						45,720,576	892,109	46,612,685

SUMMARY OF AVAILABLE COAL.

On preceding pages of this Chapter there is given at the end of the description of each of the six minable coal beds an estimate of the available tonnage of each by magisterial districts along with the total for the county. The following table, with coals arranged in descending order, gives a summary of these statements:

Summary of Available Coal by Districts in Greenbrier County
(in tons of 2,000 pounds).

Coal Seam.	Mine and Prospect Num- bers shown on Map II and Described in Chapter XI.	Meadow Bluff District.	Williamsburg District.	Falling Springs District.	Totals.
Sewell	10-230	151,602,738	22,693,016	41,483,059	215,778,813
Little Raleigh	231-257	59,102,208	23,417,856	82,520,064
Beckley	258-309	79,174,656	25,871,155	3,345,408	108,391,219
Fire Creek	310-370	312,795,648	49,065,984	81,404,928	443,266,560
No. 6 Pocahontas	408-467	273,654,374	41,148,518	8,921,088	323,723,980
No. 3 Pocahontas	484-499	45,720,576	892,109	46,612,685
Totals	922,050,200	163,088,638	135,154,483	1,220,293,321

The above summary is believed to represent approxi-
mately the amount of minable coal that was available before
commercial operations were begun in the County some thirty
years ago. The table at the beginning of the Chapter shows
that a total of 22,823,238 short tons of coal has been mined
in Greenbrier previous to December 31, 1936. The amount
of coal left in ribs and pillars that will probably never be
recovered may increase this total to 28,000,000 tons in round
figures, which sum should be deducted from the total in the
above summary. The amount of coal available after mak-
ing this deduction is, in round numbers, 1,192,000,000 short
tons, assuming an average recovery of 80 per cent. which
appears conservative under modern mining methods, and the
probable amount of coal that should eventually be recovered
in Greenbrier County, is, in round numbers, 953,600,000 short
tons.

TABLE OF COAL ANALYSES.

On the following pages are published the analyses of coal
samples collected from mines, prospects, and cores in or near
Greenbrier County. With the exception of Nos. C6, C7, C8,

C11, 278, 289, 328, 330, 351, 356, 359, 360, and 455, the analyses were made in the laboratory of the Survey, being mainly the work of Homer A. Hoskins and B. B. Kaplan. The analyses of the samples listed above were reported to the Survey, by the coal operators, and the analytical work, in each case, was done by the Commercial Testing and Engineering Company, of Charleston, W. Va. Seven analyses by the same company have been published on preceding pages.

All analyses made in the Survey laboratories were made in strict accordance with the procedure given in the U. S. Bureau of Mines Technical Paper No. 8, except in one respect, namely, moisture on the ground sample was given off in a Freas Drying Oven at a temperature of 110 degrees Centigrade. In this connection it should be noted that a considerable lapse of time, amounting in some instances to more than a year, occurred between the sampling and analyzing of the coals. In some of the samples there has been an apparent loss of moisture in storage.

Concerning the softening temperature of the ash, Hoskins makes the following statement:

"Coal was ashed and made into cones with a 10% dextrin binder, mounted on aluminum plaques and heated to various temperatures in a Denver Fire Clay Fusion Furnace. The temperatures inside this furnace were obtained by means of a Leeds and Northrup pyrometer which had been recently calibrated.

"The initial softening point was read as the temperature at which the ash cone began to deform whether by bending or sloughing. The fusion point was read as the temperature at which the cone formed a sphere, and the temperature at which this sphere melted and flowed out flat was taken as the melting point of the ash."

In the left-hand column is given the number of the sample corresponding to that shown on Map II and used to designate the mine, opening or core in the description published on a preceding page. The second column from the left gives the laboratory or sample number, with the letters indicating the identity of the collector. In that column PH=Price, R=David B. Reger, and H=Ray V. Hennen. The column headed "Mine, Prospect or Core" is self explanatory. In the fourth column from the right is given the key to the analyst by the following: HAH=Homer A. Hoskins, BBK=B. B. Kaplan, JBK=J. B. Krak, and CTE=Commercial Testing and Engineering Com-

pany, of Charleston, W. Va. Under the heading "Carbon Ratio," the fixed carbon has been calculated on the "moisture—ash free" basis. The formulas used are:

$$\text{Fixed Carbon} = \frac{100 - (\text{moisture} + \text{ash})}{100} \times \text{dry moisture—ash—free fixed carbon.}$$

$$\text{Fixed Carbon} = \frac{100 - (\text{moisture} + 1.1 \times \text{ash})}{100} \times \text{dry mineral matter—free fixed carbon.}$$

The second formula is that proposed by A. C. Fieldner, W. A. Selvig, and W. H. Frederic in, "Classification Chart of Typical Coals of the United States," U. S. Bureau of Mines Report of Investigations 3296, December, 1935. According to the classification chart given in the above-named paper, part of the coal of Greenbrier County would be "Medium-volatile bituminous coal" and part "Low-volatile bituminous coal."

The column on the right gives the page reference to the description of the mine or prospect sampled. All of the Survey samples are "channel cuts" of the mining sections of the seams, unless otherwise described, the usual method being to discard from the samples such slates or other impurities as would be rejected in ordinary commercial shipment.

The following abbreviations were used under "Coal Seam and Name of Mine or Prospect":

Greenbrier Sm. Coal Co.....Greenbrier Smokeless Coal Company.
Imperial Sm. Coal Co.....Imperial Smokeless Coal Company.
N. R. & P. Cons. Coal Co.....New River and Pocahontas Consolidated Coal Company.
Leebie Sm. Coal Co.....Leebie Smokeless Coal Company.
Low Ash Sm. Coal Co.....Low Ash Smokeless Coal Company.
Ganley C. L. Co.....Ganley Coal Land Company.

Table of Coal Analyses.

(Under "Condition of Sample", "AD" = air dried; "AR" = as received; "DB" = dry basis).

No. on Map II.		Sample Number.	Mine Prospect or Core.	Coal Seam and Name of Mine or Prospect.	Condition of Sample.		Ash.	Sulphur.	Calorimeter B. T. U. for 1 lb. of Coal.	Softening Temp. of Ash. Degrees F.			Analyst.	Carbon Ratio.		Page.		
					Moisture Lost on Air Drying.	Moisture.				Volatile Matter.	Fixed Carbon.	Initial Softening.		Fusion.	Melting.		M. & A. Free.	M. & M. M. Free.
Sowell Coal																		
16	1581H	M	Haines (J. C. Dixon)	AD	1.52	0.50	25.14	68.71	5.50	0.62	14,600	2,300	2,065	HAH	73.21	73.05	476
16	1581H	M	Haines (J. C. Dixon)	AR	2.07	24.76	67.66	5.51	0.61	14,350	2,300	2,065	HAH	73.21	73.05	476
18	1591H	M	H. J. & W. A. Fitzhugh	AD	3.48	0.73	24.36	68.34	6.57	0.82	14,390	2,110	2,695	2,772	HAH	73.72	74.25	477
18	1591H	M	H. J. & W. A. Fitzhugh	AR	4.18	28.51	65.97	6.34	9.79	13,886	2,110	2,005	2,772	HAH	73.72	74.25	477
22	901H	M	Greenbrier Sm. Coal Co. No. 1	AD	0.41	26.04	70.82	2.13	0.83	14,977	2,100	BBK	72.67	72.82	479
22	91H	M	Greenbrier Sm. Coal Co. No. 2	AD	0.56	26.14	70.51	2.79	1.03	14,924	2,060	BBK	72.95	73.17	479
26	891H	M	Crichton	AD	0.41	27.27	69.10	3.22	1.24	14,943	2,023	BBK	71.70	71.94	480
26	881H	M	Crichton	AD	0.14	0.63	26.49	70.24	2.64	1.08	14,000	2,023	BBK	72.61	72.81	480
26	81H	M	Imperial Sm. Coal Co. No. 1	AD	0.32	26.77	69.09	3.52	1.06	14,701	2,100	BBK	72.25	72.40	481
31	831H	M	Imperial Sm. Coal Co. No. 1	AD	0.26	0.22	25.05	71.97	2.76	0.80	14,019	1,935	BBK	74.18	74.39	481
31	831H	M	Imperial Sm. Coal Co. No. 1	AD	0.96	0.63	26.38	79.34	2.05	0.84	14,777	2,200	BBK	72.73	72.03	483
33	821H	M	Imperial Sm. Coal Co. No. 2	AD	0.11	0.35	26.70	69.12	3.83	0.86	14,774	2,023	BBK	72.14	72.42	483
34	801H	M	Frances	AD	9.11	0.24	25.97	70.41	3.38	1.19	14,744	2,100	BBK	73.05	73.31	484
38	841H	M	Margarette No. 2	AD	0.18	1.12	25.06	71.48	2.34	0.78	14,744	2,222	BBK	74.04	74.22	485
40	851H	M	Margarette No. 1	AD	0.44	25.13	71.01	2.62	0.80	14,812	2,280	HAH	74.19	74.30	486
43	961H	M	N. R. & P. Cons. Coal Co. No. 3	AD	0.61	25.09	71.03	2.77	1.08	14,874	2,110	BBK	74.06	74.27	487
46	921H	M	N. R. & P. Cons. Coal Co. No. 1	AD	3.23	0.60	24.48	71.19	3.73	1.61	14,800	1,075	2,150	2,459	HAH	74.41	74.70	488
51	1571H	M	Joe Nell	AD	3.81	23.74	69.73	2.72	1.40	14,410	1,975	2,150	2,459	HAH	74.60	74.82	488
51	1571H	M	Joe Nell	AR	0.44	24.68	72.28	2.60	0.90	14,992	2,166	HAH	74.65	74.75	489
54	931H	M	N. R. & P. Cons. Coal Co. No. 4	AD	0.66	26.00	71.30	2.04	0.75	15,173	2,350	HAH	73.28	73.43	489
54	941H	M	N. R. & P. Cons. Coal Co. No. 4	AD	2.15	24.31	70.91	2.63	0.85	14,939	2,286	HAH	74.47	74.68	489
54	951H	M	N. R. & P. Cons. Coal Co. No. 4	AD	1.15	3.28	24.03	70.09	2.60	9.84	14,768	2,286	HAH	74.47	74.68	489
54	951H	M	N. R. & P. Cons. Coal Co. No. 4	AR	0.63	26.36	67.07	6.04	1.40	14,639	2,100	BBK	72.06	72.44	498
95	891H	M	Loeie Sm. Coal Co. No. 2	AD	0.28	0.63	26.36	67.07	6.04	1.40	14,639	2,100	BBK	72.06	72.44	498
143	78A-1-H	M	Clear Creek Coal Co. U. Branch	AD	0.68	20.04	68.57	4.16	0.92	14,435	2,023	BBK	72.02	72.34	509
143	78B-1-H	M	Clear Creek Coal Co. L. Branch	AD	0.07	0.30	25.63	66.45	7.62	9.59	14,196	2,431	BBK	72.17	72.77	509

(Under "Condition of Sample", "AD" = air dried; "AR" = as received; "DB" = dry basis).

No. on Map II.		Sample Number.		Mine Prospect or Core.		Coal Seam and Name of Mine or Prospect.		Condition of Sample.		Proximate.		Ash.		Sulphur.		Calorimeter B. T. U. for 1 lb. of Coal.		Softening Tem- perature of Ash, Degrees F.		Analyst.		Carbon Ratio.		Page.			
								Moisture Lost on Air Drying.		Moisture.		Volatile Matter.		Fixed Carbon.				Initial Softening.		Fusion.		Melting.					
143	97PH	M	Clear Creek Coal Co.	AD	6.61	28.05	67.60	3.74	1.16	14,526	2,656	2,368	2,575	HAH	70.67	76.05	609										
150	79PH	P	Raine Lumber and Coal Co.	AD	6.60	29.14	70.30	2.61	0.96	14,827	2,656	2,160	2,575	BBK	72.96	78.09	611										
151	163PH	M	Raine Lumber and Coal Co.	AD	2.84	25.84	60.62	3.86	1.26	14,016	1,963	2,361	2,435	HAH	72.93	73.22	611										
161	163PH	M	Raine Lumber and Coal Co.	AR	3.56	25.16	67.65	3.69	1.22	14,496	1,903	2,361	2,435	HAH	72.94	73.23	611										
C-6	01470	C	Raine Lumber and Coal Co.	DB	6.73	25.49	69.32	4.51	0.90	14,800	2,126	2,423	2,527	CTE	73.67	74.14	167										
162	162PH	P	Raine Lumber and Coal Co.	AD	2.58	24.86	67.51	4.46	1.14	14,690	2,126	2,423	2,527	HAH	73.11	73.46	512										
152	152PH	P	Raine Lumber and Coal Co.	AR	3.85	27.51	68.48	3.08	1.64	14,570	1,995	2,272	2,515	HAH	71.34	71.57	512										
154	154PH	P	Raine Lumber and Coal Co.	AR	4.77	26.44	65.83	2.06	1.06	14,600	1,995	2,272	2,515	HAH	71.34	71.57	612										
154	154PH	P	Raine Lumber and Coal Co.	DB	25.43	71.27	3.36	3.36	1.25	14,025	2,326	2,326	2,326	CTE	73.72	73.95	169										
C-7	61478	C	Raine Lumber and Coal Co.	DB	25.80	70.55	3.65	3.65	0.96	14,017	2,326	2,326	2,326	CTE	73.72	73.95	169										
C-8	61480	C	Raine Lumber and Coal Co.	AR	6.71	27.63	69.34	2.32	0.84	15,316	2,326	2,326	2,326	BBK	71.61	71.68	613										
157	352R	M	Lennel Helms.	AD	4.44	26.54	70.37	2.16	1.61	15,167	1,935	2,335	2,527	HAH	72.61	72.78	524										
214	155PH	P	Raine Lumber and Coal Co.	AR	5.33	25.36	67.25	2.06	6.09	14,404	1,935	2,335	2,527	HAH	72.62	72.78	524										
216	156PH	P	Raine Lumber and Coal Co.	AD	3.36	27.92	67.94	3.64	0.73	14,300	2,272	2,396	2,659	HAH	76.87	71.14	624										
216	156PH	P	Raine Lumber and Coal Co.	AR	3.84	26.05	67.65	3.52	0.71	13,820	2,272	2,396	2,659	HAH	76.88	71.15	624										
C-11	61477	C	Raine Lumber and Coal Co.	DB	25.57	67.65	6.78	6.78	0.97	14,386	2,272	2,396	2,659	CTE	72.57	73.16	172										
222	370R	P	John A. Baile.	AR	1.66	28.86	68.05	2.09	0.76	15,160	2,264	2,756	2,783	BBK	72.96	76.42	625										
225	163PH	M	Eik Lick Coal Co.	AR	1.01	36.66	64.95	3.44	1.66	15,160	2,264	2,756	2,783	HAH	67.97	68.22	625										
225	211R	M	Eik Lick Coal Co.	AR	0.63	26.06	66.77	5.64	6.69	12,213	2,856	2,856	2,856	BBK	71.24	71.67	625										
230	02PH	M	Preston Clark Mfrs.	AD	1.57	26.09	69.88	3.40	6.08	14,756	2,674	2,236	2,550	BBK	73.16	74.18	628										
Average (Sewell)				AR	0.64	26.09	69.88	4.22	6.95	14,247	2,698	2,498	2,585														
Average (Sewell)				DB	2.79	25.86	60.98	4.56	1.03																		

Table of Coal Analyses—(Continued).

(Under "Condition of Sample", "AD" = air dried; "AR" = as received; "DB" = dry basis).

No. on Map II.	Sample Number.	Mine Prospect or Core.	Coal Seam and Name of Mine or Prospect	Condition of Sample.		Moisture.	Volatile Matter.	Fixed Carbon.	Ash.	Sulphur.	Calorimeter B. T. U. for 1 lb. of Coal.	Softening Temperature of Ash, Degrees F.			Analyst.	Carbon Ratio.		Page.
				Moisture Lost on Air Drying.	Proximate.							Initial Softening.	Fusion.	Melting.		M. & A. Free.	M. & M. Free.	
2558	150PH	M	Beckley Coal	AD	1.97	0.86	23.95	69.79	6.21	1.53	14,718	1,925	2,445	2,527	HAH	74.99	76.19	540
2558	160PH	M	Tuck Brothers	AR	2.01	23.24	68.64	9.11	1.50	14,470	1,926	2,445	2,627	HAH	74.71	76.21	540
278	40679	P	Gauley Coal	DB	25.74	69.43	4.83	1.00	14,473	2,398	CTE	72.96	73.33	544
278	40970	P	Gauley Coal	AR	2.69	25.05	67.66	4.70	0.97	14,084	2,398	CTE	72.96	73.32	544
280	40676	P	Gauley Coal	DB	25.87	62.72	11.41	0.90	CTE	70.80	71.72	548
280	40675	P	Gauley Coal	AR	6.75	24.38	59.12	10.76	0.85	CTE	70.80	71.73	548
			Average (Beckley)	AD	1.67	0.36	23.65	69.79	6.21	1.53	14,718	1,926	2,445	2,627	74.69	76.10
			Average (Beckley)	AR	3.48	24.22	65.11	7.19	1.11	14,277	1,925	2,422	2,627	72.82	73.15
			Average (Beckley)	DB	25.80	69.08	8.12	0.95	14,473	2,398	71.88	72.59
311	144PH	M	Fire Creek Coal	AD	2.19	9.48	21.74	68.93	0.16	9.92	13,869	2,296	2,916	HAH	75.94	79.72	557
311	144PH	M	L. E. McClung	AR	2.66	21.26	67.13	8.95	0.89	13,360	2,295	2,915	HAH	75.95	79.72	557
312	143PH	N	L. E. McClung	AD	3.00	0.48	25.14	71.67	2.71	9.71	15,220	2,244	2,916	HAH	74.03	74.24	557
312	143PH	M	L. E. McClung	AR	3.47	24.38	69.52	2.63	0.69	14,760	2,244	2,915	HAH	74.04	74.24	557
328	40681	P	Gauley Coal	DB	26.93	68.05	5.92	9.08	13,900	2,437	CTE	72.33	72.79	561
328	40681	P	Gauley Coal	AR	0.47	24.35	63.04	5.64	0.64	13,099	2,437	CTE	72.33	72.78	561
330	40680	P	Gauley Coal	DB	25.90	97.48	6.62	0.80	CTE	72.29	72.77	561
330	40689	P	Gauley Coal	AR	2.59	25.23	95.73	9.45	0.87	CTE	72.29	72.78	561
351	40674	P	Gauley Coal	DB	24.13	63.69	12.18	0.59	CTE	72.62	73.54	569
351	40674	P	Gauley Coal	AR	2.95	23.64	92.38	11.93	0.55	CTE	72.62	73.54	569
356	40673	P	Gauley Coal	DB	23.79	67.96	8.26	0.56	13,948	2,897	CTE	74.97	74.74	567
359	40973	P	Gauley Coal	AR	2.79	23.13	69.06	8.92	0.54	13,559	2,897	CTE	74.07	74.74	567
359	40976	P	Gauley Coal	DB	24.95	99.08	5.97	0.58	CTE	73.47	73.94	568

Table of Coal Analyses—(Continued).

(Under "Condition of Sample", "AD" = air dried; "AR" = as received; "DB" = dry basis).

No. on Map II.	Sample Number.	Mine Prospect or Core.	Coal Seam and Name of Mine or Prospect.	Condition of Sample.			Ash.	Sulphur.	Calorimeter B. T. U. for 1 lb. of Coal.	Softening Temperature of Ash, Degrees F.			Analyst.	Carbon Ratio.		Page.			
				Moisture Lost on Air Drying.	Proximate.	Fixed Carbon.				Initial Softening.	Fusion.	Melting.		M. & A. Free.	M. & M. M. Free.				
Fire Creek Coal																			
369	40676	P	Ganley Coal Land Co. No. A209.	AR	2.73	24.27	67.19	5.81	0.56	13,386	2,706	CTE	73.46	73.93	568			
360	40677	P	Ganley Coal Land Co. No. A208.	DB	24.64	66.04	9.32	0.55	12,860	2,796	CTE	72.83	73.58	569			
360	40677	P	Ganley Coal Land Co. No. A208.	AR	3.57	23.76	63.68	8.90	0.53	12,860	2,796	CTE	72.83	73.58	569			
370	162PH	P	Manning Knobs.	AR	2.14	24.42	70.50	2.94	0.90	14,500	2,110	2,314	2,600	HAH	74.27	74.50	571		
			Average (Fire Creek).....	AD	2.00	0.48	23.44	70.15	5.93	0.82	14,535	2,270	2,915	74.90	75.43	
			Average (Fire Creek).....	AR	3.16	23.83	66.20	6.81	0.69	13,706	2,216	2,715	2,600	73.53	74.00	
			Average (Fire Creek).....	DB	24.01	67.05	8.04	0.04	13,731	2,710	72.91	73.50		
Little Fire Creek Coal																			
371	871H	M	Win. Sira.	AD	0.07	0.22	23.18	72.70	3.90	0.75	14,726	2,450	BBK	75.82	75.87	239		
371	026H	M	Win. Sira.	AR	0.38	23.51	74.40	1.62	0.66	15,470	JBR	76.01	76.14	239		
374	134PH	M	Win. Bennett.	AD	3.17	0.69	21.05	69.34	8.02	0.65	14,000	2,417	2,045	HAH	76.71	77.48	240	
374	134PH	M	Win. Bennett.	AR	3.84	20.38	67.14	8.64	0.55	13,556	2,417	2,945	HAH	76.71	77.48	240	
			Average (Little Fire Creek).....	AD	1.62	0.46	22.12	71.01	6.41	0.70	14,868	2,417	2,608	76.27	76.78	
			Average (Little Fire Creek).....	AR	2.11	21.95	70.81	5.13	0.61	14,513	2,417	2,945	76.36	76.76	
No. 7 Pocahontas Coal.																			
381	135PH	M	C. N. Callison.	AD	5.20	1.76	20.73	69.83	7.08	0.63	13,700	1,925	2,100	2,250	HAH	77.11	77.77	574	
381	135PH	M	C. N. Callison.	AR	6.87	19.65	66.20	7.26	0.60	13,000	1,925	2,100	2,250	HAH	77.11	77.77	574	
383	136PH	M	E. H. Callison.	AD	2.15	0.32	20.31	73.37	0.00	1.64	14,653	2,085	2,349	2,478	HAH	78.32	78.82	575	
383	136PH	M	E. H. Callison.	AR	2.46	19.87	71.80	5.87	1.62	14,340	2,085	2,349	2,478	HAH	78.32	78.83	575	
384	146PH	M	Meadow River Coal and Land Co.	AD	3.58	0.95	23.14	69.71	6.20	1.01	14,320	2,587	2,827	HAH	75.08	75.58	575	
384	146PH	M	Meadow River Coal and Land Co.	AR	4.50	22.31	67.21	5.08	0.07	13,816	2,587	2,827	HAH	75.08	75.58	575	
			Average (No. 7 Pocahontas).....	AD	3.64	1.01	21.30	70.07	6.63	1.10	14,227	2,190	2,455	2,362	HAH	76.84	77.40
			Average (No. 7 Pocahontas).....	AR	4.61	20.61	68.40	6.88	1.06	13,710	2,197	2,455	2,362	76.84	77.40

Table of Coal Analyses—(Continued).

(Under "Condition of Sample", "AD" = air dried; "AR" = as received; "DB" = dry basis).

No. on Map II.	Sample Number.	Blue Prospect or Core.	Coal Seam and Name of Mine or Prospect.		Condition of Sample.		Proximate.			Ash.	Sulphur.	Calorimeter B. T. U. for 1 lb. of Coal.	Softening Temp. of Ash. Degrees F.	Analyst.	Carbon Ratio.	Page.	
					Moisture Lost on Air Drying.	Moisture.	Volatile Matter.	Fixed Carbon.					Initial Softening.	Fusion.	Melting.		
No. 6 Pocahontas Coal.																	
411	1321-1	M	Geo. Shaver	AD 4.01	0.17	16.99	76.60	6.18	0.93	14.630	2,214	2,805	2,860	HAH	81.86	82.40	582
411	1321-1	M	Geo. Shaver	AR	4.17	16.51	78.59	6.93	0.89	14.095	2,214	2,805	2,860	HAH	81.86	82.40	582
412	1331-1	M	Bert Hutson	AD 3.18	0.30	18.29	76.07	4.44	0.79	14.970	2,126	2,617	2,794	HAH	80.80	81.18	583
412	1331-1	M	Bert Hutson	AR	3.77	17.65	74.29	4.29	0.76	14.520	2,126	2,617	2,794	HAH	80.83	81.18	583
413	1311-1	M	J. A. & S. J. Woodruff	AD 2.93	0.17	18.79	76.82	4.22	1.21	14.908	2,059	2,539	2,728	HAH	80.34	80.70	583
413	1311-1	M	J. A. & S. J. Woodruff	AR	3.19	18.23	74.57	4.10	1.17	14.480	2,059	2,539	2,728	HAH	80.36	80.71	583
417	1301-1	M	Meadow River Fuel Co.	AD 3.05	0.26	27.48	66.84	5.42	1.25	14.810	2,243	2,575	2,695	HAH	79.87	71.27	584
417	1301-1	M	Meadow River Fuel Co.	AR	3.30	26.64	64.81	5.25	1.26	14.858	2,243	2,575	2,695	HAH	79.87	71.28	584
423	1391-1	M	Meadow River Coal and Land Co.	AD 5.35	0.60	26.10	76.65	2.65	0.55	15.067	2,214	2,935		HAH	79.22	79.44	585
423	1391-1	M	Meadow River Coal and Land Co.	AR	5.52	19.02	72.56	3.47	0.52	14.256	2,214	2,935		HAH	79.22	79.45	585
424	1261-1	M	Meadow River Smokeless Coal Co.	AR	0.32	21.60	74.61	3.47	0.54	15.210				HAH	79.23	79.45	585
425	1491-1	M	Low Ash Sm. Coal Co. (Peck)	AD 1.81	0.36	21.00	72.25	3.39	0.62	15.120	2,134	2,838		HAH	77.55	77.83	586
425	1491-1	M	Low Ash Sm. Coal Co. (Peck)	AR	2.16	23.56	70.25	3.33	0.61	14.850	2,181	2,838		HAH	75.06	75.33	586
426	1451-1	P	Gauley Coal Land Co.	AD 2.19	0.49	22.40	71.86	5.25	0.94	14.830	2,575	2,728	2,816	HAH	76.24	76.60	587
426	1451-1	P	Gauley Coal Land Co.	AR	2.97	21.81	70.07	6.12	0.92	14.460	2,575	2,728	2,816	HAH	76.24	76.66	587
428	1411-1	P	L. E. McClung	AD 2.88	0.20	23.85	74.36	2.07	0.75	15.460	2,431	2,605		HAH	76.11	76.27	588
428	1411-1	P	L. E. McClung	AR	3.07	22.68	72.24	2.01	0.73	15.015	2,431	2,605		HAH	76.11	76.27	588
429	1421-1	M	T. E. & S. T. Jones	AD 8.18	1.11	25.10	69.82	3.97	0.60	14.140	1,914	2,198	2,587	HAH	73.55	73.87	588
429	1421-1	M	T. E. & S. T. Jones	AR	9.20	23.05	64.10	3.65	0.55	12.983	1,914	2,198	2,587	HAH	73.55	73.86	588
430	1481-1	M	Joe Nell	AD 4.06	0.62	22.68	73.10	3.60	0.60	15.126	2,389	2,587	2,761	HAH	76.32	76.61	589
430	1481-1	M	Joe Nell	AR	4.65	21.57	70.13	3.45	0.58	14.512	2,389	2,587	2,761	HAH	76.31	76.60	589
431	1471-1	M	Lester Boyer	AD 2.34	0.39	26.34	69.60	3.67	0.79	15.620	2,276	2,803		HAH	72.55	72.82	589
431	1471-1	M	Lester Boyer	AR	2.72	25.52	67.98	3.58	0.77	15.240	2,276	2,803		HAH	72.55	72.83	589
444	1511-1	P	Leckie Smokeless Coal Company	AD 2.05	0.40	25.18	65.51	8.91	1.36	14.220	2,126	2,635	2,750	HAH	72.24	72.95	594

Table of Coal Analyses—(Continued).

(Under "Condition of Sample", "AD" = air dried; "AR" = as received; "DB" = dry basis).

Mine Prospect or Core.	Coal Seam and Name of Mine or Prospect.	Condition of Sample.		Proximate.			Ash.	Sulphur.	Calorimeter B. T. U. for 1 lb. of Coal.	Softening Temp. of Ash. Degrees F.			Carbon Ratio.		Page.	
		Moisture Lost on Air Drying.	Moisture.	Volatile Matter.	Fixed Carbon.	Initial Softening.				Fusion.	Melting.	Analyst.	M. & A. Free.	M. & M. Free.		
	No. 6 Pocahontas Coal.															
P	Beckle Smokeless Coal Company.....	AR	2.44	24.07	64.16	8.73	1.33	13.936	2,126	2,635	2,750	HAH	72.23	72.94	604	
P	Gauley Coal Land Co. No. A813.....	AR	10.20	25.17	56.43	8.20	0.40	OTF	69.15	69.86	597	
P	Gauley Coal Land Co. No. A813.....	DB	28.03	62.84	9.13	0.44	OTF	69.15	69.86	597	
P	Gauley C. L. C. (Hume Opening).....	AD	2.71	22.36	70.81	6.57	0.74	14.000	2,243	2,966	HAH	76.01	76.54	599	
P	Gauley C. L. C. (Hume Opening).....	AR	2.97	21.74	68.90	6.30	0.72	13.020	2,243	2,964	HAH	76.02	76.65	600	
M	Average (No. 6 Pocahontas).....	AD	3.49	22.64	72.41	4.64	0.86	14.839	2,227	2,695	2,749	HAH	76.24	76.64
M	Average (No. 6 Pocahontas).....	AR	4.06	21.08	69.29	4.67	0.78	14.305	2,227	2,695	2,749	76.56	76.81
	No. 3 Pocahontas Coal.															
M	W. H. Sims.....	AD	2.18	19.10	74.07	6.29	0.66	14.670	2,639	2,936	HAH	79.42	79.96	603	
M	W. H. Sims.....	AR	2.62	18.81	72.41	6.10	0.65	14.380	2,639	2,934	HAH	79.38	79.92	603	
M	Gyus Goddard.....	AD	4.77	0.21	18.70	78.09	0.88	15.457	2,085	2,515	2,717	HAH	80.68	80.93	604	
M	Gyus Goddard.....	AR	4.97	17.81	74.36	2.86	0.84	14.721	2,085	2,515	2,717	HAH	80.68	80.93	604	
M	Meadow River Coal and Land Co.....	AD	3.23	0.36	20.34	72.07	0.80	14.660	2,487	2,965	HAH	77.90	78.60	604	
P	Meadow River Coal and Land Co.....	AR	3.68	10.68	69.74	7.00	0.77	14.086	2,487	2,965	HAH	77.90	78.61	604	
P	Average (No. 3 Pocahontas).....	AD	3.30	0.34	19.41	74.74	0.78	14.806	2,370	2,805	79.36	79.86	
P	Average (No. 3 Pocahontas).....	AR	3.72	18.77	72.17	6.34	0.75	14.390	2,370	2,805	79.35	79.83	
	No. 1 Pocahontas Coal															
P	Ed. Grafton.....	AR	0.22	18.42	71.34	0.68	12.622	2,417	2,803	2,966	HAH	79.46	80.38	250	
	Merrimac? (Pocono) Coal															
P	Shuttle Abbott.....	AR	2.27	9.01	12.82	0.09	BRK	66.40	84.18	290	
M	W. A. Napier.....	AD	0.45	0.82	13.69	63.22	12.10	11.403	2.068	BRK	62.31	84.78	291	

included in average.

Table of Coal Analyses—(Concluded).

(Under "Condition of Sample", "AD" = air dried; "AR" = as received; "DB" = dry basis).

No. on Map II.	Sample Number.	Mine Prospect or Corr.	Coal Seam and Name of Mine or Prospect.		Condition of Sample.			Proximate.			Ash.	Sulphur.	Calorimeter B. T. U. for 1 lb. of Coal.	Initial Softening.	Fusion.	Melting.	Analyst.	Carbon Ratio.		Page.
						Moisture Lost on Air Drying.	Moisture.	Volatile Matter.	Fixed Carbon.									M. & A. Free.	M. & M. M. Free.	
604	771H	M	W. A. Napier.....	Merrimac? (Pocono) Coal	AR	0.88	1.27	13.65	62.61	22.47	12.16	1.1,454	2,068	BRK	82.10	84.60	291
607	981H	P	Near Judytown P. O.....	AD	0.88	0.65	13.58	73.23	12.54	1.60	13,422	2,264	HAI	84.36	86.60	291
607	981H	P	Near Judytown P. O.....	AR	1.62	13.44	72.62	12.42	1.58	13,200	2,264	HAI	84.38	86.62	291
608	102H	P	Floyd Childers.....	AR	0.64	10.79	76.48	12.10	0.40	13,390	2,397	HAI	87.64	88.84	292
			Average (Merrimac?) (Pocono).....	AD	0.67	0.74	13.58	68.23	17.45	6.85	12,413	2,166	83.34	86.22
			Average (Merrimac?) (Pocono).....	AR	1.11	12.63	70.57	15.69	4.74	12,711	2,422	84.71	86.46

CHAPTER XII.

LIMESTONE, ROAD MATERIAL, CLAY, BUILDING STONE, GLASS-SAND, FORESTS, AND SOILS.

LIMESTONE.

GENERAL STATEMENT.

From an economic viewpoint the limestone of Greenbrier County stands out as one of its most important assets. Next to coal and timber which are the greatest sources of revenue, limestone is the most valuable natural commodity produced in the county. The original source of wealth in order of importance was coal, timber, limestone, agricultural soils, mineral springs, water-power, iron ore, and manganese ore. Of these, the coal has been fully discussed in the preceding Chapter. The timber has been mostly removed and the possibilities of reforestation will be discussed later in this Chapter. A survey of the soils of Greenbrier County has been completed through the cooperation of the West Virginia Geological Survey and the Bureau of Chemistry and Soils of the United States Department of Agriculture and a separate report on this subject will be published in the near future. The mineral springs, water-power, iron ore, and manganese ore will be discussed in Chapter XIII.

There are three large limestone quarries located along railroads in Greenbrier County and many other small quarries scattered along the highways. The chief product of these quarries is crushed stone for railroad ballast and for black-top roads. Ground limestone that is suitable for agricultural use and for rock-dusting in coal mines is a by-product at the crushing plants. At least one quarry is now furnishing

Coal Exposure—No. 427 on Map II.

On south side of Meadow Creek, 0.5 mile from mouth; No. 6 Pocahontas Coal, elevation. 2480' B.; section reported by Mr. Wm. McChing.

			Ft.	In.
Coal (shale roof).....	2'	1"		
Parting	0	1		
Coal	1	10	4	0

Greenbrier Fire Creek Coal Company "Midland" Mine— No. 428 on Map II.

Same as Midland New Mine, formerly Midland Smokeless Coal Company; 0.8 mile northwest of Charmco, on north side of Meadow River; No. 6 Pocahontas Coal; elevation. 2580' B.

	Ft.	In.
Coal, reported.....	4	0

Post-office address, Charmco; Mine Foreman, Will Lang; on Nicholas, Fayette, and Greenbrier Railroad.

A prospect opening at approximately the same location as the above mine was measured and sampled before the mine was opened. The section measured is as follows:

			Ft.	In.
1. Shale, black, fossiliferous, Royal.....				
2. Coal, good, laminated with fusain (mineral charco- coal)	1'	10"		
3. Fusain (mineral charco- al)..	0	1		
4. Coal, clean, good.....	1	6		
5. Coal, hard	0	4	3	9

A sample (No. 141PH) was taken from Nos. 2, 3, 4, and 5 of the above section, the analysis of which is published under No. 428 in the Table of Coal Analyses at the end of this Chapter.

T. E. and S. T. Jones Mine—No. 429 on Map II.

Truck mine, formerly Ed Grafton Mine; on west side of Laurel Creek, 0.35 mile north of Charmco; No. 6 Pocahontas Coal; elevation, 2665' B.

			Ft.	In.
Coal, slightly bony (black shale roof, good).....	0'	4"		
Coal, laminated with fusain				

A sample (No. 142PH) was taken from the above section, the analysis of which is published under **No. 429** in the Table of Coal Analyses at the end of this Chapter.

Joe Neff Mine No. 1—No. 430 on Map II.

Truck mine, on Snowden Crane property; on west side of Laurel Creek, 0.85 mile north-northeast of Charmco; No. 6 Pocahontas Coal; elevation, 2670' B.

			Ft.	In.
Coal, banded, bright and dull, (shale roof)	1'	8"		
Coal, soft, partly columnar.....	1	4	3	0

A sample (No. 148PH) was taken from the above section, the analysis of which is published under **No. 430** in the Table of Coal Analyses at the end of this Chapter.

Lester Boyer Mine—No. 431 on Map II.

On west side of Laurel Creek, below public road, 1.25 miles north-east of Charmco; No. 6 Pocahontas Coal; elevation, 2665' B.

			Ft.	In.
Coal, hard, bony (black shale roof)	0'	1"		
Coal, hard	0	6		
Coal, blocky, but laminated with mineral charcoal	0	10		
Coal, columnar	1	6		
Coal, hard (slate floor).....	0	8	3	7

A sample (No. 147PH) was taken from the above section, the analysis of which is published under **No. 431** in the Table of Coal Analyses at the end of this Chapter.

The Gauley Coal Land Company has recently prospected the south end of Mill Creek Mountain. The prospecting was completed too late to be shown on Map II. However, Figure 22 shows the outcrop of the **No. 6 Pocahontas Coal** as shown for this area on Map II to which have been added the approximate locations of the following six openings:

Coal Exposure—No. 427 on Map II.

On south side of Meadow Creek, 0.5 mile from mouth; No. 6 Pocahontas Coal, elevation, 2480' B.; section reported by Mr. Wm. McClung.

			Ft.	In.
Coal (shale roof).....	2'	1"		
Parting	0	1		
Coal	1	10	4	0

Greenbrier Fire Creek Coal Company "Midland" Mine— No. 428 on Map II.

Same as Midland New Mine, formerly Midland Smokeless Coal Company; 0.8 mile northwest of Charmco, on north side of Meadow River; No. 6 Pocahontas Coal; elevation, 2580' B.

	Ft.	In.
Coal, reported.....	4	0

Post-office address, Charmco; Mine Foreman, Will Lang; on Nicholas, Fayette, and Greenbrier Railroad.

A prospect opening at approximately the same location as the above mine was measured and sampled before the mine was opened. The section measured is as follows:

			Ft.	In.
1. Shale, black, fossiliferous, Royal.....				
2. Coal, good, laminated with fusain (mineral charcoal)	1'	10"		
3. Fusain (mineral charcoal)..	0	1		
4. Coal, clean, good.....	1	6		
5. Coal, hard	0	4	3	9

A sample (No. 141PH) was taken from Nos. 2, 3, 4, and 5 of the above section, the analysis of which is published under No. 428 in the Table of Coal Analyses at the end of this Chapter.

T. E. and S. T. Jones Mine—No. 429 on Map II.


Truck mine, formerly Ed Grafton Mine; on west side of Laurel Creek, 0.35 mile north of Charmco; No. 6 Pocahontas Coal; elevation, 2665' B.

			Ft.	In.
Coal, slightly bony (black shale roof, good).....	0'	4"		
Coal, laminated with fusain (mineral charcoal).....	0	4		
Fusain (mineral charcoal).....	0	0½		
Coal, laminated	0	7		
Coal, hard	1	1		
Coal, soft	1	3		
Coal, hard	0	4	3	11½

Gauley Coal Land Company Prospect X on Figure 22.

On east side of Laurel Creek, 1.29 miles northeast of mouth; sampled by Gauley Coal Land Company and analyzed by Commercial Testing and Engineering Company, Charleston, W. Va., as reported by the former Company; No. 6 Pocahontas Coal; elevation, 2668' L.

Figure 22.—Map showing the location of recent prospect openings in the No. 6 Pocahontas Coal.



MEASURED TAKEN		110 feet in		90 feet in		45 feet in	
		Ft. in.		Ft. in.		Ft. in.	
Coal		1		2		4	
		A. S.		D. B.		A. S.	
Analyses		Per cent.		Per cent.		Per cent.	
		8.54		26.28		13.28	
Moisture		24.01		28.42		32.75	
Volatile Matter		64.88		70.01		60.77	
Fixed Carbon		2.54		2.78		5.02	
Ash		100.00		100.00		100.00	
Totals		0.50		0.52		0.63	
Sulphur		12.621		13.709		9.632	
B. T. U.		15.397		15.237		10.085	

Gauley Coal Land Company Prospect W on Figure 22.

On east side of Laurel Creek, 1.05 miles northeast of mouth; sampled by Gauley Coal Land Company and analyzed by Commercial Testing and Engineering Company, Charleston, W. Va., as reported by the former Company; No. 6 Pocahontas Coal; elevation, 2696' L.

Coal		Ft.	
13	3		
Analysis			
As Sampled.		Dry Basis.	
Percent.		Percent.	
Moisture	1.11		
Volatle Matter	22.15		
Fixed Carbon	72.81		
Ash	3.93		
Totals		100.00	100.00
Sulphur	0.72		0.73
B. T. U.	14,946		15,114

Gauley Coal Land Company Prospect V on Figure 22.

On east side of Laurel Creek, 0.55 mile northeast of mouth; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 2716' L.

Coal	2'	7"		
Coal and bone	0	4		
			2	11
Ft.		In.		

Gauley Coal Land Company Prospect U on Figure 22.

On east side of Laurel Creek, 0.55 mile east of mouth; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 2727' L.

Coal	0'	6"		
Bone	0	2		
Coal	1	3		
			1	11
Ft.		In.		

Gauley Coal Land Company Prospect T on Figure 22.

On south side of east branch of Laurel Creek, 0.55 mile southeast of Charcoal; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 2786' L.

Coal	1'	4"		
Laminated	0	2		
Coal	0	2		
State	0	1		
Coal	0	10		
			2	7
Ft.		In.		

Gauley Coal Land Company Prospect S on Figure 22.

On southwest end of Mill Creek Mountain, 0.35 mile southeast of Charneo; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 2767' L.

Coal	1'	2 3/4"	2	10
Coal and slate	1	7 3/4		

Gauley Coal Land Company Prospect No. 607— No. 432 on Map II.

On east side of Mill Creek, 2.65 miles east of Charneo and 3.05 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 2818' L.

Coal	0'	8"		
Slate	6	0		
Coal	0	6		
Slate	1	8		
Coal (fire clay floor)	2	6	11	4

Gauley Coal Land Company Prospect No. 561— No. 433 on Map II.

On east side of Mill Creek, 2.9 miles east of Charneo and 2.85 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 2887' L.

Coal	1		6
	Ft.	In.		

On south end of Big Clear Creek Mountain, 2.4 miles southeast of Charneo and 1.55 miles northwest of Rupert; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 2955' L.

Coal and bone	2		2
	Ft.	In.		

Gauley Coal Land Company Prospect M—No. 435 on Map II.

On south end of Big Clear Creek Mountain, 1.05 miles northeast of Rupert; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 3086' L.

Coal	0'	7"		
Slate	0	1		
Coal	1	2		
Slate	0	2		
Coal (fire clay floor)	2	0	4	0
	Ft.	In.			

Gauley Coal Land Company Prospect K—No. 436 on Map II.

On south end of Big Clear Creek Mountain; 0.85 mile northeast of Rupert; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 3144' L.

Coal and slate	0'	4"
Coal	1	0
Bone	0	2
Coal (fire clay floor)	1	8
.....	3	
.....	2	

Gauley Coal Land Company Prospect AF—No. 437 on Map II.

On west side of Big Clear Creek, 1.45 miles north of Rupert; used in Big Clear Creek Mountain Section; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 3093' L.

Coal and bone	2	
.....	6	

Gauley Coal Land Company Prospect AE—No. 438 on Map II.

On west side of Big Clear Creek, 1.7 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 3100' L.

Coal and bone	2	
.....	0	

Gauley Coal Land Company Prospect AD—No. 439 on Map II.

On west side of Big Clear Creek, 1.85 miles north the Rupert; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 3079' L.

Coal and slate	2	
.....	5	

Gauley Coal Land Company Prospect AC—No. 440 on Map II.

On west side of Big Clear Creek, 1.95 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 3070' L.

Coal	1	
.....	5	

Gauley Coal Land Company Prospect AB—No. 441 on Map II.

On west side of Big Clear Creek, 2.05 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 3060' L.

Coal and slate	1	
.....	5	

Gauley Coal Land Company Prospect AA—No. 442 on Map II.

On east side of Big Clear Creek, 3.1 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal; elevation, 3025' L.

Coal and bone	2	Ft.
.....	1	In.

Leckie Smokeless Coal Company Prospect—No. 443 on Map II.

On the east side of Brown Creek, 1.15 miles north of its mouth and 1.55 miles northwest of Anjean; No. 6 Pocahontas Coal; elevation, 2860' B.

Sandstone, conglomerate	5	Ft.
Coal	3	In.
Congealed	0	

Leckie Smokeless Coal Company Prospect—No. 444 on Map II.

On the east side of Pollock Mountain, 3.05 miles southwest of Duo and 1.2 miles north of Anjean; No. 6 Pocahontas Coal; elevation, 2920' B.

1. Sandstone, grayish-brown	0'	6"
2. Coal, bright, good	0	4
3. Shale	0	4
4. Coal, clean	0	2½
5. Bone	0	4
6. Coal, dull to bright	2	10
7. Shale	15	
8. Sandstone, irregular bedding	40	

A sample (No. 151PH) was taken from Nos. 2, 4, and 6 of the above section, the analysis of which is published under No. 444 in the Table of Coal Analyses at the end of this Chapter.

L. E. McClung Prospect—No. 445 on Map II.

On the east side of Big Clear Creek, 2.2 miles southwest of Duo and 2 miles northeast of Anjean; No. 6 Pocahontas Coal; elevation, 2960' B. (2940' L.?).

Coal, reported by L. E. McClung, 3' 0" to

As reported by Gauley Coal Land Company for what appears to be the same location, elevation, 2940' L.

Coal	1'	5"
Shale	0	6
Coal	1	7
Coal and slate	0	3
Coal	0	7

Gauley Coal Land Company Prospect No. 591—
No. 446 on Map II.

On the west side of the south end of Shellcamp Ridge, 0.95 mile northeast of Anjean; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 2955' L.

Bono	0'	8 1/2"
Coal	1	7
Slate	0	2 1/2
Coal	1	11
Coal	4	5

Gauley Coal Land Company Prospect No. 590E—
No. 447 on Map II.

On the west side of the south end of Shellcamp Ridge, 0.85 mile northeast of Anjean; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 2984' L.

Coal	0'	9"
Slate	0	5
Coal	0	3 1/2
Bono	0	2
Coal	1	9
Slate	0	3
Coal	1	9
Coal	1	10 1/2"
Coal and slate	3	6 1/2
Coal	5	5
Bone	2	6 1/2

Gauley Coal Land Company Prospect No. 590B—
No. 448 on Map II.

On the east side of the south end of Shellcamp Ridge, 0.65 mile northeast of Anjean; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 3020' L.

Coal and slate	1'	10 1/2"
Coal	3	6 1/2
Coal	5	5

Gauley Coal Land Company Prospect No. 590A—
No. 449 on Map II.

On the east side of the south end of Shellcamp Ridge, 0.9 mile northeast of Anjean; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 3005' L.

Coal	0'	8"
Sandstone	2	6
Coal	1	2
Coal	4	4

**Gauley Coal Land Company Prospect No. 590—
No. 450 on Map II.**

On the east side of Shellcamp Ridge, 1.9 miles northeast of Anjean and 2.45 miles south of Duo; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 3036' L.

Coal	3
Slate	0
	7	
In.		

**Gauley Coal Land Company Prospect No. 1—
No. 451 on Map II.**

On the west side of Smokehouse Branch, 1 mile north of its mouth and 1.4 miles southeast of Duo; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 3164' L.

Coal (slate roof and floor)	2
	10	
In.		

**Gauley Coal Land Company Prospect No. 11—
No. 452 on Map II.**

On the southwest end of Smokehouse Ridge, 0.3 mile northeast of the mouth of Smokehouse Branch and 2.2 miles southeast of Duo; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 3312' L.

Coal, bone, and slate (slate roof and floor)	8
	3	
In.		

**Gauley Coal Land Company Prospect No. 21—
No. 453 on Map II.**

On Oldhouse Branch, 0.15 mile north of its mouth and 2.5 miles southeast of Duo; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 3339' L.

Coal	3
	8	
In.		

**Gauley Coal Land Company Prospect No. A314—
No. 454 on Map II.**

On the north side of Little Clear Creek Mountain, 0.35 mile south-east of mouth of Old Field Branch; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 3434' L.

Coal	3
	11	
In.		

**Gauley Coal Land Company Prospect No. A313—
No. 455 on Map II.**

On the north side of Little Clear Creek Mountain, 0.55 mile southwest of mouth of Old Field Branch; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 3451' L.

Coal	2'	10½"	
State	1	0½	
Coal (fire clay floor)	1		5
			0

A sample of coal was taken at the above location by the Gauley Coal Land Company and analyzed by the Commercial Testing and Engineering Company, of Charleston, W. Va. The analysis as reported by the former company is published under No. 455 in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. A312—
No. 456 on Map II.**

On north side of Little Clear Creek Mountain, 1.2 miles east of mouth of Smokehouse Branch and 0.8 mile southwest of mouth of Old Field Branch; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 3448' L.

Coal	3		
State	3		
Coal (fire clay floor)	3		0

**Gauley Coal Land Company Prospect No. A405—
No. 457 on Map II.**

On the north side of Briery Knob, 0.4 mile southeast of mouth of Smokehouse Branch; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 3300' L.

Coal	2		
State	2		
Coal	2		6½

**Gauley Coal Land Company Prospect No. A403—
No. 458 on Map II.**

On the south side of Briery Creek, 0.9 mile east of mouth; Gauley Coal Land Company authority for this section; No. 6 Pocahontas Coal?; elevation, 3282' L.

Coal	0'	7"	
State	0	2	
Coal (fire clay floor)	1	9½	2
			6½

No. 459 on Map II.

On south side of Briery Creek, 0.7 mile southeast of mouth; Gauley Coal Land Company, authority for this section; No. 6 Pocahontas

Coal	0.	6"		
State and bone	0	3		
Coal	2	0		
Slato (fire clay floor)	0	2		
			2	11

No. 460 on Map II.

On the north side of Little Clear Creek Mountain, 0.5 mile south of mouth of Briery Creek; Ganley Coal Land Company, authority for

	In.	Ft.
Coal (slate roof)	07	"8"
Bone	0	1½
Coal (fire clay floor).....	1	10¾
	
	2	
	8	

Coal Exposure—No. 461 on Map II.

On fire road, on south side of Little Clear Creek Mountain, 1.2 miles south of Anjeau; used in Little Clear Creek Section; No. 6

Pl. In.

No. 462 on Map II.

On the south end of Kuhn Ridge, 1.1 miles north of mouth of Kuhn Branch; Gaulty Coal Land Company authority for this section; No. 6

Coal (slate roof).....	4	113
Slate, soft (fire clay floor).....	0	2

Gauley Coal Land Company "Hume" Mine (Abandoned)—
No. 463 on Map II.

Graham Smokeless Coal Company property; on east side of Point Mountain, south side of Little Clear Creek Mountain, 2 miles northeast of mouth of Kuhn Branch; No. 6 Pocahontas Coal; elevation, 3537' L.

Coal, laminated, bright with fusain (mineral charcoal), (slate roof)	0.	5"	0	0	0	1	6	4	2
Coal, good, columnar	2	0	0	0	0	0	3			
Coal, gray bands	0	0	0	0	0	0	0			
Coal, clean, hard, laminated (slate floor).....	1	0	0	0	0	0	0			

A sample (No. 129PH) was taken from the above section, the analysis of which is published under No. 463 in the Table of Coal Analyses at the end of this Chapter.

Gauley Coal Land Company Prospect No. A302—
No. 464 on Map II.

Graham Smokeless Coal Company property; on west side of Middle Mountain, east side of Little Clear Creek, 2.15 miles northeast of mouth of Kuhn Branch; No. 6 Pocahontas Coal; elevation, 3549' L.

Gauley Coal Land Company Prospect No. A301—
No. 465 on Map II.

Graham Smokeless Coal Company property; on west side of Middle Mountain, east side of Little Clear Creek, 2.1 miles northeast of mouth of Kuhn Branch; No. 6 Pocahontas Coal; elevation, 3564' L.

No. 6 Pocahontas Coal, Williamsburg and Falling Springs Districts.

In these districts the No. 6 Pocahontas Coal is almost entirely unprospected, only two prospects being noted. The bed is believed, however, to attain practically the same development as in Meadow Bluff District. The horizon of this seam, the outcrop of which is delineated on Map II, extends over a greater area than the probable minable area shown on

Figure 21. The descriptions of the prospects, both of which are in Williamsburg District, follow:

Prospect—No. 466 on Map II.

On north end of Buffalo Mountain, 3.65 miles northwest of Williamsburg; No. 6 Pocahontas Coal; elevation, 3570' B.
Coal (reported by B. M. Higginbotham)..... 1
Ft. 1
In. 8

Gauley Coal Land Company Prospect—No. 467 on Map II.

On the waters of Horgamp Run, 1.05 miles from its mouth, and 2.15 miles east of Beech Knob; authority, Gauley Coal Land Company; No. 6 Pocahontas Coal; elevation, 3055' L.

Coal	0'	10"
Slate	1	3
Coal	1	4
	3	5
	Ft.	In.

Quantity of No. 6 Pocahontas Coal Available.

The following table computed from planimetric measurement of the outcrop of the seam as shown on Map II for the area indicated on Figure 21, page 581, gives the probable amount of No. 6 Pocahontas Coal in Greenbrier County. A low figure for the average thickness was assumed so that the total would not be too great if local areas prove to be cut out or too thin for mining:

Probable Amount of No. 6 Pocahontas Coal.

District.	Thickness of Coal Assumed. Feet.	Square Miles.	Acres.	Cubic Feet of Coal.	Short Tons of Coal, (2000 Lbs.)
Meadow Bluff	2	81.80	52,352	6,841,359,360	273,654,374
Williamsburg	2	18.45	11,808	1,028,712,960	41,148,518
Falling Springs	2	4.00	2,560	223,027,200	8,921,088
Totals		104.25	66,720	8,093,099,520	323,723,980

NO. 3 POCAHONTAS COAL.


The No. 3 Pocahontas Coal, previously described in Chapter VI, constitutes a small but valuable reserve of coal in Greenbrier County. It is probably the lowest minable bed in the county. In general it is multiple-bedded, soft, co-lummar, and has been mined locally at a few points. It may reach a total thickness of over five feet but is usually impure when present in that thickness. Chemical analyses reveal an excellent fuel with a low volatile content and high fusion point of the ash.

The extent of the probable minable area of No. 3 Pocahontas Coal is shown on Figure 23. Its interval below No. 6 Pocahontas is approximately 100 feet, so that its position on Map II may be easily interpolated from the position of that seam.

SHOWING
PROBABLE AREA OF MINABLE
NO. 3 POCAHONTAS COAL

MAP

01



Coal Exposure—No. 370B on Map II.

On Cold Knob road, 0.7 mile southeast of Manning Knob and 0.9 mile northwest of Blue Knob; Fire Creek Coal; elevation, 3677' L.

Ft. In.

Coal, thickness not determined.....

Coal Exposure—No. 370C on Map II.

On Cold Knob road, 0.48 mile southwest of Blue Knob and 0.9 mile northwest of Big Bull Hill; Fire Creek Coal; elevation, 3865' B.

Ft. In.

Coal and black shale, thickness not determined.....

Coal Exposure—No. 370D on Map II.

On Cold Knob road, 0.57 mile south of Blue Knob and 0.7 mile northwest of Big Bull Hill; Fire Creek Coal; elevation, 3930' B.

Ft. In.

Coal, exposed 1 6

Coal Exposure—No. 370E on Map II.

On Cold Knob road, 0.72 mile south of Blue Knob and 0.42 mile north of Big Bull Hill; Fire Creek Coal; elevation, 3950' B.

Ft. In.

Coal, exposed 1 0

Quantity of Fire Creek Coal Available.

The following table, computed from planimetric measurement of outcrop shown on Map II for the area indicated on Figure 20, page 555, gives the probable amount of Fire Creek Coal in Greenbrier County. A low figure for the average thickness was assumed so that the total would not be too great if local areas prove to be cut out or too thin for mining:

Probable Amount of Fire Creek Coal.

District.	Thickness of Coal Assumed. Feet.	Square Miles.	Acres.	Cubic Feet of Coal.	Short Tons of Coal (2000 Lbs.)
Meadow Bluff	3	93.5	59,840	7,819,891,200	312,795,648
Williamsburg	2	22.0	14,080	1,226,649,600	49,065,984
Falling Springs	2	36.5	23,360	2,035,123,200	81,404,928
Totals		152.0	97,280	11,081,664,000	443,266,560

MINABLE COALS OF THE POCAHONTAS GROUP OF POTTSVILLE SERIES.

In general, the coals of this group are soft, columnar, and multiple-bedded. They are medium-low volatile, low sulphur, low ash, high B. T. U. coals which have a fusion point of ash ranging from 2500° F. to 2900° F. Coals with these qualities are well adapted for use in automatic stokers. In view of recent inventions it would appear that here is a large potential market for the coals of Greenbrier County.

As previously mentioned in Chapter VI, it has been necessary to consider No. 7 Pocahontas and No. 6 Pocahontas Coals together. To avoid confusion, however, the openings in each seam have been numbered consecutively and will be described separately on the following pages:

NO. 7 POCAHONTAS COAL.***No. 7 Pocahontas Coal, Meadow Bluff District.***

In general, the coal that is here provisionally correlated as the No. 7 Pocahontas Coal is soft, columnar, and multiple-bedded, and ranges in thickness from 4 feet to the vanishing point. This coal has been mined on the south end of Little Sewell Mountain, near the mouth of Meadow Creek, in the vicinity of Charmco and on the south end of Big Clear Creek

Mountain. These are all truck mines and only two (Nos. 385A and 400) are in regular operation at the present time (1936).

Due to the apparently irregular nature of this seam the present information is not considered sufficient to predict the probable minable area or to estimate the available tonnage.

W. H. Sims Coal Prospect—No. 379 on Map II.

On the east side of Sims Mountain, 400 to 500 feet due south of Sims School and 2 miles due south of Rainelle; described by Ray V. Hemmen under No. 264 on page 321, Fayette County Report; No. 7 Pocahontas Coal?; elevation, 3035' B.

			Ft.	In.
Coal, bony and slaty	1'	0"		
Coal, soft, conchoidal	1	6	2	6

Coal Prospect—No. 380 on Map II.

On Wm. Bennett land, on east side of road on Little Sewell Mountain, 2.1 miles southwest of Rupert and 1.7 miles northeast of Meadowvale School; No. 7 Pocahontas Coal; elevation, 3200' B.

	Ft.	In.
Coal (opening partially filled with water).....	2	8+

C. N. Callison Mine (Abandoned)—No. 381 on Map II.

On southwest side of Little Sewell Mountain, 2.2 miles southwest of Rupert and 1.75 miles northeast of Meadowvale School; No. 7 Pocahontas Coal; elevation, 3230' B.

	Ft.	In.
Coal (shale roof)	3	5+

The opening was partially filled with water. A sample (No. 135PH) was taken from the upper 3 feet 5 inches of coal, the analysis of which is published under No. 381 in the Table of Coal Analyses at the end of this Chapter.

Coal Mine (Abandoned)—No. 382 on Map II.

On southwest side of Little Sewell Mountain, 2.25 miles southwest of Rupert and 1.85 miles northeast of Meadowvale School; No. 7 Pocahontas Coal; elevation, 3250' B.

	Ft.	In.
Coal, reported (fallen shut)	4	0

Fossil collection No. 140 taken from roof shales.

E. H. Callison Mine (Abandoned)—No. 383 on Map II.

On south end of Little Sewell Mountain, 2.5 miles south of Rupert and 1.8 mile east of Meadowvale School; No. 7 Pocahontas Coal; elevation, 3285' B.

			Ft.	In.
Coal, bony	0'	3"		
Coal, block	0	3		
Pyrite	0	0½		
Coal, columnar, laminated with fusain (mineral charcoal)..	0	9		
Coal, hard	0	5		
Coal, laminated with fusain (mineral charcoal) and pyrite	1	4		
Coal, bony	0	4	3	4½

A sample (No. 136PH) was taken from the above section, the analysis of which is published under **No. 383** in the Table of Coal Analyses at the end of this Chapter.

Evely Mine (Abandoned)—No. 384 on Map II.

On south side of Meadow Creek, 0.2 mile east of mouth; No. 7 Pocahontas Coal; elevation, 2525' B.

			Ft.	In.
1. Slate, black (roof not good).....			0	6
2. Coal, small blocks	1'	10"		
3. Coal, soft, thin streaks of bone (sandstone floor) ..	1	8	3	6

A sample (No. 146PH) was taken from Nos. 2 and 3 of the above section, the analysis of which is published under **No. 384** in the Table of Coal Analyses at the end of this Chapter.

Coal Mine (Abandoned)—No. 385 on Map II.

Truck mine, on northwest side of Charmco-Quinwood road, 0.95 mile northeast of Charmco; used in Charmco Section; No. 7 Pocahontas Coal; elevation, 2695' B.

	Ft.	In.
Coal, fallen slnt, visible 2' to	3	0

Lester Boyer Mine—No. 385A on Map II.

Truck mine, on west side of Laurel Creek, 1.35 miles northeast of Charmco; No. 7 Pocahontas Coal; elevation, 2695' B.

			Ft.	In.
Coal	0'	6"		
Bone	1	0		
Coal, dirty, with bone partings..	2	0	3	6

The above section was reported by one of the men working at the mine.

Gauley Coal Land Company Prospect AR—No. 386 on Map II.

On east side of Mill Creek, 2.7 miles east of Charmeo and 3 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal?; elevation, 2878' L.

	Ft.	In.
Coal	1	6

Gauley Coal Land Company Prospect AQ—No. 387 on Map II.

On east side of Mill Creek, 2.6 miles southeast of Charmeo and 2.1 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 2990' L.

	Ft.	In.
Coal and bone	2	2

Gauley Coal Land Company Prospect AP—No. 388 on Map II.

On east side of Mill Creek, 2.5 miles southeast of Charmeo and 2.05 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 2982' L.

		Ft.	In.
Coal (sandstone roof)	0' 1"		
Slate	0 2		
Bone	0 3		
Coal	2 11½		
Bone (fire clay floor)	0 2	3	7½

Gauley Coal Land Company Prospect AN—No. 389 on Map II.

On east side of Mill Creek, 2.55 miles southeast of Charmeo and 1.95 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 3001' L.

	Ft.	In.
Coal and bone	2	4

Gauley Coal Land Company Prospect AM—No. 390 on Map II.

On east side of Mill Creek, 2.6 miles southeast of Charmeo and 1.9 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 3006' L.

	Ft.	In.
Coal and bone	1	11

Gauley Coal Land Company Prospect AL—No. 391 on Map II.

On east side of Mill Creek, 2.75 miles southeast of Charmeo and 1.75 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 3036' L.

	Ft.	In.
Coal and bone	2	7

Gauley Coal Land Company Prospect AK—No. 392 on Map II.

On east side of Mill Creek, 2.4 miles southeast of Charmeo and 1.8 miles northwest of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 3003' L.

	Ft.	In.
Coal	2	2

Gauley Coal Land Company Prospect AJ—No. 393 on Map II.

On the east side of Mill Creek, 2.2 miles southeast of Charmeo and 1.95 miles northwest of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 2966' L.

			Ft.	In.
Coal	1'	3"		
Bone	0	2		
Coal (fire clay floor)	1	2	2	7

Gauley Coal Land Company Prospect AI—No. 394 on Map II.

On the east side of Mill Creek, 2.2 miles southeast of Charmeo and 1.95 miles northwest of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 2956' L.

			Ft.	In.
Coal	0'	5"		
Bone	0	1		
Coal (fire clay floor)	2	6	3	0

**Gauley Coal Land Company Prospect No. 603—
No. 395 on Map II.**

On east side of Mill Creek Mountain, 2.15 miles southeast of Charmeo and 1.95 miles northwest of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 2957' L.

	Ft.	In.
Coal	1	6

Gauley Coal Land Company Prospect AH—No. 396 on Map II.

On east side of Mill Creek, 2.15 miles southeast of Charmeo and 1.85 miles northwest of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 2946' L.

	Ft.	In.
Coal	1	6

Gauley Coal Land Company Prospect J—No. 397 on Map II.

On the west side of Big Clear Creek Mountain, 1.05 miles north of Rupert; No. 7 Pocahontas Coal; Gauley Coal Land Company authority for this section; elevation, 3180' L.

			Ft.	In.
Coal	3'	8"		
Slate	0	0½		
Coal and bone (fire clay floor) ..	0	3	3	11½

Gauley Coal Land Company Prospect I—No. 398 on Map II.

On west side of Big Clear Creek, 1.1 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 3168' L.

			Ft.	In.
Coal (slate roof)	0'	3"		
Slate	0	1		
Coal	3	1½		
Slate	0	1		
Coal (fire clay floor)	0	7½	4	2

Gauley Coal Land Company Prospect H—No. 399 on Map II.

On the west side of Big Clear Creek, 1.4 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 3154' L.

			Ft.	In.
Coal (slate roof)	3'	11½"		
Bone (fire clay floor)	0	2½	4	2

Amick Mine—No. 400 on Map II.

Truck mine, on west side of Big Clear Creek, 1.45 miles north of Rupert; No. 7 Pocahontas Coal; elevation, 3135' L.

			Ft.	In.
Sandstone				
Bone, 1' to			2	0
Coal, bright	1'	8"		
Coal, dull	2	0	3	8

Gauley Coal Land Company Prospect F—No. 401 on Map II.

On west side of Big Clear Creek, 2.1 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal?; elevation, 3093' L.

			Ft.	In.
Coal and slate			1	0

Gauley Coal Land Company Prospect E—No. 402 on Map II.

On west side of Big Clear Creek, 3.1 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 3070' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	2	8

Gauley Coal Land Company Prospect D—No. 403 on Map II.

On west side of Big Clear Creek, 3.2 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 3066' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	3	11

Gauley Coal Land Company Prospect C—No. 404 on Map II.

On west side of Big Clear Creek, 3.3 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 3069' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	4	4

Gauley Coal Land Company Prospect A—No. 405 on Map II.

On west side of Big Clear Creek, 1.9 miles southwest of Anjean and 3.5 miles north of Rupert; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal; elevation, 3068' L.

	Ft.	In.
Coal (fire clay floor)	3	2

**Gauley Coal Land Company Prospect No. A311—
No. 406 on Map II.**

On the north side of Joe Knob, 1 mile southeast of mouth of Smokehouse Branch; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal?; elevation, 3449' L.

	Ft.	In.
Coal (slate roof and floor)	2	6

**Gauley Coal Land Company Prospect No. A310—
No. 407 on Map II.**

On the northeast side of Briery Knob, 0.45 mile southeast of Smokehouse Branch; Gauley Coal Land Company authority for this section; No. 7 Pocahontas Coal?; elevation, 3374' L.

			Ft.	In.
Coal	0'	5½"		
Slate	2	4½		
Coal (slate floor).....	1	7	4	5

NO. 6 POCAHONTAS COAL.

The No. 6 Pocahontas Coal, previously described in Chapter VI, ranks second in Greenbrier County in available tonnage. In general it is soft, columnar, multiple-bedded, and ranges from 1 to 5 feet in thickness. In chemical properties it is an excellent fuel, having a volatile content of 21 to 23 per cent., ash content of from 3 to 6 per cent., and a B. T. U. value that often exceeds 15,000. This coal has been mined at several points and a few small mines are in regular operation at the present time.

Figure 21 shows the probable minable extent of the No. 6 Pocahontas Coal, and its detailed outcrop is shown on Map II.

FIGURE 21

MAP


OF

GREENBRIER COUNTY

SHOWING

PROBABLE AREA OF MINABLE

NO. 6 POCAHONTAS COAL



No. 6 Pocahontas Coal, Meadow Bluff District.

Nearly all of the prospecting and all of the mines opened in the No. 6 Pocahontas Coal are in this district. Its stratigraphic position and thickness are shown in the sections published in Chapter V for Big Clear Creek, Little Clear Creek, Little Sewell Mountain—Southeast, Little Sewell Mountain—West Side, Sims Mountain—North End, Sims School, and Sims Station, page references to which are given in the Index; and in the records of Coal Test Borings Nos. 1, 5C, 6, 11, 12, 13, 14, and 15 published on preceding pages of this Chapter. The following openings and prospects were noted:

Bellwood Coal Company Mine No. 1—No. 408 on Map II.

Fayette County, Quinnimont District; on east side of Quinton Branch, 0.85 mile south of its mouth, near Bellwood; No. 6 Pocahontas Coal; elevation, 2823.0' L.

No section at mine mouth; see borings Nos. 143-152 for thickness.

Bellwood Coal Company Mine No. 2—No. 409 on Map II.

Fayette County, Quinnimont District; on east side of Quinton Branch, 1.05 miles south of its mouth, near Bellwood; No. 6 Pocahontas Coal; elevation, 2832.2' L.

No section at mine mouth; see borings Nos. 143-152 for thickness.

Bellwood Coal Company Mine No. 3—No. 410 on Map II.

Fayette County, Quinnimont District; on west side of Quinton Branch, 1.1 miles south of its mouth, near Bellwood; No. 6 Pocahontas Coal; elevation, 2817.8' L.

No section at mine mouth; see borings Nos. 143-152 for thickness.

George Shawver Mine—No. 411 on Map II.

Fayette County, Quinnimont District; farm mine, 2.15 miles east of Springdale and 1.05 miles southwest of Coal Hollow School; No. 6 Pocahontas Coal; elevation, 3065' B.

	Ft.	In.
Shale, black, Royal, Lingula fossils abundant.....		
Coal, clean, columnar, with layers of fusain (mineral charcoal), (slate roof and floor).....	3	2

A sample (No. 132PH) was taken from the above section, the analysis of which is published under No. 411 in the Table of Coal Analyses at the end of this Chapter.

A section was measured at the same mine by Ray V. Hennen and reported under No. 576 in the Fayette County Report, page 855, as follows:

	Ft.	In.
Shale, black, Royal, <i>Lingula fossilis</i> abundant.....		
Coal, soft	4	0
Shale, dark and gray	10	0

Coal Exposure—No. 411A on Map II.

On public road, 2.2 miles east of Springdale and 1 mile southwest of Coal Hollow School; No. 6 Pocahontas Coal, Lower Bench; elevation, 3055' B.

	Ft.	In.
Coal, exposed	2	0

Bert Hutsonpillar Mine—No. 412 on Map II.

Farm mine, operated by C. C. Helmick, on north side of Turniphole Mountain, 2.5 miles east of Springdale; No. 6 Pocahontas Coal; elevation, 3065' B.

		Ft.	In.
1. Coal, bony (shale roof, poor)	0' 4"		
2. Coal, hard	0 10		
3. Coal, columnar	0 10		
4. Coal, hard	0 4		
5. Coal, bright, laminated with fusain (mineral charcoal)	1 5		
6. Coal, bony	0 4	4	1

A sample (No. 133PH) was taken from Nos. 1 to 5 inclusive of the above section, the analysis of which is published under No. 412 in the Table of Coal Analyses at the end of this Chapter.

J. A. and S. J. Wooldridge Coal Mine—No. 413 on Map II.

Farm mine, on southeast side of Turniphole Mountain, 2.9 miles northwest of Dawson and 2.85 miles east of Springdale; No. 6 Pocahontas Coal?; elevation, 3120' B.

		Ft.	In.
1. Sandstone, grayish-brown, cross-bedded, micaceous		50	0
2. Concealed		30	0
3. Shale, dark-gray to brown		10	0
4. Coal, bony (slate roof).....	0' 4"		
5. Coal, clean, columnar	1 10		
6. Coal, hard	0 4		
7. Coal, clean, columnar (slate floor)	2 1	4	7
8. Concealed		12	0
9. Coal (reported)		3	0

A sample (No. 131PH) was taken from Nos. 4, 5, 6, and 7 of the above section, the analysis of which is published under **No. 413** in the Table of Coal Analyses at the end of this Chapter.

Coal Prospect—No. 414 on Map II.

On Sims Mountain, on public road, 2.7 miles south of Rainelle; used in Sims Station Section, page 158; No. 6 Pocahontas Coal?; elevation, 3082' L.

	Ft.	In.
Coal, reported 8" to	1	0

Coal Exposure—No. 415 on Map II.

On north end of Sims Mountain, on public road, 0.4 mile south of East Rainelle; used in Sims Mountain Section—North End; No. 6 Pocahontas Coal?; elevation, 2840' B.

	Ft.	In.
Coal, exposed	0	2

S. H. Samples Mine (Abandoned)—No. 416 on Map II.

Farm mine, on east side of Goddard Mountain, 2.3 miles southeast of East Rainelle and 3.2 miles southwest of Rupert; No. 6 Pocahontas Coal?; elevation, 3085' B.

	Ft.	In.
Shale	6	0
Coal	1	10

Meadow River Fuel Company "Lincoln" Mine (Abandoned)— No. 417 on Map II.

On northwest end of Little Sewell Mountain, 0.3 mile northeast of East Rainelle; No. 6 Pocahontas Coal; elevation, 2780' B.

		Ft.	In.
Coal, laminated with bright and fusain (mineral charecoal), (slate roof)	0' 6"		
Coal, columnar, good.....	1 2		
Coal, laminated	0 10	2	6

A sample (No. 130PH) was taken from the above section, the analysis of which is published under **No. 417** in the Table of Coal Analyses at the end of this Chapter. At the point of sampling the coal measured 30 inches; however, it is reported to be 34 to 36 inches thick in most places in the mine.

Coal Exposure—No. 418 on Map II.

On west side of Little Sewell Mountain, 1 mile southeast of Rainelle; used in Little Sewell Mountain Section—West Side; No. 6 Pocahontas Coal; elevation, 2885' B.

	Ft.	In.
Coal, soft, exposed	0	9

Coal Exposure—No. 419 on Map II.

On west side of Little Sewell Mountain, 0.7 mile northeast of Dennis and 2.25 miles southwest of Rupert; No. 6 Pocahontas Coal; elevation, 3185' B.

	Ft.	In.
Coal, (exposed in spring on land of Mr. Dunn).....	1	0

Coal Prospect—No. 420 on Map II.

On west side Little Sewell Mountain, north side of road, 0.65 mile northeast of Dennis and 2.2 miles southwest of Rupert; No. 6 Pocahontas Coal; elevation, 3200' B.

	Ft.	In.
Coal, fallen shut, thickness reported.....	3	0

Coal Exposure—No. 421 on Map II.

On south end of Little Sewell Mountain, 4.1 miles southeast of East Rainelle and 2.35 miles south of Rupert; No. 6 Pocahontas Coal; elevation, 3270' B.

	Ft.	In.
Coal	3	0

Meadow River Lumber Company Prospect—No. 422 on Map II.

On the east side of Little Sewell Mountain, 1.75 miles west of Rupert; No. 6 Pocahontas Coal; elevation, 3055' B.

	Ft.	In.
Shale, dark		
Coal blossom		
Shale, numerous plants	3	0
Coal, fallen shut, reported.....	5	0

**Meadow River Lumber Company Mine (Abandoned)—
No. 423 on Map II.**

Farm mine, 0.3 mile north of Rainelle; No. 6 Pocahontas Coal; elevation, 2710' B.

	Ft.	In.
1. Shale, black. Royal, Lingua.....	8	0
2. Coal, dirty	0'	1"
3. Coal, laminated, soft, poor..	0	4
4. Coal, columnar	0	5
5. Coal, hard	0	5
6. Coal, columnar, soft	0	10
7. Coal, laminated (slate floor)	0	11
	3	0

A sample (No. 139PH) was taken from Nos. 3, 4, 5, 6, and 7 of the above section, the analysis of which is published under No. 423 in the Table of Coal Analyses at the end of this Chapter.

The section of the following abandoned mine with comments by Ray V. Hennen is reprinted from page 852 of the Fayette County Report:

**Meadow River Smokeless Coal Company "Dwyer" Mine
(Abandoned)—No. 424 on ap II.**

Formerly J. W. Dwyer, same as Tuck Brothers Mine, owned by Meadow River Coal and Land Company; located just east of Fayette-Greenbrier County line, 0.9 mile northeast of Rainelle; No. 6 Pocahontas Coal; elevation, 2605' B.

Section and sample by Ray V. Hennen at rib at starting point of crop entry off 1st left.

	Ft.	In.
1. Slate, black		
2. Slate, black, harder, cannelly, usually draw slate....	0	8
3. Coal, soft, clean (slate, black, floor).....	3	8

"This mine was opened by J. W. Dwyer in 1914 who operated it until May, 1916; principal office, Rainelle, W. Va.; lease of Meadow River Lumber (Coal and Land) Company; output, 100 tons, 9-hour day; men employed, 25 inside and 5 outside; ship run-of-mine coal only and mostly for steam purposes, both east and west, and gives perfect satisfaction; rises rapidly southeast; J. M. Suttle, Foreman, authority for mine data."

A sample (926H) was collected from No. 3 of the above section by Hennen, the analysis of which is published under No. 424 in the Table of Coal Analyses at the end of this Chapter.

**Low Ash Smokeless Coal Company "Green Siding" Mine—
No. 425 on Map II.**

Fayette County, Sewell Mountain District; also known as Peck Mine, 1.75 miles northwest of Rainelle, on west bank of western tributary of Meadow River; No. 6 Pocahontas Coal; elevation, 2470' B.

	Ft.	In.
1. Shale, black, Royal, pelecypods, concretions.....	5	0+
2. Coal, draw in part, dull to bright	0'	6"
3. Coal, laminated with bright and dull	1	5
4. Coal, columnar	0	8
5. Fusain (mineral charcoal)..	0	0½
6. Coal	0	1
7. Fusain (mineral charcoal)..	0	0½
8. Coal, columnar, lumps well	1	2
9. Coal, draw in part.....	0	4
	4	2½

A sample (No. 149PH) was taken from Nos. 2, 3, 4, 5, 6, 7, and 8 of the above section, the analysis of which is published under **No. 425** in the Table of Coal Analyses at the end of this Chapter.

An abandoned opening in the No. 6 (?) Pocahontas Coal was reported near the location of the above mine but on the east bank of the small stream joining Meadow River at Aldrich Camp. The section as shown in the Maywood-Aldrich Camp Section, page 203, of the Fayette Report, is as follows:

	Ft.	In.
Coal, No. 6 Pocahontas, at closed opening (No. 569 on Map II of Fayette County Report) of Meadow Lumber (Coal and Land) Company, reported clean and 48" thick by W. F. Hall, Superintendent.....	4	0
Interval to Meadow River at Aldrich Camp.....	140	0

As noted above, Hennen reports the abandoned opening as 140 feet above the river at Aldrich Camp. Mine No. 425, however, is only 75 feet above the railroad at Aldrich Camp and not over 100 feet above the river at that point. It would appear that the abandoned opening was probably in the No. 7 Pocahontas Coal.

Gauley Coal Land Company Prospect—No. 426 on Map II.

On northwest side of Meadow Creek, 0.8 mile from mouth; No. 6 Pocahontas Coal; elevation, 2460' B.

	Ft.	In.
1. Shale, black, Royal, large pelecypods.....		
2. Coal, blocky, impure.....	0'	4"
3. Coal, banded	1	2
4. Coal, hard, bony.....	0	1
5. Coal, columnar	1	8
6. Coal, banded bright and dull	0	4
7. Shale, sandy		

A sample (No. 145PH) was taken from Nos. 2, 3, 4, 5, and 6 of the above section, the analysis of which is published under **No. 426** in the Table of Coal Analyses at the end of this Chapter.

Fire Creek Coal, Meadow Bluff District.

The best development of the Fire Creek Coal in this district is on Little Clear Creek Mountain where there is a large area of coal with a thickness in excess of 5 feet. In a large part of this district, however, this coal attains a thickness of only 2 to 3 feet. This thinner coal probably could not be mined profitably at the present time but it is here considered as a minable reserve.

The thickness and stratigraphic position of the Fire Creek Coal are noted in the Sims Station Section, Sims Mountain—North End Section, and the Big Clear Creek Mountain Section, all published in Chapter V, and in the records of coal test borings Nos. 5, 11, 13, and 14, all published on preceding pages of this chapter. The stratigraphic position of this coal is shown in the partial records of coal test borings Nos. 5A, 5B, 5D, 5F, 5H, and 5I, published on preceding pages of this chapter. There are no actively operating mines in this coal at the present time (1936). The following prospects and openings were noted:

Gauley Coal and Land Company Prospect (Closed)— No. 310 on Map II.

On the west side of Burdette Creek, 0.2 mile east of Meadow River; No. 545 in Fayette Report; Fire Creek Coal; elevation, 2330' B.; examined by Ray V. Hennen.

	Ft.	In.
Coal, reported by Mr. H. W. Osborne	1	6

"Sulphur spring here. This coal belongs immediately on top of a grayish-white, quartzite sandstone (Pineville?) ellip, 30 to 50 feet thick."

Gauley Coal and Land Company Prospect (Closed)— No. 310A on Map II.

On south bank of Burdette Creek, 0.35 mile east of Meadow River; No. 546 in Fayette Report; Fire Creek Coal; elevation, 2340' B.; examined by Ray V. Hennen.

	Ft.	In.
Coal, reported by Mr. H. W. Osborne	1	8

The above two prospects and comments by Ray V. Hennen are reprinted from page 820 of the Fayette County Report.

Coal exposure **No. 310B on Map II** is published in connection with the Sims Mountain—North End Section in Chapter V, page 158.

The following prospect is reprinted from page 819 of the Fayette County Report:

Thos. Stead Coal Prospect—No. 310C on Map II.

Fayette County, on south hillside of Meadow River, 2.4 miles southeast of Russellville; **Fire Creek Coal**; elevation, 2120' B.; examined by Ray V. Heunen.

			Ft.	In.
Shale, gray, argillaceous, visible.....			3	0
Coal, soft	0'	7"		
Slate, black	0	0½		
Coal, soft (slate floor)	1	8	2	3½

The contours shown on the United States Geological Survey's topographic maps for the region about one mile west of Charmco do not agree with conditions found there. As a result mines Nos. 311, 312, and 428 were very difficult to locate on the map and the correlations of them are doubtful, as indicated by the question marks in the descriptive headings of these mines.

**Midland Smokeless Coal Company "Midland" Mine No. 1
(Abandoned)—No. 311 on Map II.**

On the property of L. E. McClung; on the southwest side of Laurel Creek Mountain, 0.65 mile northwest of Charmco and 0.27 mile north of Meadow River; **Fire Creek? (No. 7 Pocahontas?) Coal**; elevation, 2735' B.

			Ft.	In.
1. Coal, hard, dull (slate roof)	0'	8"		
2. Coal, bright	0	2½		
3. Fusulu (mineral charcoal) ..	0	0½		
4. Coal, good	0	6		
5. Coal, dull, hard	1	0		
6. Coal, laminated dull and bright	0	7		
7. Shale	0	1		
8. Coal, bony	1	0	4	1

A sample (No. 144PH) was taken from Nos. 2, 3, 4, 5, and 6 of the above section, the analysis of which is published under **No. 311** in the Table of Coal Analyses at the end of this chapter.

**Midland Smokeless Coal Company "Midland" Mine No. 2
(Abandoned)—No. 312 on Map II.**

On property of L. E. McClung; on the southwest side of Laurel Creek Mountain, 0.67 mile northwest of Charmco and 0.4 mile north of Meadow River; Fire Creek? (No. 6 Pocahontas?) Coal; elevation, 2700' B.

			Ft.	In.
Coal, bright, laminated (shale roof)	1'	6"		
Coal, hard	0	4		
Coal, soft, columnar	0	10		
Coal, hard (slate floor).....	0	4	3	0

A sample (No. 143PH) was taken from the above section, the analysis of which is published under No. 312 in the Table of Coal Analyses at the end of this chapter.

**Gauley Coal Land Company Prospect No. 562—
No. 313 on Map II.**

On west side of Mill Creek, 3.65 miles north of Rupert; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 2980' L.

	Ft.	In.
Coal and slate	5	0

Gauley Coal Land Company Prospect P—No. 314 on Map II.

On east side of Mill Creek, 2.8 miles east of Charmco and 3.2 miles north of Rupert; Gauley Coal Land Company authority for this section; Fire Creek Coal?; elevation, 3005' L.

	Ft.	In.
Coal	2	0

**Gauley Coal Land Company Prospect No. 558—
No. 315 on Map II.**

On west side of Big Clear Creek Mountain, 1.85 miles north of Rupert; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3204' L.

	Ft.	In.
Coal	1	10

**Gauley Coal Land Company Prospect No. 559—
No. 316 on Map II.**

On west side of Big Clear Creek Mountain, 1.4 miles north of Rupert; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3294' L.

	Ft.	In.
Coal and slate	3	0

**Gauley Coal Land Company Prospect No. 560—
No. 317 on Map II.**

On west side of Big Clear Creek Mountain, 1.3 miles north of Rupert; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3299' L.

	Ft.	In.
Coal	1	11

**Gauley Coal Land Company Prospect No. 557—
No. 318 on Map II.**

On west side of Big Clear Creek, 1.9 miles north of Rupert; Gauley Coal Land Company authority for this section; Fire Creek Coal?; elevation, 3273' L.

	Ft.	In.
Coal	2	1

Leckie Smokeless Coal Company Prospect—No. 319 on Map II.

On west side of Brown Creek, 1.65 miles north of mouth; Fire Creek Coal; elevation, 3028.95' L.

		Ft.	In.
Coal (shale roof)	1' 7"		
Fusain (mineral charecoal).....	0 0½		
Coal	0 10½		
Bone	0 0½		
Coal	0 8		
Bone	0 3		
Coal	0 8		
Bone	0 2		
Coal	0 8	4	11½

**Gauley Coal Land Company Prospect No. 10—
No. 320 on Map II.**

On the west side of the south end of Smokehouse Ridge, 1.05 miles northeast of the mouth of Smokehouse Branch and 2.25 miles southeast of Duo; Gauley Coal Land Company authority for this section; Fire Creek Coal?; elevation, 3529' L.

	Ft.	In.
Coal (slate floor).....	1	8

**Gauley Coal Land Company Prospect No. 13—
No. 321 on Map II.**

On the south end of Smokehouse Ridge, 0.45 mile west of the mouth of Job Knob Branch and 2.5 miles southeast of Duo; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3567' L.

			Ft.	In.
Coal	2'	2"		
Bone	0	3		
Coal (slate floor)	1	6	3	11

**Gauley Coal Land Company Prospect No. 16—
No. 322 on Map II.**

On the east side of Smokehouse Ridge, 0.65 mile north of the mouth of Job Knob Branch and 2.4 miles southeast of Dno; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3564' L.

			Ft.	In.
Coal	1'	5"		
Slate	0	2		
Coal	0	9		
Slate	0	1		
Coal	1	3		
Coal and slate (slate floor).....	1	4	5	0

**Gauley Coal Land Company Prospect No. 20—
No. 323 on Map II.**

On the east side of Smokehouse Ridge, 0.2 mile northwest of the mouth of Oldhouse Branch and 2.4 miles southeast of Duo; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3526' L.

			Ft.	In.
Coal (slate roof)	0'	5"		
Clay	0	0½		
Coal	0	10		
Coal and slate (slate floor).....	1	9	3	4½

**Gauley Coal Land Company Prospect No. A326—
No. 324 on Map II.**

On the southeast side of Job Knob Branch, 1.95 miles northeast of mouth; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3647' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	1	10

**Gauley Coal Land Company Prospect No. A325—
No. 325 on Map II.**

On the southeast side of Job Knob Branch, 1.55 miles northeast of mouth; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3638' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	2	11

**Gauley Coal Land Company Prospect No. A324—
No. 326 on Map II.**

On the southeast side of Job Knob Branch, 1.35 miles northeast of mouth; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3650' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	3	2½

**Gauley Coal Land Company Prospect No. A322—
No. 327 on Map II.**

On the southeast side of Job Knob Branch, 1.2 miles northeast of mouth; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3669' L.

	Ft.	In.
Coal	2	9

**Gauley Coal Land Company Prospect No. A321—
No. 328 on Map II.**

On the southeast side of Job Knob Branch, 1 mile northeast of mouth; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3669' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	3	8½

A sample was collected from the above section by the Gauley Coal Land Company and analyzed by the Commercial Testing and Engineering Company, of Charleston, W. Va. The analysis as reported by the former company is published under No. 328 in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. A320—
No. 329 on Map II.**

On the north side of Old Field Branch, 0.7 mile northeast of mouth; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3690' L.

			Ft.	In.
Bone and slate (slate roof).....	0'	7"		
Coal (fire clay floor).....	3	0	3	7

**Gauley Coal Land Company Prospect No. A319—
No. 330 on Map II.**

On the north side of Old Field Branch, 0.85 mile northeast of mouth; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3701' L.

	Ft.	In.
Coal (fire clay floor).....	3	1

A sample was taken from the above section by the Gauley Coal Land Company and analyzed by the Commercial Testing and Engineering Company, of Charleston, W. Va. The analysis as reported by the former company is published under No. 330 in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. A318—
No. 331 on Map II.**

On the north side of Old Field Branch, 1.1 miles northeast of mouth; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3704' L.

	Ft.	In.
Coal (fire clay floor).....	3	2

**Gauley Coal Land Company Prospect No. A317—
No. 332 on Map II.**

On the north side of Old Field Branch, 1.25 miles northeast of mouth; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3715' L.

	Ft.	In.
Coal (fire clay floor)	3	5

**Gauley Coal Land Company Prospect No. A316—
No. 333 on Map II.**

On the north side of Old Field Branch, 1.35 miles northeast of mouth; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3712' L.

		Ft.	In.
Coal, slate, and bone.....	0' 8"		
Coal (fire clay floor).....	2 10	3	6

**Gauley Coal Land Company Prospect No. A315—
No. 334 on Map II.**

On north side of Old Field Branch, 1.7 miles northeast of mouth; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3735' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	3	2

**Gauley Coal Land Company Prospect No. A232—
No. 335 on Map II.**

On the north side of Little Clear Creek Mountain, 0.5 mile southeast of mouth of Old Field Branch and 1.95 miles east of mouth of Smokehouse Branch; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3593' L.

			Ft.	In.
Coal (slate roof)	1'	4"		
Bone	0	1		
Coal	0	11½		
Coal, bony	0	1½		
Coal (slate floor)	1	1½	3	7½

**Gauley Coal Land Company Prospect No. A231—
No. 336 on Map II.**

On property of Graham Smokeless Coal Company; on the north side of Little Clear Creek Mountain, 0.75 mile southeast of mouth of Old Field Branch and 1.95 miles east of mouth of Smokehouse Branch; Gauley Coal Land Company authority for this section; **Fire Creek Coal**; elevation, 3634' L.

			Ft.	In.
Coal (slate roof)	1'	1½"		
Bone	0	2½		
Coal (slate floor).....	1	10½	3	2½

**Gauley Coal Land Company Prospect No. A230—
No. 337 on Map II.**

On north side of Little Clear Creek Mountain, 0.7 mile south of mouth of Old Field Branch and 1.75 miles southeast of mouth of Smokehouse Branch; Gauley Coal Land Company authority for this section; **Fire Creek Coal**; elevation, 3639' L.

	Ft.	In.
Coal, fallen shnt, exposed	2	0

**Gauley Coal Land Company Prospect No. A229—
No. 338 on Map II.**

On the north side of Little Clear Creek Mountain, 1.45 miles east of mouth of Smokehouse Branch and 0.6 mile southwest of mouth of Old Field Branch; Gauley Coal Land Company authority for this section; **Fire Creek Coal**; elevation, 3639' L.

	Ft.	In.
Coal (fire clay floor).....	3	9

**Gauley Coal Land Company Prospect No. A228—
No. 339 on Map II.**

On the north side of Little Clear Creek Mountain, 1.4 miles southeast of mouth of Smokehouse Branch and 3.95 miles northeast of Anjean; Gauley Coal Land Company authority for this section; **Fire Creek Coal**; elevation, 3658' L.

			Ft.	In.
Coal and slate (slate roof).....	0'	3"		
Coal	3	4		
Bone	0	0½		
Coal	1	1½		
Bone (sandstone floor)	0	2	4	11

**Gauley Coal Land Company Prospect No. A227—
No. 340 on Map II.**

On the north side of Joe Knob, 1.1 miles southeast of mouth of Smokehouse Branch and 3.2 miles east of Anjean; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3628' L.

			Ft.	In.
Coal	3'	4"		
Coal and slate	0	6		
Coal (slate floor)	1	10	5	8

**Gauley Coal Land Company Prospect No. A226—
No. 341 on Map II.**

On the north side of Little Clear Creek Mountain, 0.7 mile southeast of mouth of Smokehouse Branch and 3 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3550' L.

			Ft.	In.
Coal (slate roof)	1'	7"		
Slate	0	1½		
Coal	2	10½		
Bone	0	3		
Coal (fire clay floor)	1	5	6	3

**Gauley Coal Land Company Prospect No. A158—
No. 342 on Map II.**

On the east side of Briery Knob, 0.6 mile south of mouth of Smokehouse Branch and 2.9 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Fire Creek Coal?; elevation, 3520' L.

			Ft.	In.
Coal	1'	10"		
Bone	0	0½		
Coal (slate floor)	0	7½	2	6

**Gauley Coal Land Company Prospect No. A225—
No. 343 on Map II.**

On the north side of Briery Knob, 2.8 miles northeast of Anjean and 0.45 mile south of mouth of Smokehouse Branch; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3439' L.

			Ft.	In.
Bone	0'	2"		
Coal	1	8½		
Slate	0	1		
Coal	1	0½		
Bone	0	3		
Coal	0	11		
Slate	0	1½		
Coal (slate floor)	1	2	5	5½

Gauley Coal Land Company Prospect No. A224—**No. 344 on Map II.**

On the north side of Brlery Knob, 2.6 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3397' L.

			Ft.	In.
Coal (slate roof)	3'	3"		
Bone	0	3		
Coal	1	3		
Bone	0	3		
Fire clay	0	8		
Coal (slate floor)	1	11	7	7

Gauley Coal Land Company Prospect No. A223—**No. 345 on Map II.**

On the northwest side of Briery Knob, 0.6 mile northeast of mouth of Brlery Creek and 2.15 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3393' L.

			Ft.	In.
Coal	3'	6½"		
Slate	0	11		
Coal and slate	0	3		
Coal (fire clay floor).....	1	7	6	3½

Gauley Coal Land Company Prospect No. A222—**No. 346 on Map II.**

On the north side of Brlery Creek, 1.25 miles east of mouth and 2.75 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3527' L.

			Ft.	In.
Coal (slate roof).....	2'	11"		
Bone	0	4		
Coal	1	0		
Slate	0	2		
Coal	0	7		
Coal and bone (slate floor).....	0	5	5	5

Gauley Coal Land Company Prospect No. A221—**No. 347 on Map II.**

On the south side of Briery Creek, 1.3 miles southeast of mouth and 2.65 miles east of Anjean; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3618' L.

			Ft.	In.
Coal and slate (slate roof).....	0'	3"		
Coal	2	10½		
Bone	0	4		
Coal (fire clay floor).....	2	2	5	7½

**Gauley Coal Land Company Prospect No. A220—
No. 348 on Map II.**

On the south side of Briery Creek, 1.05 miles southeast of mouth and 2.35 miles east of Anjean; Gauley Coal Land Company authority for this section; **Fire Creek Coal**; elevation, 3553' L.

	Ft.	In.
Coal (slate roof; sandstone floor).....	6	0

**Gauley Coal Land Company Prospect No. A219—
No. 349 on Map II.**

On the south side of Briery Creek, 1.1 miles southeast of mouth and 2.3 miles east of Anjean; Gauley Coal Land Company authority for this section; **Fire Creek Coal**; elevation, 3605' L.

	Ft.	In.
Coal (fire clay floor)	4	4+

**Gauley Coal Land Company Prospect No. A218—
No. 350 on Map II.**

On the north side of Little Clear Creek Mountain, 1.8 miles east of Anjean and 0.7 mile southeast of mouth of Briery Creek; Gauley Coal Land Company authority for this section; **Fire Creek Coal**; elevation, 3493' L.

			Ft.	In.
Coal	3'	0"		
Slate	0	1½		
Coal (fire clay floor)	2	1½	5	3

**Gauley Coal Land Company Prospect No. A217—
No. 351 on Map II.**

On the north side of Little Clear Creek Mountain, 1.35 miles east of Anjean and 0.6 mile south of mouth of Briery Creek; Gauley Coal Land Company authority for this section; **Fire Creek Coal**; elevation, 3457' L.

	Ft.	In.
Coal (fire clay floor)	5	3

A sample was collected from the above section by the Gauley Coal Land Company and analyzed by the Commercial Testing and Engineering Company, of Charleston, W. Va. The analysis as reported by the former company is published under **No. 351** in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. A216—
No. 352 on Map II.**

On north side of Little Clear Creek Mountain, 0.9 mile southeast of Anjean and 1.1 miles southwest of mouth of Brlerly Creek; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3440' L.

		Ft.	In.
Coal and slate (slate roof).....	0' 10"		
Coal (fire clay floor).....	2 7	3	5

**Gauley Coal Land Company Prospect No. A215—
No. 353 on Map II.**

On the north side of Little Clear Creek Mountain, 1.15 miles southeast of Anjean and 1.3 miles southwest of mouth of Brlerly Creek; Gauley Coal Land Company authority for this section; Fire Creek Coal?; elevation, 3503' L.

	Ft.	In.
Coal	2	1

**Gauley Coal Land Company Prospect No. A214—
No. 354 on Map II.**

On the north side of Little Clear Creek Mountain, 0.65 mile southeast of Anjean and 1.4 miles southwest of mouth of Briery Creek; Gauley Coal Land Company authority for this section; Fire Creek Coal?; elevation, 3449' L.

	Ft.	In.
Coal	1	3

**Gauley Coal Land Company Prospect No. A213—
No. 355 on Map II.**

On the north side of Little Clear Creek Mountain, 0.85 mile southeast of Anjean and 1.6 miles southwest of the mouth of Brlerly Creek; Gauley Coal Land Company authority for this section; Fire Creek Coal?; elevation, 3468' L.

	Ft.	In.
Coal	1	3

**Gauley Coal Land Company Prospect No. A212—
No. 356 on Map II.**

On the south side of Little Clear Creek Mountain, 1.9 miles southeast of Anjean and 4.15 miles south of Duo; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3534' L.

	Ft.	In.
Coal (fire clay floor)	6	7

A sample was collected from the above section by the Gauley Coal Land Company and analyzed by the Commercial Testing and Engineering Company, of Charleston, W. Va. The analysis as reported by the former company is published under **No. 356** in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. A211—
No. 357 on Map II.**

On the south side of Little Clear Creek Mountain, west side of Hog Run, 2.1 miles southeast of Anjean; Gauley Coal Land Company authority for this section; **Fire Creek Coal**; elevation, 3591' L.

	Ft.	In.
Coal (fire clay floor)	5	1½

**Gauley Coal Land Company Prospect No. A210—
No. 358 on Map II.**

On the south side of Little Clear Creek Mountain, west side of Hog Run, 2.4 miles east of Anjean; Gauley Coal Land Company authority for this section; **Fire Creek Coal**; elevation, 3595' L.

	Ft.	In.
Coal (fire clay floor)	6	6

**Gauley Coal Land Company Prospect No. A209—
No. 359 on Map II.**

On the south side of Joe Knob, 3.4 miles east of Anjean; Gauley Coal Land Company authority for this section; **Fire Creek Coal**; elevation, 3695' L.

		Ft.	In.
Coal and slate (slate roof).....	0' 1½"		
Coal	2. 3		
Coal and bone	0 4		
Coal (slate floor).....	1 6	4	2½

A sample was collected from the above section and analyzed by the Commercial Testing and Engineering Company, of Charleston, W. Va. The analysis as reported by the former company is published under **No. 359** in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. A208—
No. 360 on Map II.**

On the south side of Little Clear Creek Mountain, 0.65 mile northeast of Joe Knob and 1.55 miles southeast of mouth of Smokehouse Branch; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3667' L.

			Ft.	In.
Coal (slate roof)	2'	0"		
Slate	0	0½		
Coal (sandstone floor)	2	5	4	5½

A sample was collected from the above section by the Gauley Coal Land Company and analyzed by the Commercial Testing and Engineering Company, of Charleston, W. Va. The analysis is published under No. 360 in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. A207—
No. 361 on Map II.**

On the property of the Graham Smokeless Coal Company; on south side of Little Clear Creek Mountain, 0.65 mile east of Joe Knob; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3683' L.

			Ft.	In.
Coal (slate roof)	0'	3"		
Parting	0	1		
Coal (fire clay floor)	4	2	4	6

**Gauley Coal Land Company Prospect No. A206—
No. 362 on Map II.**

On the property of Graham Smokeless Coal Company; on the east side of Kuhn Branch, 1.9 miles north of mouth, and 4.3 miles east of Anjean; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3645' L.

			Ft.	In.
Coal and slate (slate roof).....	9'	4"		
Coal	4	9	5	1
Slate (fire clay floor)			0	8

**Gauley Coal Land Company Prospect No. A205—
No. 363 on Map II.**

On property of Graham Smokeless Coal Company; on east side of Polut Mountain, 1.8 miles northeast of mouth of Kuhu Branch; Gauley

Coal Land Company authority for this section; Fire Creek Coal; elevation, 3685' L.

		Ft.	In.
Coal (slate roof)	0' 3½"		
Slate	0 1½		
Coal (slate floor)	4 11½	5	4½

Gauley Coal Land Company Prospect No. A203— No. 364 on Map II.

On property of Graham Smokeless Coal Company; on east side of Little Clear Creek near head, 1.9 miles southeast of Old Field Branch, 5.2 miles east of Anjean; Gauley Coal Land Company authority for this section; Fire Creek Coal; elevation, 3735' L.

		Ft.	In.
Coal (slate roof).....	2' 7"		
Slate and bone	0 5½		
Coal (fire clay floor).....	1 5	4	5½

Prospect—No. 365 on Map II.

On head of Stony Run, 4.55 miles east of Rupert and 2.1 miles north of Kleffer; Fire Creek Coal?; elevation, 3565' B.

	Ft.	In.
Coal, (reported by B. M. Higginbotham).....	3	0

Deats Mine (Abandoned)—No. 366 on Map II.

Farm mine, on west side of Cross Mountain, 4.75 miles east of Rupert and 1.75 miles north of Kleffer; Fire Creek Coal?; elevation, 3610' B.

	Ft.	In.
Coal, (reported by B. M. Higginbotham)	3	6

Fire Creek Coal, Williamsburg District.

An area of 22 square miles in this district should contain the Fire Creek Coal. It is believed that this coal is of minable thickness over much if not all of this area and that further prospecting is warranted. The following prospects and openings were noted:

Mine (Abandoned)—No. 367 on Map II.

Farm mine, on east side of Buffalo Mountain, 3.5 miles west of Williamsburg; Fire Creek Coal?; elevation, 3810' B.

	Ft.	In.
Coal, (opening fallen shut, thickness reported).....	2'	11

**Gauley Coal Land Company Prospect No. 150—
No. 368 on Map II.**

On the east end of Little Clear Creek Mountain, 0.15 mile southeast of Drill hole No. 14; Gauley Coal Land Company authority for this section; **Fire Creek Coal?**; elevation, 4034' L.

	Ft.	In.
Coal	1	0

Prospect—No. 369 on Map II.

On northeast side of Laurel Creek, 3.6 miles northeast of Beech Knob and 2.3 miles southwest of Baber School; **Fire Creek Coal**; elevation, 3261'.

	Ft.	In.
Coal, reported to be dirty with a thickness of.....	4	2

Fire Creek Coal, Falling Springs District.

An area of 36.5 square miles in this district should contain **Fire Creek Coal**. It is believed that this coal is of minable thickness over much if not all of this area and that further prospecting is warranted. The following prospect and exposures were noted:

**Cherry River Boom and Lumber Company? Prospect—
No. 370 on Map II.**

Located 0.75 mile southeast of Manning Knob and 5 miles northeast of Clearco; **Fire Creek Coal**; elevation, 3675' B.

	Ft.	In.
Shale, black, pelecypods		
Coal (fallen shut, reported).....	3	2

The above opening had fallen shut but a sample of coal was collected from the dump. The analysis of this sample (No. 162PH) is published under **No. 370** in the Table of Coal Analyses at the end of this Chapter.

Coal Exposure—No. 370A on Map II.

On Cold Knob road, 0.4 mile southeast of Manning Knob and 4 miles south of Richwood; **Fire Creek Coal**; elevation, 3375' B.

	Ft.	In.
Coal	0'	6"
Fire clay	5	0
Coal	0	6
	6	0

tain whether it is a true split off the Beekley Coal or a separate lenticular coal bed. This lower seam is 10 to 40 feet below the Beekley Coal and shows a variable but sometimes good section of coal. Descriptions of the Beekley and "Split" seam prospects on Joe Knob follow:

**Gauley Coal Land Company Prospect No. A116—
No. 283 on Map II.**

On the northeast side of Joe Knob, 3.25 miles east of Anjean; Gauley Coal Land Company authority for this section; Beekley Coal; elevation, 3718' L.

Coal (slate floor).....	Ft.	In
	3	

**Gauley Coal Land Company Prospect No. A115—
No. 284 on Map II.**

On the north side of Joe Knob, 3.1 miles east of Anjean; Gauley Coal Land Company authority for this section; Beekley Coal; elevation 3714' L.

Coal and slate (slate floor).....	1'	5"	Ft.
Coal (slate floor)	3	2	4

The "Split" seam has been opened immediately below No. 284 and shows the following section:

**Gauley Coal Land Company Prospect No. 157—
Not Shown on Map II.**

On the north side of Joe Knob, 3.1 miles east of Anjean; Gauley Coal Land Company authority for this section; "Split" Coal elevation 3688' L.

Bone (slate roof)	0'	5"	Ft.
Slate	0	0½	
Coal (sandstone floor)	4	4	

**Gauley Coal Land Company Prospect No. A11
No. 285 on Map II.**

On west side of Joe Knob, 2.85 miles east of Anjean; Gauley Coal Land Company authority for this section; Beekley Coal

The "Split" seam has been opened immediately below No. 285 and shows the following:

**Gauley Coal Land Company Prospect No. 154—
Not Shown on Map II.**

On west side of Joe Knob, 2.85 miles east of Anjean; Gauley Coal Land Company authority for this section; "Split" Coal; elevation, 3658' L.

	Ft.	In.
Coal (no top; sandstone floor).....	2	6½

**Gauley Coal Land Company Prospect No. A109—
No. 286 on Map II.**

On the southwest side of Joe Knob, 2.9 miles east of Anjean; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3694' L.

		Ft.	In.
Coal	1' 11"		
Slate	0 0½		
Coal	0 11		
Slate	0 0½		
Coal	2 6½		
Coal and slate (fire clay floor).. <hr/>	0 3½	5	9

The "Split" seam has been opened 0.2 mile southeast of No. 286 and shows the following:

**Gauley Coal Land Company Prospect No. 153—
Not Shown on Map II.**

On south side of Joe Knob, 3.1 miles east of Anjean; Gauley Coal Land Company authority for this section; "Split" Coal; elevation, 3696' L.

	Ft.	In.
Coal (no top; fire clay floor).....	2	6

**Gauley Coal Land Company Prospect No. A108—
No. 287 on Map II.**

On the south side of Joe Knob, 3.3 miles east of Anjean; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3771' L.

		Ft.	In.
Coal	2' 10"		
Coal and slate	0 3		
Coal (slate floor)	1 8	4	9

tain whether it is a true split off the Beekley Coal or a separate lenticular coal bed. This lower seam is 10 to 40 feet below the Beekley Coal and shows a variable but sometimes good section of coal. Descriptions of the Beekley and "Split" seam prospects on Joe Knob follow:

**Gauley Coal Land Company Prospect No. A116—
No. 283 on Map II.**

On the northeast side of Joe Knob, 3.25 miles east of Anjean; Gauley Coal Land Company authority for this section; Beekley Coal; elevation, 3718' L.

	Ft.	In.
Coal (slate floor).....	3	7

**Gauley Coal Land Company Prospect No. A115—
No. 284 on Map II.**

On the north side of Joe Knob, 3.1 miles east of Anjean; Gauley Coal Land Company authority for this section; Beekley Coal; elevation 3714' L.

		Ft.	In.
Coal and slate (slate floor).....	1' 5"		
Coal (slate floor)	3 2	4	7

The "Split" seam has been opened immediately below No. 284 and shows the following section:

**Gauley Coal Land Company Prospect No. 157—
Not Shown on Map II.**

On the north side of Joe Knob, 3.1 miles east of Anjean; Gauley Coal Land Company authority for this section; "Split" Coal elevation, 3688' L.

		Ft.	In.
Bone (slate roof)	0' 5"		
Slate	0 0½		
Coal (sandstone floor)	4 4	4	9½

**Gauley Coal Land Company Prospect No. A114—
No. 285 on Map II.**

On west side of Joe Knob, 2.85 miles east of Anjean; Gauley Coal Land Company authority for this section; Beekley Coal; elevation, 3671' L.

	Ft.	In.
Coal (slate floor).....	5	6½

COMMERCIAL COAL.

Gauley Coal Land Company Prospect No. A119—
No. 288 on Map II.

On the east side of Briery Knob, 2.8 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3578' L.

Bone (slate roof)	0'	93"
Coal (fire clay floor)	2	9

Fl. In. 64

Gauley Coal Land Company Prospect No. A118—
No. 289 on Map II.

On the north side of Briery Knob, 2.6 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3530' L.

A sample was collected from the above section by Gauley Coal Land Company and analyzed by the Commercial Testing and Engineering Company, of Charleston, W. The analysis as reported by the former company is published under No. 289 in the Table of Coal Analyses at the end of this chapter.

Gauley Coal Land Company Prospect No. A117—
No. 290 on Map II.

On the northwest side of Briery Knob, 2.5 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3539' L.

0'	6"
0	10
0	8
0	2
1	1
0	10
1	2

Coal (fire clay floor)

Gauley Coal Land Company Prospect No. A113—
No. 291 on Map II.

On the north side of Little Clear Creek Mountain, 1.8 miles west of Joe Knob; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3568' L.

**Gauley Coal Land Company Prospect No. A119—
No. 288 on Map II.**

On the east side of Briery Knob, 2.8 miles northeast of Anjean;
Gauley Coal Land Company authority for this section; Beckley Coal;
elevation, 3578' L.

			Ft.	In.
Bone (slate roof)	0'	9½"		
Coal (fire clay floor)	2	9	3	6½

**Gauley Coal Land Company Prospect No. A118—
No. 289 on Map II.**

On the north side of Briery Knob, 2.6 miles northeast of Anjean;
Gauley Coal Land Company authority for this section; Beckley Coal;
elevation, 3530' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	3	8½

A sample was collected from the above section by the
Gauley Coal Land Company and analyzed by the Commercial
Testing and Engineering Company, of Charleston, W. Va.
The analysis as reported by the former company is published
under **No. 289** in the Table of Coal Analyses at the end of
this chapter.

**Gauley Coal Land Company Prospect No. A117—
No. 290 on Map II.**

On the northwest side of Briery Knob, 2.5 miles northeast of
Anjean; Gauley Coal Land Company authority for this section; Beck-
ley Coal; elevation, 3539' L.

			Ft.	In.
Slate and bone (slate roof).....	0'	6"		
Coal	0	10		
Slate	0	8		
Slate	1	2		
Coal	0	10		
Slate and coal	1	2	5	2
Coal (fire clay floor).....				

**Gauley Coal Land Company Prospect No. A113—
No. 291 on Map II.**

On the north side of Little Clear Creek Mountain, 1.8 miles east of
Anjean and 1.35 miles west of Joe Knob; Gauley Coal Land Company
authority for this section; Beckley Coal?; elevation, 3568' L.

	Ft.	In.
Coal (slate roof)	0	10

Gauley Coal Land Company Prospect No. A119—
No. 288 on Map II.

On the east side of Briery Knob, 2.8 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3578' L.

Bone (slate roof)	0'	9 3/4"	2	9	3	6 1/2
Coal (fire clay floor)							

Gauley Coal Land Company Prospect No. A118—
No. 289 on Map II.

On the north side of Briery Knob, 2.6 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3530' L.

Coal (slate roof; fire clay floor)	3	8 1/2
--	---	-------

A sample was collected from the above section by the Gauley Coal Land Company and analyzed by the Commercial Testing and Engineering Company, of Charleston, W. Va. The analysis as reported by the former company is published under No. 289 in the Table of Coal Analyses at the end of this chapter.

Gauley Coal Land Company Prospect No. A117—
No. 290 on Map II.

On the northwest side of Briery Knob, 2.5 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3539' L.

Slate and bone (slate roof)	0'	6"	0	10	8	1	2	5
Coal									
Slate									
Coal									
Slate and coal									
Coal (fire clay floor)									

Gauley Coal Land Company Prospect No. A113—
No. 291 on Map II.

On the north side of Little Clear Creek Mountain, 1.8 miles east of Anjean and 1.35 miles west of Joe Knob; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3568' L.

Coal (slate roof)

**Gauley Coal Land Company Prospect No. A119—
No. 288 on Map II.**

On the east side of Briery Knob, 2.8 miles northeast of Anjean;
Gauley Coal Land Company authority for this section; Beckley Coal;
elevation, 3578' L.

			Ft.	In.
Bone (slate roof)	0'	9½"		
Coal (fire clay floor)	2	9	3	6½

**Gauley Coal Land Company Prospect No. A118—
No. 289 on Map II.**

On the north side of Briery Knob, 2.6 miles northeast of Anjean;
Gauley Coal Land Company authority for this section; Beckley Coal;
elevation, 3530' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	3	8½

A sample was collected from the above section by the
Gauley Coal Land Company and analyzed by the Commercial
Testing and Engineering Company, of Charleston, W. Va.
The analysis as reported by the former company is published
under No. 289 in the Table of Coal Analyses at the end of
this chapter.

**Gauley Coal Land Company Prospect No. A117—
No. 290 on Map II.**

On the northwest side of Briery Knob, 2.5 miles northeast of
Anjean; Gauley Coal Land Company authority for this section; Beck-
ley Coal; elevation, 3539' L.

			Ft.	In.
Slate and bone (slate roof).....	0'	6"		
Coal	0	10		
Slate	0	8		
Coal	1	2		
Slate and coal	0	10		
Coal (fire clay floor).....	1	2	5	2

**Gauley Coal Land Company Prospect No. A113—
No. 291 on Map II.**

On the north side of Little Clear Creek Mountain, 1.8 miles east of
Anjean and 1.35 miles west of Joe Knob; Gauley Coal Land Company
authority for this section; Beckley Coal?; elevation, 3568' L.

	Ft.	In.
Coal (slate roof)	0	10

**Gauley Coal Land Company Prospect No. A112—
No. 292 on Map II.**

On the north side of Little Clear Creek Mountain, 1.35 miles east of Anjean and 1.8 miles west of Joe Kuob; Gauley Coal Land Company authority for this section; Beckley Coal?; elevation, 3534' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	0	10

**Gauley Coal Land Company Prospect No. A111—
No. 293 on Map II.**

On north side of Little Clear Creek Mountain, 1.35 miles southeast of Anjean and 2.1 miles southwest of Joe Knob; Gauley Coal Land Company authority for this section; Beckley Coal?; elevation, 3556' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	1	5½

**Gauley Coal Land Company Prospect No. A110—
No. 294 on Map II.**

On the south side of Little Clear Creek Mountain, 1.9 miles southeast of Anjean and 1.4 miles southwest of Joe Knob; Gauley Coal Land Company authority for this section; Beckley Coal?; elevation, 3612' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	1	10

**Gauley Coal Land Company Prospect No. A107—
No. 295 on Map II.**

On the south side of Little Clear Creek Mountain, 3.8 miles east of Anjean and 3.8 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3746' L.

	Ft.	In.
Coal (slate top; slate floor)	2	10

**Gauley Coal Land Company Prospect No. A106—
No. 296 on Map II.**

On Graham Smokeless Coal Company property; on the south side of Little Clear Creek Mountain, 4.1 miles east of Anjean and 4.2 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3737' L.

	Ft.	In.
Coal (slate top; fire clay floor)	3	2

Gauley Coal Land Company Prospect No. A112—
No. 292 on Map II.

On the north side of Little Clear Creek Mountain, 1.35 miles east of Aujean and 1.8 miles west of Joe Knob; Gauley Coal Land Company authority for this section; Beckley Coal?; elevation, 3534' L.
Coal (slate roof; fire clay floor)..... 0
Ft. In. 10

Gauley Coal Land Company Prospect No. A111—
No. 293 on Map II.

On north side of Little Clear Creek Mountain, 1.35 miles southeast of Aujean and 2.1 miles southwest of Joe Knob; Gauley Coal Land Company authority for this section; Beckley Coal?; elevation, 3556' L.
Coal (slate roof; fire clay floor)..... 1
Ft. In. 54

Gauley Coal Land Company Prospect No. A110—
No. 294 on Map II.

On the south side of Little Clear Creek Mountain, 1.9 miles south-east of Aujean and 1.4 miles southwest of Joe Knob; Gauley Coal Land Company authority for this section; Beckley Coal?; elevation, 3612' L.
Coal (slate roof; fire clay floor)..... 1
Ft. In. 10

Gauley Coal Land Company Prospect No. A107—
No. 295 on Map II.

On the south side of Little Clear Creek Mountain, 3.8 miles east of Aujean and 3.8 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3746' L.
Coal (slate top; slate floor)..... 2
Ft. In. 10

Gauley Coal Land Company Prospect No. A106—
No. 296 on Map II.

On Graham Smokeless Coal Company property; on the south side of Little Clear Creek Mountain, 4.1 miles east of Aujean and 4.2 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3737' L.
Coal (slate top; fire clay floor)..... 3
Ft. In. 2

**Gauley Coal Land Company Prospect No. A112—
No. 292 on Map II.**

On the north side of Little Clear Creek Mountain, 1.35 miles east of Anjean and 1.8 miles west of Joe Knob; Gauley Coal Land Company authority for this section; Beckley Coal?; elevation, 3534' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	0	10

**Gauley Coal Land Company Prospect No. A111—
No. 293 on Map II.**

On north side of Little Clear Creek Mountain, 1.35 miles southeast of Anjean and 2.1 miles southwest of Joe Knob; Gauley Coal Land Company authority for this section; Beckley Coal?; elevation, 3556' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	1	5½

**Gauley Coal Land Company Prospect No. A110—
No. 294 on Map II.**

On the south side of Little Clear Creek Mountain, 1.9 miles southeast of Anjean and 1.4 miles southwest of Joe Knob; Gauley Coal Land Company authority for this section; Beckley Coal?; elevation, 3612' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	1	10

**Gauley Coal Land Company Prospect No. A107—
No. 295 on Map II.**

On the south side of Little Clear Creek Mountain, 3.8 miles east of Anjean and 3.8 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3746' L.

	Ft.	In.
Coal (slate top; slate floor)	2	10

**Gauley Coal Land Company Prospect No. A106—
No. 296 on Map II.**

On Graham Smokeless Coal Company property; on the south side of Little Clear Creek Mountain, 4.1 miles east of Anjean and 4.2 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3737' L.

	Ft.	In.
Coal (slate top; fire clay floor)	3	2

**Gauley Coal Land Company Prospect No. A105—
No. 297 on Map II.**

On Graham Smokeless Coal Company property; on the south side of Little Clear Creek Mountain, 4.3 miles east of Anjean and 3.9 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3727' L.

	Ft.	In.
Coal (slate top; fire clay floor).....	3	2

**Gauley Coal Land Company Prospect No. A104—
No. 298 on Map II.**

On Graham Smokeless Coal Company property; on the south side of Little Clear Creek Mountain, 4.65 miles east of Anjean and 4.35 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3748' L.

	Ft.	In.
Coal	3	5

**Gauley Coal Land Company Prospect No. A102—
No. 299 on Map II.**

On the head of Little Clear Creek on the south side of Little Clear Creek Mountain, 6 miles east of Anjean and 4.5 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal?; elevation, 3904' L.

	Ft.	In.
Coal (slate roof)	2	1

**Gauley Coal Land Company Prospect No. A101—
No. 300 on Map II.**

On the east side of Little Clear Creek near its source, 5.65 miles east of Anjean and 4.6 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3865' L.

	Ft.	In.
Coal (slate roof and floor)	3	3

**Gauley Coal Land Company Prospect No. A100—
No. 301 on Map II.**

On the east side of Little Clear Creek near its source, 5.55 miles east of Anjean and 4.55 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3855' L.

	Ft.	In.
Coal (slate roof and floor)	3	2½

Beckley Coal, Williamsburg District.

Only one farm mine and three prospects in the Beckley Coal were noted in this district. The horizon of this coal is present over a larger area in this district than is shown as minable on Figure 19, page 531, and warrants further prospecting.

Gauley Coal Land Company Prospect No. 151— No. 302 on Map II.

On the head of Flynn Creek, 2.9 miles northwest of Trout P. O.; Gauley Coal Land Company authority for this section; Beckley Coal?; elevation, 4040' L.

	Ft.	In.
Coal	0	10

Gauley Coal Land Company Prospect—No. 303 on Map II.

On the waters of Hogcamp Run, 1.2 miles southwest from its mouth and 2.1 miles east of Beech Knob; Ganley Coal Land Company authority for this section; Beckley Coal; elevation, 3273' L.

		Ft.	In.
Coal	0' 6"		
Slate	1 0		
Coal	1 0		
Slate	0 4		
Coal	0 6	3	4

Gauley Coal Land Company Prospect No. 465— No. 304 on Map II.

1.55 miles northeast of Lile and 1.5 miles northeast of Beech Knob; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3223' L.

	Ft.	In.
Coal	3	0

Laurel Manufacturing Company Mine—No. 305 on Map II.

Farm mine, on the south side of McMillion Creek, 1.7 miles northeast of Lile; observation by Reger; Beckley Coal; elevation, 3000' B.

		Ft.	In.
Coal, bony (slate roof).....	0' 11"		
Slate, bony	0 4		
Coal, soft (slate floor)	2 0	3	3

This opening was described by Reger under No. 1276 in the Nicholas County Report, page 715, as occurring in the Fire Creek seam, but after tracing this coal across Greenbrier it apparently proves to be at the Beckley horizon.

Beckley Coal, Falling Springs District.

With the exception of one reported opening on Buffalo Mountain, the two abandoned mine openings and the prospect near them on Lost Flat are the only points at which the Beckley Coal was noted in this district. As shown on the Lost Flat Mine map the thickness of the coal ranged from one and one-half to six and one-half feet. The coal was locally absent in one part of the mine and from the reported circumstances it was probably cut out by the overlying Lower Raleigh Sandstone. It is probable that this cut-out was local and if this is true it would appear that further prospecting at this horizon in the district would be profitable.

Elk Lick Coal Company Prospect—No. 306 on Map II.

On northeast side of Lost Flat, 1.9 miles northeast of Manning Knob and 5.7 miles southeast of Richwood; Beckley Coal; elevation, 3600' B.

	Ft.	In.
Coal	1	6

Elk Lick Coal Company "Lost Flat" Mine (Abandoned)— No. 307 on Map II.

On the east side of Lost Flat, 2.35 miles northeast of Manning Knob and 6 miles southeast of Richwood; Beckley Coal; elevation, 3640' B.

	Ft.	In.
Coal	3	7

Elk Lick Coal Company "Old Lost Flat" Mine (Abandoned)— No. 308 on Map II.

On southeast end of Lost Flat, 2.1 miles northeast of Manning Knob and 6.1 miles southeast of Richwood; Beckley Coal; elevation, 3650' B.

	Ft.	In.
Coal	4	0

Prospect—No. 309 on Map II.

On east side of Buffalo Mountain, 3.4 miles west of Williamsburg;
Beckley Coal?; elevation, 3865' B.

	Ft.	In.
Coal, (reported by B. M. Higginbotham), 0' 10" to.....	1	0

Quantity of Beckley Coal Available.

The following table, computed from planimetric measurement of outcrop outlined on work sheets for the area indicated on Figure 19, page 531, gives the probable amount of Beekley Coal in Greenbrier County. The assumed thicknesses of coal shown in the table are **average** thicknesses and should not be used in any tabulation of coal reserves by thicknesses. A low **average** figure was used so that the total tonnage would not be too great if local areas prove to be cut out or too thin for mining:

Probable Amount of Beckley Coal.

District.	Thickness of Coal Assumed. Feet.	Square Miles.	Acres.	Cubic Feet of Coal.	Short Tons of Coal (2000 Lbs.)
Meadow Bluff	2	35.5	22,720	1,979,366,400	79,174,656
Williamsburg	2	11.6	7,424	646,778,880	25,871,155
Falling Springs	3	1.0	640	83,635,200	3,345,408
Totals		48.1	30,784	2,709,780,480	108,391,219

Figures of the Department of Mines show that 119,522 tons of coal have been produced from mines operating the Beekley seam in Greenbrier County, all of which came from the Lost Flat Mine in Falling Springs District.

FIRE CREEK COAL.

The Fire Creek Coal, previously described in Chapter VI, page 238, ranks first in available coal in Greenbrier County. In general it varies from 1 to 6 feet in thickness and in comparatively local areas it may be entirely cut out by the overlying Quinnimont Sandstone. The ash content of the Fire Creek Coal appears to be somewhat higher than the Sewell Coal but in other respects it compares favorably with the latter coal. The probable minable area of the Fire Creek Coal is shown on Figure 20, and its detailed outcrop is outlined in blue on Map II.

FIGURE 20

MAP


OF

GREENBRIER COUNTY

SHOWING

PROBABLE AREA OF MINABLE

FIRE CREEK COAL



LITTLE RALEIGH COAL.

The Little Raleigh Coal, previously discussed in Chapter VI, pages 235-6, is generally persistent throughout a considerable part of Greenbrier County. In general it ranges in thickness from 2 to 3 feet but scattered sections show a thickness of 4 feet of clean coal. Chemical analyses indicate that the ash content is fairly high but it is probable that some of the openings sampled were not driven in far enough to reach the best coal. This coal has been mined at several points for local use but at present (1936) there are no actively operating mines in the area. In appearance and rank the Little Raleigh Coal is quite similar to the other Pottsville coals of the region.

Greenbrier County is the only county in the State in which the Little Raleigh Coal is known to be of minable thickness. Figure 19, page 531, shows the probable area of minable Little Raleigh Coal but its outcrop is not outlined on Map II. The position of the outcrop of this seam can be easily found by reference to the Sewell Coal structure contour lines since it is generally 130 to 160 feet below the Sewell horizon.

FIGURE 19

MAP

OF


GREENBRIER COUNTY

SHOWING

PROBABLE AREA OF MINABLE

LITTLE RALEIGH COAL

BECKLEY COAL



Little Raleigh Coal, Meadow Bluff Distriet.

In Meadow Bluff Distriet the Little Raleigh Coal is noted in coal test borings Nos. 5, 5A, 5C, 5H, 5I, 5K, 5M, 6, and 11, the details of which are published on preceding pages. The prospects and openings noted are as follows:

Gauley Coal Land Company Prospect No. 8— No. 231 on Map II.

On west side of Mill Creek Mountain, 1.85 miles northeast of Charmco and 3.9 miles northwest of Rupert; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3123' L.

	Ft.	In.
Coal (slate roof and floor)	2	1

Coal Exposure No. 231A of the Little Raleigh Coal is published in the Charmco Section, page 164, where a thickness of one foot was noted at an elevation of 2925' B.

Gauley Coal Land Company Prospect No. 7— No. 232 on Map II.

On Mill Creek Mountain, 1.95 miles northeast of Charmco and 3.35 miles northwest of Rupert; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3193' L.

	Ft.	In.
Coal (slate floor)	2	6

Gauley Coal Land Company Prospect No. 6— No. 233 on Map II.

On Mill Creek Mountain, 1.8 miles northeast of Charmco and 3.15 miles northwest of Rupert; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3193' L.

	Ft.	In.
Coal (slate floor)	2	6

Gauley Coal Land Company Prospect No. 5— No. 234 on Map II.

On Mill Creek Mountain, 1.05 miles east of Charmco; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3127.7' L.

			Ft.	In.
Coal (slate roof)	2'	3½"		
Coal, bony (slate floor).....	0	3	2	6½

**Gauley Coal Land Company Prospect No. 4—
No. 235 on Map II.**

On Mill Creek Mountain, 1.8 miles east of Charmco and 3 miles northwest of Rupert; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3213' L.

	Ft.	In.
Coal (slate roof and floor).....	2	7

The following is an analysis made by the Commercial Testing and Engineering Company, of Charleston, W. Va., of a sample collected from above prospect by the Gauley Coal Land Company, as reported under Laboratory No. 86186 by the latter company:

	As Received. Per cent.	Dry Basls. Per cent.
Molsture	1.40
Volatile Matter	24.04	24.38
Fixed Carbon	66.70	67.65
Ash	7.86	7.97
	<hr/> 100.00	<hr/> 100.00
Sulphur	1.68	1.78
B. T. U.	14,093	14,293

Leslie Hines Mine (Abandoned)—No. 236 on Map II.

Gauley Coal Land Company Prospect No. 9, on Mill Creek Mountain, 1.7 miles east of Charmco and 2.75 miles northwest of Rupert; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3218' L.

	Ft.	In.
Coal	3	2

The following is an analysis made by the Commercial Testing and Engineering Company, of Charleston, W. Va., of a sample collected from the above opening by the Gauley Coal Land Company, as reported under Laboratory No. 86187 by the latter company:

	As Received. Per cent.	Dry Basls. Per cent.
Molsture	1.51
Volatile Matter	22.09	22.43
Fixed Carbon	67.21	68.24
Ash	9.19	9.33
	<hr/> 100.00	<hr/> 100.00
Sulphur	1.39	1.41
B. T. U.	13,830	14,042

**Gauley Coal Land Company Prospect No. 11—
No. 237 on Map II.**

On south end of Mill Creek Mountain, 1.6 miles east of Charmco and 2.65 miles northwest of Rupert; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3182' L.

	Ft.	In.
Coal	3	0

**Gauley Coal Land Company Prospect No. 10—
No. 238 on Map II.**

On Mill Creek Mountain, 1.75 miles east of Charmco and 2.6 miles northwest of Rupert; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3232' L.

	Ft.	In.
Coal (slate roof and floor).....	3	2

**Gauley Coal Land Company Prospect No. 3—
No. 239 on Map II.**

On east side of Mill Creek Mountain, 0.65 mile south of Big Branch School; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3218' L.

	Ft.	In.
Coal (slate roof and floor)	2	10

**Gauley Coal Land Company Prospect No. 2—
No. 240 on Map II.**

On east side of Mill Creek Mountain, 0.25 mile southeast of Big Branch School; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3193' L.

	Ft.	In.
Coal (slate roof and floor)	2	8

The following is an analysis made by the Commercial Testing and Engineering Company, of Charleston, W. Va., of a sample collected from the above opening by the Gauley Coal Land Company, as reported under Laboratory No. 86185 by the latter company:

	As Received. Per cent.	Dry Basis. Per cent.
Molsture	3.09
Volatile Matter	22.83	23.56
Fixed Carbon	63.98	66.02
Ash	10.10	10.42
	<hr/>	<hr/>
	100.00	100.00
Sulphur	1.32	1.36
B. T. U.	12,948	13,361

**Gauley Coal Land Company Prospect No. 1—
No. 241 on Map II.**

On east side of Mill Creek Mountain, 0.2 mile northeast of Big Brauch School; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3174' L.

	Ft.	In.
Coal (slate roof and floor).....	2	1

**Gauley Coal Land Company Prospect No. 504—
No. 242 on Map II.**

On the north bank of North Fork of Big Clear Creek, 2 miles southwest of Clearco and 1.1 miles north of Duo; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3290' L.

	Ft.	In.
Coal	2	10

**Gauley Coal Land Company Prospect No. 503—
No. 243 on Map II.**

On the south side of Beech Ridge, 1.7 miles southwest of Clearco; and 1.3 miles north of Duo; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3282' L.

		Ft.	In.
Coal	1' 3"		
Slate	0 2		
Coal	1 7	3	0

**Gauley Coal Land Company Prospect No. 502—
No. 244 on Map II.**

0.9 mile northwest of Job Knob and 0.8 mile southeast of Clearco; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3510' L(?).

	Ft.	In.
Coal	3	6

**Gauley Coal Land Company Prospect No. 6—
No. 245 on Map II.**

On the east side of Smokehouse Branch, 1.85 miles northeast of its mouth and 1.5 miles southeast of Duo; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3555' L.

		Ft.	In.
Coal	1' 2"		
Coal and slate (slate floor).....	1 8½	2	10½

**Gauley Coal Land Company Prospect No. 5—
No. 246 on Map II.**

On the east side of Smokehouse Branch, 1.5 miles northeast of its mouth and 1.7 miles southeast of Duo; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3637' L.

	Ft.	In.
Coal (slate roof and floor).....	1	11

**Gauley Coal Land Company Prospect No. 3—
No. 247 on Map II.**

On the west side of Smokehouse Ridge, 1.4 miles northeast of mouth of Smokehouse Branch; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3639' L.

	Ft.	In.
Coal (slate floor)	2	11

**Gauley Coal Land Company Prospect No. 19—
No. 248 on Map II.**

On the east side of Smokehouse Ridge, 0.25 mile northwest of the mouth of Oldhouse Branch and 2.3 miles southeast of Duo; Gauley Coal Land Company authority for this section; Little Raleigh Coal?; elevation, 3716' L.

	Ft.	In.
Coal (slate floor)	2	0

Prospect—No. 249 on Map II.

On southwest end of Hickory Ridge, 3.85 miles northeast of Rupert and 2.45 miles south of Anjean; Little Raleigh Coal?; elevation, 3630' B.

	Ft.	In.
Coal, (reported by Golden Jones), 3' to	4	0

Prospect—No. 250 on Map II.

On northeast end of Hickory Ridge, 5.2 miles northeast of Rupert and 2.9 miles southeast of Anjean; Little Raleigh Coal?; elevation, 3930' B.

	Ft.	In.
Coal, (reported by B. M. Higginbotham)	4	0

Prospect—No. 251 on Map II.

On west end of Long Point, 3.9 miles east of Rupert; Little Raleigh Coal?; elevation, 3645' B.

	Ft.	In.
Coal, (reported by B. M. Higginbotham)	4	0

Prospect—No. 252 on Map II.

On west side of Buffalo Mountain, 3.65 miles west of Williamsburg; Little Raleigh Coal; elevation, 3930' B.

	Ft.	In.
Coal, (reported by B. M. Higginbotham).....	2	4

Little Raleigh Coal, Williamsburg District.

Very little prospecting for Little Raleigh Coal has been done in Williamsburg District. The sections shown in the following prospects and openings indicate that further prospecting is highly desirable:

**Gauley Coal Land Company Prospect No. 466—
No. 253 on Map II.**

On the northeast side of Beech Ridge, 1.55 miles east of Cleareo; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3517' L.

		Ft.	In.
Coal	4' 2"		
Slate	3 0		
Coal	0 5	7	7

**Gauley Coal Land Company Prospect No. 465B—
No. 254 on Map II.**

On the waters of Hogcamp Run, 1.25 miles southwest of its mouth and 2.1 miles east of Beech Knob; Gauley Coal Land Company authority for this section; Little Raleigh Coal; elevation, 3386' L.

		Ft.	In.
Coal	3' 10"		
Slate	1 3		
Coal	0 8	5	9

T. B. Lilly Mine—No. 255 on Map II.

Farm mine, on the north side of McMillion Creek, 2.4 miles south 9° east of Eureka School and 1.1 miles northeast of Lille; observation by Reger; Little Raleigh Coal; elevation, 3040' B.

	Ft.	In.
Fallen shut, coal reported 3' 0" to.....	4	0

The above opening was described by Reger under No. 1264 in the Nicholas County Report, page 344, as occurring in the Beckley seam, but after tracing the coal across Greenbrier it apparently proves to be at the Little Raleigh horizon.

Levi Lilly Heirs Mine—No. 256 on Map II.

Farm mine, on north bank of McMillion Creek, 2.1 miles south 30° east of Eureka School and 1.6 miles northeast of Lille; observation by Reger; **Little Raleigh Coal**; elevation, 3015' B.

			Ft.	In.
Sandstone, massive				
Coal, soft	1'	4"		
Coal, bony	0	8	2	0
Slate, bony			2	0

The above opening was described by Reger under No. 1263 in the Nicholas County Report, page 344, as occurring in the Beekley seam, but after tracing this coal across Greenbrier it apparently proves to be at the Little Raleigh horizon.

Gauley Coal Land Company (?) Mine—No. 257 on Map II.

On the Jetsville-Manning Knob road, 4.4 miles northeast of Lille and 4 miles north of Cleareo; **Little Raleigh Coal**; elevation, 3485' B.

			Ft.	In.
Coal	1'	8"		
Shale	0	2		
Coal	0	6	2	4

Coal Blossom—No. 257A on Map II.

On Cold Knob road, 4.15 miles south of Richwood and 0.7 mile southeast of Manning Knob; **Little Raleigh Coal**; elevation, 3520' B.

	Ft.	In.
Coal blossom, thickness not determined.....

Quantity of Little Raleigh Coal Available.

The estimates in the following table have been computed from planimetric measurement of outcrop outlined on work sheets for the areas indicated on Figure 19, page 531, and show the probable amount of Little Raleigh Coal in Greenbrier County:

Probable Amount of Little Raleigh Coal.

District.	Thickness of Coal Assumed. Feet.	Square Miles.	Acres.	Cubic Feet of Coal.	Short Tons of Coal (2000 Lbs.)
Meadow Bluff	2	26.5	16,960	1,477,555,200	59,102,208
Willamsburg	3	7.0	4,480	585,446,400	23,417,856
Totals		33.5	21,440	2,063,001,600	82,520,064

BECKLEY COAL.

The Beekley Coal previously discussed in Chapter VI, pages 236-7, ranks fourth in available tonnage within the county. In general this coal is from 2 to 5 feet thick. In appearance and chemical properties the Beekley Coal is quite similar to the Sewell seam and in at least one instance this fact has led to confusion in correlating from one area to another.

The first commercial mine (Lost Flat—No. 307 on Map II) in the county was opened in this seam in Falling Springs District in 1907. This mine was abandoned in 1910 when the same company opened the Spruce Knob Mine (No. 225 on Map II) on North Fork of Cherry River. At present (1936) there are no actively operating mines in this seam.

The probable minable area of Beekley Coal is shown on Figure 19, page 531, but the coal is not outcropped on Map II, as it is only 50 to 80 feet above the Fire Creek and its position may readily be interpolated from the position of the latter.

Beckley Coal, Meadow Bluff District.

In this district the Beekley Coal reaches its best development on Little Clear Creek Mountain where there is a fairly large area of coal with a thickness of 3 to 5 feet. Over much

of the rest of the area indicated on Figure 19, page 531, for this district, this coal is generally between 2 and 3½ feet thick and while it may not be profitable to mine this thinner coal at the present time, it should be considered a minable reserve.

Because of insufficient information, the Beekley Coal is not shown on Figure 19 as minable in Hickory Ridge, Cross Mountain, and Buffalo Mountain. Future prospecting, however, may prove it to be present in these areas in commercial thickness.

The Beekley Coal is noted in the records of coal test borings Nos. 5, 6, 11, 12, and 14, the details of which are published on preceding pages. In addition its stratigraphic position is shown in the partial records of coal test borings Nos. 5A, 5B, 5C, 5D, 5F, 5G, 5H, and 5K. The following openings and prospects were noted:

Tuck Brothers Mine—No. 258 on Map II.

On Fayette-Greenbrier County line, 0.8 mile north of Rainelle; Beekley Coal; elevation, 2815' B.

			Ft.	In.
1. Slate, draw			1	6
2. Coal, bony	0'	2"		
3. Bone	0	1		
4. Coal, columnar	0	6		
5. Fusain (mineral charcoal)..	0	0½		
6. Coal	0	3		
7. Fusain (mineral charcoal)..	0	0½		
8. Coal, columnar	1	7½		
9. Coal, bony	0	4	3	0

A sample (No. 150PH) was taken from Nos. 4, 5, 6, 7, and 8 of the above section, the analysis of which is published under No. 258 in the Table of Coal Analyses at the end of this Chapter.

Meadow River Lumber Company? Prospect— No. 259 on Map II.

On north end of Little Sewell Mountain, 0.95 mile northeast of East Rainelle; Beekley Coal; elevation, 3080' B.

	Ft.	In.
Shale, black, concretions.....	8	0
Coal (slate floor)	2	8

**Gauley Coal Land Company? Mine (Abandoned)—
No. 260 on Map II.**

Truck mine, on the west side of Big Clear Creek, at the stream and road crossing, 0.95 mile southwest of Duo and 3.2 miles northeast of Anjean; has been mined for local use; Beckley Coal; elevation, 3160' B.

	Ft.	In.
Sandstone, grayish-white		
Coal, good	2	6
Shale	10	2
Sandstone, shaly		

**Raine Lumber and Coal Company Prospect—
No. 261 on Map II.**

On the east side of Big Clear Creek, 1.2 miles southwest of Duo and 3 miles northeast of Anjean; Beckley Coal; elevation, 3190' B.

		Ft.	In.
Coal, bony (sandstone roof).....	0' 3"		
Coal, weathered	1 11	2	2
Concealed			

L. E. McClung Prospect—No. 262 on Map II.

On the west side of Shellcamp Ridge, 2 miles southwest of Duo and 2.25 miles northeast of Anjean; fallen shut, reported by L. E. McClung; Beckley Coal; elevation, 3285' B.

		Ft.	In.
Shale, black			
Coal, shale partings.....	3' 4"		
Shale	2 0		
Coal	1 10	7	2

L. E. McClung Prospect—No. 263 on Map II.

On the west side of Shellcamp Ridge, 2.15 miles southwest of Duo and 2.1 miles northeast of Anjean; Beckley Coal; elevation, 3310' B.

		Ft.	In.
Shale, black			
Coal	0' 5"		
Shale	0 1		
Coal	0 0½		
Shale	0 0½		
Coal	0 3½		
Fusain (mineral charcoal).....	0 0½		
Coal, laminated	2 6	3	5

**Gauley Coal Land Company Prospect No. 4—
No. 264 on Map II.**

On the east side of Smokehouse Branch, 1.45 miles northeast of its mouth and 1.75 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3544' L.

	Ft.	In.
Coal (slate roof and floor)	1	10

**Gauley Coal Land Company Prospect No. 2—
No. 265 on Map II.**

On the east side of Smokehouse Branch, 1.25 miles northeast of its mouth and 1.7 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3513' L.

	Ft.	In.
Coal (slate floor)	2	2

**Gauley Coal Land Company Prospect No. 9—
No. 266 on Map II.**

On the east side of Smokehouse Branch, 1 mile northeast of its mouth and 1.8 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3514' L.

		Ft.	In.
Coal (slate roof).....	0' 6"		
Fire clay	0 2		
Coal (slate floor).....	2 7	3	3

**Gauley Coal Land Company Prospect No. 14—
No. 267 on Map II.**

On the east side of Smokehouse Ridge, 0.35 mile north of the mouth of Job Knob Branch and 2.6 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3640' L.

	Ft.	In.
Coal (slate floor).....	2	0

**Gauley Coal Land Company Prospect No. 15—
No. 268 on Map II.**

On the east side of Smokehouse Ridge, 0.6 mile north of mouth of Job Knob Branch and 2.45 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3646' L.

	Ft.	In.
Coal (slate roof and floor).....	2	8

**Gauley Coal Land Company Prospect No. 18—
No. 269 on Map II.**

On the east side of Smokehouse Ridge, 2.35 miles southeast of Duo and 5.15 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3615' L.

	Ft.	In.
Coal (slate roof and floor).....	2	6

**Gauley Coal Land Company Prospect No. 22—
No. 270 on Map II.**

On the east side of Oldhouse Branch, 0.55 mile north of its mouth and 2.6 miles east of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3591' L.

	Ft.	In.
Bone and coal (slate roof).....	0'	4"
Coal (slate floor).....	2	1½
	2	5½

**Gauley Coal Land Company Prospect No. A237—
No. 271 on Map II.**

On the north side of Rockcamp Ridge, 1.25 miles south of Job Knob; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3734' L.

	Ft.	In.
Coal	2	11

**Gauley Coal Land Company Prospect No. A236—
No. 272 on Map II.**

On the north side of Rockcamp Ridge, 1.4 miles south of Job Knob; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3793' L.

	Ft.	In.
Coal	2	5

**Gauley Coal Land Company Prospect No. A235—
No. 273 on Map II.**

On the south side of Rockcamp Ridge, 1.6 miles southeast of Job Knob; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3616' L.

	Ft.	In.
Coal and slate	1	9

**Gauley Coal Land Company Prospect No. A234—
No. 274 on Map II.**

On the south side of Rockcamp Ridge, 1.5 miles southeast of Job Knob; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3863' L.

	Ft.	In.
Coal and slate	2	0½

**Gauley Coal Land Company Prospect No. A130—
No. 275 on Map II.**

On Old Field Mountain just south of Grassy Knob, 5.45 miles east of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 4213' L.

	Ft.	In.
Coal	2	2

**Gauley Coal Land Company Prospect No. A129—
No. 276 on Map II.**

On Old Field Mountain, 7 miles east of Anjean and 5.4 miles south-east of Duo; Beckley Coal; elevation, 4162' L.

		Ft.	In.
Coal	2' 11"		
Coal and slate (fire clay floor)....	1 0	3	11

**Gauley Coal Land Company Prospect No. A127—
No. 277 on Map II.**

On north side of Little Clear Creek Mountain, 6.35 miles north-east of Anjean and 4.5 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3879' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	3	11

**Gauley Coal Land Company Prospect No. A126—
No. 278 on Map II.**

On north side of Little Clear Creek Mountain just above core test No. 13, 6.15 miles east of Anjean and 4.55 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3817' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	3	5

A sample was taken from the above section by the Gauley Coal Land Company and analyzed by the Commercial Testing and Engineering Company, of Charleston, W. Va. Its analysis is published under No. 278 in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. A125—
No. 279 on Map II.**

On the north side of Little Clear Creek Mountain, 5.35 miles northeast of Anjean and 3.6 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3706' L.

			Ft.	In.
Bone and coal (slate roof).....	0'	2½"		
Coal	0	10		
Slate and coal	0	1		
Coal (fire clay floor).....	2	3½	3	5

**Gauley Coal Land Company Prospect No. A122—
No. 280 on Map II.**

On north side of Little Clear Creek Mountain, 4.3 miles northeast of Anjean and 3.2 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3714' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	2	7

**Gauley Coal Land Company Prospect No. A121—
No. 281 on Map II.**

On the north side of Little Clear Creek Mountain, 4.2 miles northeast of Anjean and 3.35 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3732' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	3	1

**Gauley Coal Land Company Prospect No. A120—
No. 282 on Map II.**

On the north side of Little Clear Creek Mountain, 4 miles northeast of Anjean and 3.45 miles southeast of Duo; Gauley Coal Land Company authority for this section; Beckley Coal; elevation, 3740' L.

	Ft.	In.
Coal (slate roof; fire clay floor).....	2	7

The above opening may be in the "Split" seam coming a few feet under the Beckley proper.

In the vicinity of Joe Knob so many prospect openings have been made in or near the Beckley Coal horizon that it was not possible to show all of them on Map II. A seam between the Beckley and Fire Creek has been opened in several places on the knob. This seam is classified as the "Split" seam on the Gauley Coal Land Company's maps but it is not cer-

**Gauley Coal Land Company Prospect No. 396—
No. 171 on Map II.**

On the west side of Little Trap Ridge, 2 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3060' B.

	Ft.	In.
Coal	3	11

**Gauley Coal Land Company Prospect No. 397—
No. 172 on Map II.**

On the north end of Little Trap Ridge, 2.9 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2831' B.

	Ft.	In.
Coal	2	2

Gauley Coal Land Company Prospect—No. 173 on Map II.

On the north side of Hominy Creek, 3.6 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2821' B.

	Ft.	In.
Coal	5	5

**Gauley Coal Land Company Prospect No. 398—
No. 174 on Map II.**

On the east side of Little Trap Ridge, 2.7 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2932' B.

	Ft.	In.
Coal	5	7½

**Gauley Coal Land Company Prospect No. 399—
No. 175 on Map II.**

On the east side of Little Trap Ridge, 2.5 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3060' B.

	Ft.	In.
Coal	3	11

**Gauley Coal Land Company Prospect No. 401—
No. 176 on Map II.**

On the northwest side of Big Clear Creek Mountain, 2.8 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3208' B.

	Ft.	In.
Coal	1'	6"
Slate	0	1
Coal	2	6
	4	1

**Gauley Coal Land Company Prospect No. 402—
No. 177 on Map II.**

On the northwest side of Big Clear Creek Mountain, 3 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3234' B.

			Ft.	In.
Coal	1'	4"		
Slato	6	0		
Coal	2	10	10	2

**Gauley Coal Land Company Prospect No. 404—
No. 178 on Map II.**

On the northwest side of Big Clear Creek Mountain, 3.2 miles northwest of Duo and 3 miles northeast of Marfrance; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3234' L.

	Ft.	In.
Coal	2	10

**Gauley Coal Land Company Prospect No. 405—
No. 179 on Map II.**

On the headwaters of Hominy Creek, 3.2 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3191' L.

	Ft.	In.
Coal	2	10

**Gauley Coal Land Company Prospect No. 406—
No. 180 on Map II.**

On the headwaters of Hominy Creek, 3.1 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3212' L.

	Ft.	In.
Coal	3	6

**Gauley Coal Land Company, Prospect No. 407—
No. 181 on Map II.**

On the headwaters of Hominy Creek, 2.6 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3313' L.

	Ft.	In.
Coal	5	6

**Gauley Coal Land Company Prospect No. 407A—
No. 182 on Map II.**

On the headwaters of Hominy Creek, 2.5 miles northwest of Duo;
Gauley Coal Land Company authority for this section; Sewell Coal;
elevation, 3424' L.

	Ft.	In.
Coal	2	6

**Gauley Coal Land Company Prospect No. 409—
No. 183 on Map II.**

On the south side of Blue Ridge, 2.7 miles northwest of Duo;
Gauley Coal Land Company authority for this section; Sewell Coal;
elevation, 3377' L.

	Ft.	In.
Coal	3	2

**Gauley Coal Land Company Prospect No. 410—
No. 184 on Map II.**

On the west side of Blue Ridge, 2.3 miles northwest of Duo; Gauley
Coal Land Company authority for this section; Sewell Coal, elevation,
3210' L.

	Ft.	In.
Coal	3	1

**Gauley Coal Land Company Prospect No. 412—
No. 185 on Map II.**

On the north side of Blue Ridge, 2.3 miles southwest of Beech
Knob; Gauley Coal Land Company authority for this section; Sewell
Coal; elevation, 3259' L.

	Ft.	In.
Coal	2	5

**Gauley Coal Land Company Prospect No. 413—
No. 186 on Map II.**

On the north side of Blue Ridge, 1.65 miles southwest of Beech
Knob; Gauley Coal Land Company authority for this section; Sewell
Coal; elevation, 3362' L.

	Ft.	In.
Coal	2	1

Perry Amick Mine—No. 187 on Map II.

Farm mine. 1.75 miles southwest of Lile and 1 mile southeast of
White Buck School; authority, David B. Reger (No. 1217, Nicholas
Report, page 702); Sewell Coal; elevation, 3120' B.

	Ft.	In.
Coal (slate roof and floor)	3	2

**Gauley Coal Land Company Prospect No. 416—
No. 188 on Map II.**

1.4 miles southwest of Lile and 1.15 miles southeast of White Buck School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3232' L.

			Ft.	In.
Coal	1'	2"		
Slate	0	10		
Coal	3	4	5	4

**Gauley Coal Land Company Prospect No. 418—
No. 189 on Map II.**

1.3 miles southwest of Lile and 1 mile southeast of White Buck School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3215' L.

			Ft.	In.
Coal	1'	4"		
Slate	1	2		
Coal	2	11	5	5

**Gauley Coal Land Company Prospect No. 419—
No. 190 on Map II.**

1.1 miles southwest of Lile and 1.2 miles southeast of White Buck School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3245' L.

			Ft.	In.
Coal	2'	4½"		
Slate	1	5		
Coal	2	5	6	2½

**Gauley Coal Land Company Prospect No. 420—
No. 191 on Map II.**

0.85 mile west of Lile and 1.3 miles southeast of White Buck School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3215' L.

			Ft.	In.
Laminated	0'	10"		
Coal	2	6	3	4

**Gauley Coal Land Company Prospect No. 422—
No. 192 on Map II.**

On the headwaters of Brushy Meadow Creek, 1.25 miles northwest of Lile and 0.85 mile east of White Buck School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3066' L.

			Ft.	In.
Coal			3	11

**Gauley Coal Land Company Prospect No. 423—
No. 193 on Map II.**

On the headwaters of Brushy Meadow Creek, 1.2 miles northwest of Lille and 0.95 mile east of White Buck School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3096' L.

			Ft.	In.
Coal	1'	2"		
Slate	0	1		
Coal	2	9	4	0

**Gauley Coal Land Company Prospect No. 424—
No. 194 on Map II.**

On the headwaters of Brushy Meadow Creek, 0.9 mile northwest of Lille; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3137' L.

			Ft.	In.
Coal	1'	3"		
Slate	0	8		
Coal	2	6½	4	5½

**Gauley Coal Land Company Prospect No. 425—
No. 195 on Map II.**

On the headwaters of Brushy Meadow Creek, 0.6 mile northwest of Lille; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3223' L.

	Ft.	In.
Coal	3	7

Sewell Coal, Williamsburg District.

Of the estimated 22,693,016 tons of Sewell Coal in Williamsburg District, about 95 per cent. is on Beech Ridge. The other 5 per cent. is accounted for by small isolated areas of coal on or near Manning Knob and small areas on Kerless and Sugar Knobs. The following openings and prospects were noted:

Marshall Amick Mine (Abandoned)—No. 196 on Map II.

Farm mine, on the head of Paek Fork of Laurel Creek, 3.9 miles south 69° east of Leivasy and 1 mile northeast of Lille; authority, David B. Reger (No. 1184, Nicholas Report, page 694); Sewell Coal; elevation, 3070' B.

	Ft.	In.
Shale, sandy	10	0
Coal, soft (slate floor).....	2	5

Levi Lilly Mine (Abandoned)—No. 197 on Map II.

Farm mine, on the north bank of McMillon Creek, 1.7 miles north-east of Lile and 4.6 miles south 80° east of Leivasy; authority, David B. Reger (No. 1183, Nicholas Report, page 693); Sewell Coal; elevation, 3155' B.

	Ft.	In.
Slate, dark
Coal, soft, columnar	2	6
Shale, gray, with plant roots

Gauley Coal Land Company Prospect No. 31— No. 198 on Map II.

On southeast side of McMillon Creek, 0.2 mile east of Lile; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3317' L.

	Ft.	In.
Coal	3	1

Gauley Coal Land Company Prospect No. 32— No. 199 on Map II.

On the headwaters of Beech Run, 0.9 mile east of Lile and 0.85 mile north of Beech Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3426' L.

	Ft.	In.
Coal	2	5

Gauley Coal Land Company Prospect No. 33— No. 200 on Map II.

1.25 miles northeast of Lile and 1.2 miles northeast of Beech Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3417' L.

	Ft.	In.
Coal	2	10

Gauley Coal Land Company Prospect No. 34— No. 201 on Map II.

1.45 miles northeast of Lile and 1.2 miles northeast of Beech Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3496' L.

	Ft.	In.
Coal	3	0

Gauley Coal Land Company Prospect No. 35— No. 202 on Map II.

On the headwaters of Hogeamp Run, 0.55 mile east of Little Beech Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3621' L.

	Ft.	In.
Coal	3	8

**Gauley Coal Land Company Prospect No. 423—
No. 193 on Map II.**

On the headwaters of Brushy Meadow Creek, 1.2 miles northwest of Lile and 0.95 mile east of White Buck School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3096' L.

			Ft.	In.
Coal	1'	2"		
Slate	0	1		
Coal	2	9	4	0

**Gauley Coal Land Company Prospect No. 424—
No. 194 on Map II.**

On the headwaters of Brushy Meadow Creek, 0.9 mile northwest of Lile; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3137' L.

			Ft.	In.
Coal	1'	3"		
Slate	0	8		
Coal	2	6½	4	5½

**Gauley Coal Land Company Prospect No. 425—
No. 195 on Map II.**

On the headwaters of Brushy Meadow Creek, 0.6 mile northwest of Lile; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3223' L.

	Ft.	In.
Coal	3	7

Sewell Coal, Williamsburg District.

Of the estimated 22,693,016 tons of Sewell Coal in Williamsburg District, about 95 per cent. is on Beech Ridge. The other 5 per cent. is accounted for by small isolated areas of coal on or near Manning Knob and small areas on Kerless and Sugar Knobs. The following openings and prospects were noted:

Marshall Amick Mine (Abandoned)—No. 196 on Map II.

Farm mine, on the head of Pack Fork of Laurel Creek, 3.9 miles south 69° east of Leivasy and 1 mile northeast of Lile; authority, David B. Reger (No. 1184, Nicholas Report, page 694); Sewell Coal; elevation, 3070' B.

	Ft.	In.
Shale, sandy	10	0
Coal, soft (slate floor).....	2	5

Levi Lilly Mine (Abandoned)—No. 197 on Map II.

Farm mine, on the north bank of McMillon Creek, 1.7 miles north-east of Lile and 4.6 miles south S0° east of Leivasy; authority, David B. Reger (No. 1183, Nicholas Report, page 693); Sewell Coal; elevation, 3155' B.

	Ft.	In.
Slate, dark
Coal, soft, columnar	2	6
Shale, gray, with plant roots

Gauley Coal Land Company Prospect No. 31— No. 198 on Map II.

On southeast side of McMillon Creek, 0.2 mile east of Lile; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3317' L.

	Ft.	In.
Coal	3	1

Gauley Coal Land Company Prospect No. 32— No. 199 on Map II.

On the headwaters of Beech Run, 0.9 mile east of Lile and 0.85 mile north of Beech Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3426' L.

	Ft.	In.
Coal	2	5

Gauley Coal Land Company Prospect No. 33— No. 200 on Map II.

1.25 miles northeast of Lile and 1.2 miles northeast of Beech Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3417' L.

	Ft.	In.
Coal	2	10

Gauley Coal Land Company Prospect No. 34— No. 201 on Map II.

1.45 miles northeast of Lile and 1.2 miles northeast of Beech Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3496' L.

	Ft.	In.
Coal	3	0

Gauley Coal Land Company Prospect No. 35— No. 202 on Map II.

On the headwaters of Hogeamp Run, 0.55 mile east of Little Beech Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3621' L.

	Ft.	In.
Coal	3	8

**Gauley Coal Land Company Prospect No. 36—
No. 203 on Map II.**

On the north side of Beech Ridge, 1.9 miles east of Beech Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3525' L.

	Ft.	In.
Coal	4	1

**Gauley Coal Land Company Prospect No. 36A—
No. 204 on Map II.**

On north side of Beech Ridge, 2 miles east of Beech Knob; Gauley Coal Land Coal Company authority for this section; Sewell Coal; elevation, 3521' L.

	Ft.	In.
Coal	4	10

**Gauley Coal Land Company Prospect No. 37—
No. 205 on Map II.**

On the north side of Beech Ridge, 2.1 miles east of Beech Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3513' L.

	Ft.	In.
Coal	0'	2½"
Clay	0	0½
Coal	5	0
	5	3

**Gauley Coal Land Company Prospect No. 38—
No. 206 on Map II.**

On the north side of Beech Ridge, 2.45 miles east of Beech Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3527' L.

	Ft.	In.
Coal	2'	5"
Slate	0	1
Coal	1	3
Bone	0	1
	3	10

**Gauley Coal Land Company Prospect No. 39—
No. 207 on Map II.**

On the north side of Beech Ridge, 1.3 miles northeast of Clearco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3545' L.

	Ft.	In.
Bone	0'	6"
Coal	3	8
Laminated	0	2½
Coal	0	8
Bone	0	1
Coal	0	8
Bone	0	2
	5	11½

**Gauley Coal Land Company Prospect No. 39A—
No. 208 on Map II.**

On the northeast side of Beech Ridge, 1.25 miles east of Clearco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3602' L.

	Ft.	In.
Coal	4	5

**Gauley Coal Land Company Prospect No. 39B—
No. 209 on Map II.**

On the northeast side of Beech Ridge, 1.3 miles east of Clearco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3622' L.

	Ft.	In.
Coal	4	2

**Gauley Coal Land Company Prospect No. 40—
No. 210 on Map II.**

On the northeast side of Beech Ridge, 1.4 miles east of Clearco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3656' L.

	Ft.	In.
Coal	4	2

**Gauley Coal Land Company Prospect No. 40A—
No. 211 on Map II.**

On the northeast side of Beech Ridge, 1.5 miles east of Clearco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3656' L.

	Ft.	In.
Coal	4	2½

Gauley Coal Land Company Prospect—No. 212 on Map II.

0.7 mile northeast of Job Knob and 1.7 miles southeast of Clearco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3733' L.

	Ft.	In.
Coal	2	9

Gauley Coal Land Company Prospect—No. 213 on Map II.

0.6 mile northeast of Job Knob and 1.7 miles southeast of Clearco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3753' L.

	Ft.	In.
Coal	3	0

Raine Coal and Land Company Prospect—No. 214 on Map II.

0.9 mile northeast of Job Knob and 2.2 miles southeast of Clearco;
Sewell Coal; elevation, 3809' L.

			Ft.	In.
1. Coal	3'	0"		
2. Coal (reported by J. W. Raine)	1	5	4	5

A sample (No. 155PH) was taken from No. 1 of the above section and its composition is published under No. 214 in the Table of Coal Analyses at the end of this Chapter.

Raine Lumber and Coal Company Mine (Abandoned)— No. 215 on Map II.

On the north side of Beech Ridge, 2.7 miles southeast of Clearco and 4.5 miles east of Duo; section measured 50 feet in from mine mouth; Sewell Coal; elevation, 3946' L.

			Ft.	In.
Shale roof				
Coal, blocky, some bone at top..	1'	0"		
Coal, columnar	1	6		
Coal, hard, laminated.....	1	0		
Coal, hard, bony at base.....	0	11	4	5

A sample (No. 156PH) was taken from the above section and its analysis is published under No. 215 in the Table of Coal Analyses at the end of this Chapter.

Raine Lumber and Coal Company Prospect— No. 216 on Map II.

On the north side of Beech Ridge, 3.1 miles southeast of Clearco, and 4.75 miles east of Duo; Sewell Coal; elevation, 4013' L.

	Ft.	In.
Coal, (reported by J. W. Raine), 3' to.....	4	0

Coal Blossom—No. 217 on Map II.

On the Jetsville-Manning Knob road, 4 miles northeast of Clearco and 1.7 miles west of Manning Knob; Sewell Coal; elevation, 3665' B.

	Ft.	In.
Coal, exposed	1	6

**Gauley Coal Land Company (?) Prospect—
No. 218 on Map II.**

5.3 miles northeast of Lile and 1.6 miles west of Manning Knob;
Sewell Coal; elevation, 3620' B.

	Ft.	In.
Coal, reported	3	4

**Gauley Coal Land Company (?) Prospect—
No. 219 on Map II.**

On the Jetsville-Manning Knob road, 5.3 miles northeast of Lile
and 1.5 miles northwest of Manning Knob; Sewell Coal; elevation,
3660' B.

	Ft.	In.
Coal, reported	3	4

**Gauley Coal Land Company (?) Mine (Abandoned)—
No. 220 on Map II.**

On the Jetsville-Manning Knob road, 5.7 miles northeast of Lile
and 1.4 miles northwest of Manning Knob; Sewell Coal; elevation,
3700' B.

	Ft.	In.
Coal, reported	2	2

Coal Prospect—No. 221 on Map II.

On the east side of Manning Knob, on the west side of the Cold
Knob road; Sewell Coal; elevation, 3870' B.

	Ft.	In.
Coal, reported by Sam Howard to be 1' 6" to	2	0

Coal Blossom—No. 221A on Map II.

On the east side of Manning Knob, on the west side of Cold Knob
road; Sewell Coal; elevation, 3870' B.

	Ft.	In.
Coal, thickness not determined.....

John A. Bailes Coal Stripping—No. 222 on Map II.

On south side of Nixon Branch of Laurel Creek, 2.8 miles S. 22°
E. of Saxman; authority David B. Reger (No. 1182 Nicholas Report,
page 693); Sewell Coal; elevation, 2995' B.

	Ft.	In.
1. Slate, black.....		
2. Coal, soft	4	5
3. Slate, pavement		

A sample (No. 376R) was collected from No. 2 of section,
the composition of which is published under No. 222 in the
Survey Table of Coal Analyses at the end of this Chapter.

Coal Blossom—No. 223 on Map II.

On private road at Greenbrier-Nicholas County line, 1.9 miles south of Richwood; Sewell Coal; elevation, 2990' B.

	Ft.	In.
Coal, thickness not determined.....

Sewell Coal, Falling Springs District.

In Falling Springs District, only a small fraction of the 18.6 square miles believed to be underlain by Sewell Coal has been prospected. Six openings and prospects in the Sewell Coal were found and they are all in the immediate vicinity of the two abandoned mines of the Elk Lick Coal Company. The thickness of the coal in and around these mines varies from two feet to four feet and ten inches; however, due to the uncertainties involved, the low average thickness of two feet was used in computing the estimate of 41,483,059 tons of Sewell Coal present in Falling Springs District. It is believed that a large part of the 18.6 square miles is underlain by Sewell Coal with a thickness in excess of 3 feet and that the estimate is conservative.

Elk Lick Coal Company "Turkey Run" Mine (Abandoned)— No. 224 on Map II.

On the east side of Turkey Run of North Fork of Cherry River, 4.6 miles southeast of Richwood; Sewell Coal; elevation, 3370' L.

	Ft.	In.
Coal	4	8

The coal production statistics given at the beginning of this Chapter include the production from this mine with the production of the following mine under the "Spruce Knob" mine.

**Elk Lick Coal Company "Spruce Knob" Mine (Abandoned)—
No. 225 on Map II.**

On the south side of North Fork of Cherry River, 5 miles southeast of Richwood; Sewell Coal; elevation, 3379' L.

		Ft.	In.
1. Shale, black, weathers red, thin and platy, fossil col- lection 149		5	0
2. Coal, laminated	1' 6½"		
3. Coal, columnar	1 1½		
4. Coal, laminated	0 7		
5. Coal, columnar	0 8		
6. Coal, laminated	0 7	4	6
7. Sandstone, shaly			

A sample (No. 163PH) was taken from Nos. 2, 3, 4, 5, and 6 of the above section and its analysis is published under No. 225 in the Table of Coal Analyses at the end of this Chapter.

The following section and comment by Reger is reprinted from page 688 of the Nicholas County Report:

	Ft.	In.
1. Slate, dark		
2. Coal, soft, columnar	4	6
3. Slate, pavement		

"Principal office, Richwood, W. Va.; daily output, 100 tons; 15 miners and 9 laborers employed; mule haulage; greatest rise, southeast; coal used for railroad fuel by Cherry River Boom & Lumber Company; H. C. Livesay, Timekeeper, authority for mine data; sample (No. 211R) collected in First Left from No. 2 of section, for composition of which see (No. 225) in the Survey Table of Coal Analyses at the end of this Chapter."

As noted in the headings, mines Nos. 224 and 225 are now abandoned. Any correspondence concerning them should be addressed to the Cherry River Boom & Lumber Company, Richwood, W. Va.

Elk Lick Coal Company Prospect—No. 226 on Map II.

On the north side of Briery Run, 4.3 miles southeast of Richwood; Sewell Coal; elevation, 3400' B.

	Ft.	In.
Coal thickness undetermined.....

Elk Lick Coal Company Prospect—No. 227 on Map II.

On the head of Beech Lick Run, 4.7 miles southeast of Richwood; authority, Elk Lick Coal Company's mine map; Sewell Coal; elevation, 3456' L.

	Ft.	In.
Coal	4	0

Elk Lick Coal Company Prospect—No. 228 on Map II.

On the north side of Rocky Run, 5.5 miles southeast of Richwood; authority, Elk Lick Coal Company's mine map; Sewell Coal; elevation, 3570' L.

	Ft.	In.
Coal	4	0

Elk Lick Coal Company Prospect—No. 229 on Map II.

On the waters of Rocky Run, on the south side of Bearwallow Knob, 6.1 miles southeast of Richwood; authority, Elk Lick Coal Company's mine map; Sewell Coal; elevation, 3670' L.

	Ft.	In.
Coal	2	0

The following section is of a mine in Pocahontas County near the Greenbrier line. The section with comments by Paul H. Price is reprinted from page 297 of the Pocahontas County Report:

Preston Clark Heirs Prospect—No. 230 on Map II.

Pocahontas County, Little Levels District; on west side of Briery Knob, one-half mile northwest of triangulation station and Fire-Tower; Sewell Coal; elevation, 4225' B.

		Ft.	In.
Shale, Hartridge; plants and pelecypods.....			
Coal, good, clean	2' 2"		
Shale, argillaceous	1 10		
Coal, clean	1 3		
Coal, bony.....	1 0	6	3
Concealed			

"It is doubtful if the complete thickness of the coal here is revealed by this section. Mr. Lee Clark, who had been in the mine, reported as much as eleven feet at certain points."

A sample (No. 62PH) of this coal was taken, the composition of which appears under Mine No. 230 in the Table of Coal Analyses at the end of this Chapter.

Quantity of Sewell Coal Available.

The following table, calculated from planimetric measurement of the Sewell Coal outcrop on Map II, for the minable areas as indicated in Figure 17, shows the probable amount of Sewell Coal in Greenbrier County. The assumed average thicknesses of coal and the total tonnage are believed to be quite conservative:

Probable Amount of Sewell Coal.

District.	Thickness of Coal Assumed, Feet.	Square Miles.	Acres.	Cubic Feet of Coal.	Short Tons of Coal (2000 Lbs.)
Meadow Bluff	2	13.20	8,448	735,989,760	29,439,590
Meadow Bluff	3	6.65	4,256	556,174,080	22,246,963
Meadow Bluff	3½	12.00	7,680	1,170,892,800	46,835,712
Meadow Bluff	4	2.00	1,280	223,027,200	8,921,088
Meadow Bluff	4½	8.80	5,632	1,103,984,640	44,159,385
Williamsburg	2½	2.30	1,472	160,300,800	6,412,032
Williamsburg	3	2.60	1,664	217,451,520	8,698,060
Williamsburg	3½	0.80	512	78,059,520	3,122,380
Williamsburg	4	1.00	640	111,513,600	4,460,544
Falling Springs	2	18.60	11,904	1,037,076,480	41,438,059
Totals		67.95	43,488	5,394,470,400	215,778,813

According to the records of the State Department of Mines, the total coal mined at operations in the Sewell Coal of Greenbrier County to the end of the calendar year 1936, is 22,382,111 short tons. Assuming a recovery factor of 80 per cent., the 215,778,813 tons above would be reduced to 172,623,050 short tons of Sewell Coal in Greenbrier County, from which should be deducted the amount already mined, leaving a recoverable tonnage on above basis of 150,240,939.

**Gauley Coal Land Company Prospect No. 105—
No. 110 on Map II.**

On the west side of Sam Rldge, 1.2 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3491' L.

			Ft.	In.
Slate	0'	5"		
Coal	3	4		
Bone	2	4	6	1

**Gauley Coal Land Company Prospect No. 104—
No. 111 on Map II.**

On the west side of Sam Rldge, 1.3 miles west of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3486' L.

			Ft.	In.
Slate	0'	5"		
Coal	3	3	3	8

**Gauley Coal Land Company Prospect No. 103—
No. 112 on Map II.**

On the west side of Sam Rldge, 1.3 miles southwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3515' L.

		Ft.	In.
Coal		3	1

**Gauley Coal Land Company Prospect No. 102—
No. 113 on Map II.**

On the south end of Sam Rldge, 1 mile southwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3486' L.

		Ft.	In.
Coal		3	10

**Gauley Coal Land Company Prospect No. 101—
No. 114 on Map II.**

On the east side of Sam Rldge, 0.7 mile southwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3487' L.

		Ft.	In.
Coal		3	10

**Gauley Coal Land Company Prospect No. 100—
No. 115 on Map II.**

On the east side of Sam Ridge, 0.7 mile northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3425' L. (?) (3452').

	Ft.	In.
Coal	3	9

**Gauley Coal Land Company Prospect No. 99—
No. 116 on Map II.**

On the east side of Sam Ridge, 0.8 mile northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3448' L.

	Ft.	In.
Coal	3'	4"
Slate	0	6
	3	10

**Gauley Coal Land Company Prospect No. 96—
No. 117 on Map II.**

On the west bank of Elijah Branch, 1.5 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3490' L.

	Ft.	In.
Coal	3	2

**Gauley Coal Land Company Prospect No. 95—
No. 118 on Map II.**

On the west bank of Elijah Branch, 1.7 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3509' L.

	Ft.	In.
Bone	1'	8"
Coal	2	6
	4	2

**Gauley Coal Land Company Prospect No. 94—
No. 119 on Map II.**

On the west bank of Elijah Branch, 1.8 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3504' L.

	Ft.	In.
Bone	1'	8"
Coal	2	6
	4	2

**Gauley Coal Land Company Prospect No. 92—
No. 120 on Map II.**

On the east bank of Elijah Branch, 1.9 miles northwest of Duo;
Gauley Coal Land Company authority for this section; Sewell Coal;
elevation 3483' L.

	Ft.	In.
Coal	3	0

**Gauley Coal Land Company Prospect No. 91—
No. 121 on Map II.**

On the east bank of Elijah Branch, 1.7 miles northwest of Duo;
Gauley Coal Land Company authority for this section; Sewell Coal;
elevation, 3485' L.

	Ft.	In.
Coal	3	1

**Gauley Coal Land Company Prospect No. 90—
No. 122 on Map II.**

On the east bank of Elijah Branch, 1.5 miles northwest of Duo;
Gauley Coal Land Company authority for this section; Sewell Coal;
elevation, 3468' L.

	Ft.	In.
Coal	3	3

**Gauley Coal Land Company Prospect No. 89—
No. 123 on Map II.**

On the east bank of Elijah Branch, 1.3 miles northwest of Duo;
Gauley Coal Land Company authority for this section; Sewell Coal;
elevation, 3475' L.

	Ft.	In.
Coal	2	7

**Gauley Coal Land Company Prospect No. 88A—
No. 124 on Map II.**

On the west bank of Road Branch, 1.2 miles north of Duo; Gauley
Coal Land Company authority for this section; Sewell Coal; elevation,
3450' L.

			Ft.	In.
Coal	3'	4"		
Bone	1	2		
Coal	1	2		
Bone	0	3	5	11

**Gauley Coal Land Company Prospect No. 88—
No. 125 on Map II.**

On the west bank of Road Branch, 1.7 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3476' L.

			Ft.	In.
Coal	3'	2"		
Bone	0	9		
Coal and bone, interlaminated..	1	3	5	2

**Gauley Coal Land Company Prospect No. 87—
No. 126 on Map II.**

On the west bank of Road Branch, 2 miles north of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3507' L.

		Ft.	In.
Coal		2	6

**Gauley Coal Land Company Prospect No. 86—
No. 127 on Map II.**

On the east bank of Road Branch, 1.5 miles north of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3491' L.

			Ft.	In.
Coal and bone, interlaminated..	1'	0"		
Coal	3	7	4	7

**Gauley Coal Land Company Prospect No. 84—
No. 128 on Map II.**

On the east bank of Road Branch, 1.9 miles southwest of Cleareo and 1.25 miles north of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3470' L.

			Ft.	In.
Bone	1'	2½"		
Coal	0	6		
Bone	0	0½		
Coal	2	3½	4	0½

**Gauley Coal Land Company Prospect No. 83C—
No. 129 on Map II.**

On the north bank of North Fork of Big Clear Creek, 1.8 miles west of Clearco and 1.4 miles north of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3495' L.

			Ft.	In.
Bone	0'	7"		
Coal	2	7		
Slate	0	2½		
Coal	0	3½	3	8

**Gauley Coal Land Company Prospect No. 83B—
No. 130 on Map II.**

On the south side of Beech Ridge, 1.7 miles northwest of Clearco and 2.1 miles north of Duo; Gauley Coal Land Company authority for this section; Sewell Coal, elevation, 3542' L.

		Ft.	In.
Coal		2	7

**Gauley Coal Land Company Prospect No. 83A—
No. 131 on Map II.**

On the south side of Beech Ridge, 1.6 miles northwest of Clearco and 2.2 miles north of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3552' L.

		Ft.	In.
Coal		3	7

**Gauley Coal Land Company Prospect No. 83—
No. 132 on Map II.**

On the south side of Beech Ridge, 1.5 miles southwest of Clearco and 1.3 miles northeast of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3467' L.

			Ft.	In.
Coal	0'	5"		
Slate	0	0½		
Coal	3	1		
Coal and bone, Interlaminated..	0	3	3	9½

**Gauley Coal Land Company Prospect No. 82—
No. 133 on Map II.**

On the south side of Beech Ridge, 1 mile northwest of Clearco and 2.1 miles northeast of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation 3496' L.

			Ft.	In.
Coal and bone, interlaminated..	0'	8"		
Coal	2	6		
Slate	0	1		
Coal	1	0		
Slate	0	4		
Coal	0	2	4	9

**Gauley Coal Land Company Prospect No. 81—
No. 134 on Map II.**

On the south side of Beech Ridge, 0.85 mile northwest of Clearco; and 2.1 miles northeast of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3497' L.

			Ft.	In.
Coal	0'	8"		
Slate	0	7		
Coal	3	0		
Slate	0	7		
Coal	0	11		
Slate	0	4		
Coal	1	3		
Slate	0	3	7	7

**Gauley Coal Land Company Prospect No. 80—
No. 135 on Map II.**

On the south side of Beech Ridge, 0.8 mile west of Clearco and 2 miles northeast of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3497' L.

			Ft.	In.
Coal	4'	4"		
Slate	0	7		
Coal	0	4	5	3

**Gauley Coal Land Company Prospect No. 79—
No. 136 on Map II.**

On the south side of Beech Ridge, 0.8 mile southwest of Clearco and 1.9 miles northeast of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3513' L.

			Ft.	In.
Coal and bone, interlaminated..	1'	5"		
Coal	3	5	4	10

**Gauley Coal Land Company Prospect No. 78—
No. 137 on Map II.**

On the south side of Beech Ridge, 0.8 mile southwest from Clearco and 1.75 miles northeast of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3519' L.

			Ft.	In.
Coal and bone, interlaminated....	1'	5"		
Coal	2	11	4	4

**Gauley Coal Land Company Prospect No. 77—
No. 138 on Map II.**

On the south side of Beech Ridge, 0.5 mile west of Clearco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3518' L.

			Ft.	In.
Coal	3'	9"		
Slate	1	5		
Coal	0	11	6	1

**Gauley Coal Land Company Prospect No. 76—
No. 139 on Map II.**

On the south side of Beech Ridge, 0.4 mile west of Clearco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3503' L.

			Ft.	In.
Coal	3'	8"		
Slate	1	1		
Coal	1	4	6	1

**Gauley Coal Land Company Prospect No. 75—
No. 140 on Map II.**

On the south side of Beech Ridge, 0.4 mile northeast of Clearco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3526' L.

			Ft.	In.
Coal	0'	5"		
Slate	0	0½		
Coal	3	8	4	1½

**Gauley Coal Land Company Prospect No. 74—
No. 141 on Map II.**

On the south side of Beech Ridge, 0.2 mile northwest of Clearco; Sewell Coal; elevation, 3526' L.

			Ft.	In.
Shale, dark, sandy			15	0
Coal, bony	1'	0"		
Coal, clean	0	5		
Slate, 0' 1" to	0	2		
Coal, columnar	3	0		
Slate and shale	1	1		
Coal, clean (slate floor).....	1	2	6	10

**Clear Creek Coal Company "Brooke" Mine No. 2—
No. 142 on Map II.**

On the south side of Beech Ridge, 0.3 mile northeast of Clearco; section as shown on mine map; Sewell Coal; elevation, 3554' L.
Location of section; on 1st left, 500' from main entry.

			Ft.	In.
Bone	1'	3"		
Coal	3	8		
Slate	1	2		
Coal	1	3	7	4

Mine post-office address and shipping point, Clearco; mine superintendent, R. J. Holmes; on Nicholas, Fayette, and Greenbrier Railroad.

**Clear Creek Coal Company "Brooke" Mine No. 1—
No. 143 on Map II.**

On the south side of Beech Ridge, 0.3 mile northeast of Clearco; Sewell Coal; elevation, 3567' L.

Section measured at the face of main entry, 350' (east-northeast) from mine mouth.

			Ft.	In.
1. Coal, very hard (slate roof)	0'	3"		
2. Bone	0	1		
3. Coal, good, columnar	3	0		
4. Bone	0	4		
5. Coal, laminated (slate floor)	1	1	4	9

A sample (No. 97PH) was taken from Nos. 1, 2, 3, and 5 of the above section and its analysis is published under **No. 143** in the Table of Coal Analyses at the end of this Chapter.

Previous to the opening of the mine two samples were taken from a prospect opening at the same locality. The section is as follows:

			Ft.	In.
1. Coal, columnar (dark shale roof)	3'	8"		
2. Slate	1	1		
3. Coal, hard (slate floor).....	1	1	5	10

Sample No. 78A-PH was taken from No. 1 and sample No. 78B-PH was taken from No. 3 of the above section and their analyses are published under **No. 143** in the Table of Coal Analyses at the end of this Chapter. These samples show a lower volatile and higher ash content than the one taken inside the mine. As indicated by the above sections the parting thins rapidly to the east.

**Clear Creek Coal Company Prospect—
No. 144 on Map II.**

On the south side of Beech Ridge, 0.4 mile east of Clearco; Sewell Coal; elevation, 3565' B.

			Ft.	In.
Coal, columnar (shale roof)	3'	9"		
Shale and slate	1	1		
Coal, hard (slate floor).....	1	1	5	11

**Gauley Coal Land Company Prospect No. 72—
No. 145 on Map II.**

On the south side of Beech Ridge, 0.6 mile east of Clearco; Sewell Coal; elevation, 3567' L.

			Ft.	In.
Coal (shale roof, dark, sandy)	0'	3"		
Slate	0	2		
Coal, columnar	2	4		
Slate	0	4		
Coal (slate floor).....	1	10	4	11

**Gauley Coal Land Company Prospect No. 71—
No. 146 on Map II.**

1.1 miles northwest of Job Kuob and 0.6 mile southeast of Clearco; Sewell Coal; elevation, 3578' L.

			Ft.	In.
Shale, dark			5	0
Coal	0'	2"		
Slate	0	3½		
Coal, columnar	2	10		
Slate	1	2		
Coal (slate floor)	1	1	5	6½

**Gauley Coal Land Company Prospect No. 70—
No. 147 on Map II.**

1.2 miles northwest of Job Knob and 0.4 mile southeast of Clearco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3603' L.

			Ft.	In.
Coal	2'	0"		
Bone	0	1		
Coal	1	6		
Coal and bone, interlamated....	0	1	3	8

**Gauley Coal Land Company Prospect No. 69—
No. 148 on Map II.**

0.7 mile northwest of Job Knob and 1 mile southeast of Clearco;
Gauley Coal Land Company authority for this section; Sewell Coal;
elevation, 3640' L.

	Ft.	In.
Coal	3	5½

**Raine Lumber and Coal Company Prospect—
No. 149 on Map II.**

On the east side of Big Clear Creek, 0.5 mile north of Duo; Sewell
Coal; elevation, 3431' L.

	Ft.	In.
Shale, dark, weathers brown, conerations	8	0
Shale, black, slaty, small pelecypods	0	7
Coal some bone.....	0'	8"
Coal, columnar	2	2
Coal, hard, blocky (shale floor)	0	7
	3	5

Coal Prospect—No. 150 on Map II.

On the property of the Raine Lumber and Coal Company, behind
the "Old House" at Duo; Sewell Coal; elevation, 3422' L.

	Ft.	In.
Shale, dark, Hartridge		
Coal, clean (slate floor).....	3	4

A sample (No. 79PH) was taken from the above section
and its composition is published under No. 150 in the Table
of Coal Analyses at the end of this Chapter.

**Raine Lumber and Coal Company "Duo" Mine—
No. 151 on Map II.**

On the west side of Shelleamp Ridge, 0.5 mile southeast of Duo;
Sewell Coal; elevation, 3485' L.

	Ft.	In.
Coal, hard (black slate roof)....	0'	5"
Coal, blocky, laminated	0	7
Coal, columnar	0	7
Coal, soft, columnar	1	6
Coal, hard (slate floor).....	0	8
	3	9

The above section was measured at a prospect opening
before the mine was opened. The prospect was at the same
location as that of the mine and was driven in from the out-
crop 50 feet.

A sample (No. 153PH) was taken from the above section and its analysis is given under No. 151 in the Table of Coal Analyses at the end of this Chapter.

Post-office address and shipping point, Duo; superintendent, J. W. Rulne; on Nicholas, Fayette, and Greenbrier Railroad.

Raine Lumber and Coal Company Coal Stripping— No. 152 on Map II.

On the west side of Shellcamp Ridge, 1.3 miles south of Duo; Sewell Coal; elevation, 3567' L.

			Ft.	In.
Coal, bony (black shale roof).....	0'	9½"		
Coal, blocky	0	7		
Coal, columnar	1	6		
Coal, hard, blocky	0	4		
Bone	0	0½		
Coal (shale floor)	0	2	3	5

A sample (No. 152PH) was taken from the above section and its composition is published under No. 152 in the Table of Coal Analyses at the end of this Chapter.

Raine Lumber and Coal Company Coal Stripping— No. 153 on Map II.

On Shellcamp Ridge, 1.5 miles south of Duo; Sewell Coal; elevation, 3600' B.

	Ft.	In.
Coal (small boulder in top), 3' to.....	4	0

Raine Lumber and Coal Company Prospect— No. 154 on Map II.

On the west side of Smokehouse Branch, 0.9 mile southeast of Duo; Sewell Coal; elevation, 3574' L.

			Ft.	In.
1. Shale, with coal streaks	0'	10"		
2. Coal, draw	0	3		
3. Coal, blocky and laminated	1	3		
4. Coal, columnar.....	1	3		
5. Coal, blocky, hard, laminated	0	3		
6. Coal, hard, some bone (slate floor)	0	8	4	6

A sample (No. 154PH) was taken from Nos. 2, 3, 4, 5, and 6 of the above section and its analysis is published under

No. 154 in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. 23—
No. 155 on Map II.**

On the east side of Smokehouse Ridge, 2.1 miles east of Duo and 5.35 miles northeast of Anjean; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3816' L.

	Ft.	In.
Coal (slate roof and floor)	3	3

**Raine Lumber and Coal Company Prospect—
No. 156 on Map II.**

On the waters of Oldhouse Branch, 2.3 miles east of Duo and 0.9 mile west of Job Knob; Sewell Coal; elevation, 3752' L.

	Ft.	In.
Coal, thickness reported by Joe Raine as.....	3	8

Lemuel Hellems Mine—No. 157 on Map II.

Farm mine, on the west side of Peaser Ridge, 0.15 mile south of Hellem School; authority, David B. Reger (No. 1239, Nicholas Report, pages 707-8); Sewell Coal; elevation, 2770' B.

		Ft.	In.
1. Coal, soft (dark slate roof)	4' 7"		
2. Slate, dark	0 6		
3. Coal, soft (shale floor)	1 4	6	5

A sample (No. 352R) was taken from Nos. 1 and 3 of the above section and its analysis is published under No. 157 in the Table of Coal Analyses at the end of this Chapter.

Lemuel Hellems Mine—No. 158 on Map II.

Farm mine, on the west side of Peaser Ridge, 0.1 mile northwest of Hellem School; authority David B. Reger (No. 1238, Nicholas Report, page 707); Sewell Coal; elevation, 2725' B.

		Ft.	In.
Coal, soft (dark shale roof).....	3' 6"		
Shale, dark	0 6		
Coal, bony	0 5		
Coal, soft (fire clay floor)	1 4	5	9

Gauley Coal Land Company Prospect—No. 159 on Map II.

On the west side of Peaser Branch, 2.8 miles northwest of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2730' B.

	Ft.	In.
Coal	5	2

Gauley Coal Land Company Prospect—No. 160 on Map II.

On the west side of Peaser Branch, 2.6 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2780' B.

	Ft.	In.
Coal	4	1

Gauley Coal Land Company Prospect No. 379— No. 161 on Map II.

On the west side of Peaser Branch, 2.3 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2815' B.

	Ft.	In.
Coal	4	7

Gauley Coal Land Company Prospect No. 383— No. 162 on Map II.

On the west side of Peaser Branch, 2.2 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2859' B.

	Ft.	In.
Coal	5	2

Gauley Coal Land Company Prospect No. 384— No. 163 on Map II.

On the west side of Peaser Branch, 1.45 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2962' B.

	Ft.	In.
Coal	6	0

Gauley Coal Land Company Prospect No. 385— No. 164 on Map II.

On the west side of Peaser Branch, 1.55 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2960' B.

	Ft.	In.
Coal	6	6

Gauley Coal Land Company Prospect No. 386—

Gauley Coal Land Company Prospect—No. 160 on Map II.

On the west side of Peascer Branch, 2.6 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2780' B.

Coal
Ft. 4
In. 1

Gauley Coal Land Company Prospect No. 379—

No. 161 on Map II.

On the west side of Peascer Branch, 2.3 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2815' B.

Coal
Ft. 4
In. 7

Gauley Coal Land Company Prospect No. 383—

No. 162 on Map II.

On the west side of Peascer Branch, 2.2 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2859' B.

Coal
Ft. 5
In. 2

Gauley Coal Land Company Prospect No. 384—

No. 163 on Map II.

On the west side of Peascer Branch, 1.45 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2962' B.

Coal
Ft. 6
In. 0

Gauley Coal Land Company Prospect No. 385—

No. 164 on Map II.

On the west side of Peascer Branch, 1.55 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2960' B.

Coal
Ft. 6
In. 6

Gauley Coal Land Company Prospect No. 386—

No. 165 on Map II.

Gauley Coal Land Company Prospect—No. 160 on Map II.

On the west side of Peaser Branch, 2.6 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2780' B.

	Ft.	In.
Coal	4	1

Gauley Coal Land Company Prospect No. 379— No. 161 on Map II.

On the west side of Peaser Branch, 2.3 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2815' B.

	Ft.	In.
Coal	4	7

Gauley Coal Land Company Prospect No. 383— No. 162 on Map II.

On the west side of Peaser Branch, 2.2 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2859' B.

	Ft.	In.
Coal	5	2

Gauley Coal Land Company Prospect No. 384— No. 163 on Map II.

On the west side of Peaser Branch, 1.45 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2962' B.

	Ft.	In.
Coal	6	0

Gauley Coal Land Company Prospect No. 385— No. 164 on Map II.

On the west side of Peaser Branch, 1.55 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2960' B.

	Ft.	In.
Coal	6	6

Gauley Coal Land Company Prospect No. 386— No. 165 on Map II.

**Gauley Coal Land Company Prospect No. 387—
No. 166 on Map II.**

On the west side of Peaser Branch, 1.7 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3112' B.

			Ft.	In.
Coal	1'	1"		
Slate band		
Coal	3	9	4	10

**Gauley Coal Land Company Prospect No. 388—
No. 167 on Map II.**

On the west side of the head of Peaser Branch, 1.8 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3170' B.

			Ft.	In.
Coal	1'	2"		
Slate	0	4		
Coal	3	10	5	4

**Gauley Coal Land Company Prospect No. 389—
No. 168 on Map II.**

On the head of Peaser Branch, 1.9 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3202' B.

			Ft.	In.
Coal			5	1

**Gauley Coal Land Company Prospect No. 392—
No. 169 on Map II.**

On the head of Peaser Branch, 2 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3245' B.

			Ft.	In.
Coal	1'	4"		
Slate	1	0		
Coal	2	10	5	2

**Gauley Coal Land Company Prospect No. 394—
No. 170 on Map II.**

On the head of Peaser Branch, 2.4 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3213' B.

			Ft.	In.
Coal	1'	3"		
Slate	0	1½		
Coal	2	10	4	2½

A sample (No. 85PH) was taken from Nos. 2, 3, 4, and 5 of the above section and its analysis is published under No. 40 in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. 200—
No. 41 on Map II.**

On the south side of Meadow Creek, 0.6 mile southeast of Quinwood; Gauley Coal Land Company authority for this section, old opening; Sewell Coal; elevation, 3115' B.

	Ft.	In.
Coal	4	11

**New River & Pocahontas Consolidated Coal Company
Mine No. 2, Abandoned—No. 42 on Map II.**

On the south side of Meadow Creek, 0.3 mile southwest of Quinwood; Sewell Coal; elevation, 3059' B?

	Ft.	In.
Coal, worked out, thickness reported	6	6

The above mine was known as the "Nelson No. 2" of the Nelson Fuel Company prior to 1929.

Mines Nos. 43, 46, and 54 are intereconnected and their production is reported to the Department of Mines under the Leslie Mine. Mine No. 46 was known as the "Nelson No. 1" of the Nelson Fuel Company prior to 1929.

Main office post-office address, Fayetteville, W. Va.; mine post-office address and shipping point, Leslie; mine superintendent, E. H. Marrs; on Nicholas, Fayette, and Greenbrier Railroad.

**New River & Pocahontas Consolidated Coal Company
Mine No. 3—No. 43 on Map II.**

On the head of Little Fork of Meadow Creek, 0.65 mile southeast of Quinwood; Sewell Coal; elevation, 3135' L.

Location of sample; main heading at property line.

		Ft.	In.
1. Coal, soft, laminated (slate roof)	1' 5"		
2. Slate parting	0 9		
3. Coal, hard, columnar	2 4		
4. Coal, laminated with fusain (mineral charcoal)	1 0		
5. Coal, soft, columnar	1 6		
6. Coal, bony (not mined)	1 2	8	2

A sample (No. 96PH) was taken from Nos. 1, 3, 4, and 5 of the above section, the analysis of which is published under No. 43 in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. 196—
No. 44 on Map II.**

On the south side of Little Fork of Meadow Creek, 0.6 mile south of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3100' B.

	Ft.	In.
Coal	7	0

**Gauley Coal Land Company Prospect No. 195—
No. 45 on Map II.**

On the south side of Little Fork of Meadow Creek, 3.5 miles northeast of Charmco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3016' B.

		Ft.	In.
Coal	1' 0"		
Slate	0 2		
Coal	5 0	6	2

**New River and Pocahontas Consolidated Coal Company
Mine No. 1—No. 46 on Map II.**

On the south side of Little Fork of Meadow Creek, 3.45 miles northeast of Charmco and 0.6 mile southwest of Quinwood; Sewell Coal; elevation, 2996' L.

Location of sample; 5th right, No. 4 room.

		Ft.	In.
1. Coal, laminated with bone (slate roof)	0' 8"		
2. Coal, hard, columnar	1 10		
3. Coal, soft, laminated with fusain (mineral charecoal)	1 2		
4. Coal, soft, columnar	0 10		
5. Bone	0 2		
6. Coal, soft, laminated	1 0		
7. Coal, laminated with bone..	0 8	6	4

A sample (No. 92PH) was taken from Nos. 2, 3, 4, and 6 of the above section, the analysis of which is published under No. 46 in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. 192—
No. 47 on Map II.**

On the south side of Little Fork of Meadow Creek, 3.35 miles northeast of Charmeo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2976' B.

			Ft.	In.
Coal	6'	2"		
Slate	1	0		
Coal	0	7	7	9

**Gauley Coal Land Company Prospect No. 191—
No. 48 on Map II.**

On the east side of Meadow Creek, 3.2 miles northeast of Charmeo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2947' B.(?)

			Ft.	In.
Coal	5'	1"		
Slate	0	7		
Coal	0	7	6	3

**Gauley Coal Land Company Prospect No. 190—
No. 49 on Map II.**

On the east side of Meadow Creek, 2.6 miles northeast of Charmeo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3020' B.(?); calculated from mine map, 2970'.

		Ft.	In.
Coal		4	6

**Gauley Coal Land Company Prospect No. 188—
No. 50 on Map II.**

On the east side of Meadow Creek, 2.25 miles northeast of Charmeo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3016' B.

			Ft.	In.
Coal	0'	10"		
Slate	0	6		
Coal	4	3	5	7

Joe Neff Mine—No. 51 on Map II.

Truck mine on Snowden Crane property; on Laurel Creek Mountain, 1 mile south of Bellburn and 1.7 miles north of Charmeo; Sewell Coal; elevation, 3065' B.

		Ft.	In.
1. Coal, hard, laminated, and blocky (slate roof)	0'	11½"	

			Ft.	In.
2. Shale parting, with coal streaks	0	4		
3. Coal, columnar, soft.....	1	1		
4. Fusain (mineral charcoal)..	0	0½		
5. Coal, laminated bright and dull	0	11½		
6. Coal, laminated, soft.....	0	4		
7. Coal, hard	0	2		
8. Shale parting, (reported by Joe Neff)	1	0		
9. Coal, (reported by Joe Neff)	1	3	6	1½

A sample (No. 157PH) was taken from Nos. 1, 3, 4, 5, 6, and 7 of the above section, the analysis of which is published under No. 51 in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. 187—
No. 52 on Map II.**

On the west side of Laurel Creek, 2.35 miles northeast of Charmco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3051' B.

	Ft.	In.
Coal	6	4

**Gauley Coal Land Company Prospect No. 186—
No. 53 on Map II.**

On the north side of the head of Laurel Creek, 2.9 miles northeast of Charmco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3088' L.

	Ft.	In.
Coal	5	0

**New River and Pocahontas Consolidated Coal Company
Mine No. 4—No. 54 on Map II.**

Near the head of Laurel Creek, 2.85 miles northeast of Charmco; Sewell Coal; elevation, 3120' L.

Location of sample; 3rd right, No. 1 room.

			Ft.	In.
1. Bone, (slate roof)	0'	3"		
2. Coal, soft, laminated	1	5		
3. Coal, hard, columnar	1	11		
4. Coal, soft, laminated	0	9		
5. Coal, soft, columnar	0	11		
6. Bone	0	1	5	4

Three samples were taken from the above section; No. 93PH from No. 2 of section; No. 94PH from No. 3 of section, and No. 95PH from Nos. 4 and 5 of section. The analyses of these samples are published under No. 54 in the Table of Coal Analyses at the end of this Chapter.

**Gauley Coal Land Company Prospect No. 185—
No. 55 on Map II.**

On the east side of Laurel Creek, 2.4 miles northeast of Charmco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3104' B.

			Ft.	In.
Coal	5'	2"		
Slate	0	4		
Coal	0	7	6	1

**Gauley Coal Land Company Prospect No. 184—
No. 56 on Map II.**

On the east side of Laurel Creek, 2.5 miles northeast of Charmco; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3159' B.

			Ft.	In.
Coal			4	0

**Gauley Coal Land Company Prospect No. 181—
No. 57 on Map II.**

On Mill Creek Mountain, 0.25 mile southwest of Big Branch School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3243' B.

			Ft.	In.
Coal	4'	0"		
Coal and slate	0	8	4	8

**Gauley Coal Land Company Prospect No. 180—
No. 58 on Map II.**

On the east side of Mill Creek Mountain 0.25 mile north of Big Branch School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3243' B.

			Ft.	In.
Coal	1'	2"		
Slate	1	4		
Coal	4	6	7	0

**Gauley Coal Land Company Prospect No. 179—
No. 59 on Map II.**

On the west side of Mill Creek, 0.6 mile northeast of Blg Branch School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3253' B.

			Ft.	In.
Coal	2'	1"		
Slate	3	9		
Coal	3	11		
Coal and slate	1	8	11	5

**Gauley Coal Land Company Prospect No. 178—
No. 60 on Map II.**

On the west side of Mill Creek, 1 mile northeast of Blg Branch School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3237' B.

			Ft.	In.
Coal	4'	0"		
Coal and slate	2	0	6	0

**Gauley Coal Land Company Prospect No. 167—
No. 61 on Map II.**

On the east side of Mill Creek, 1.4 miles northeast of Blg Branch School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3280' B.

		Ft.	In.
Coal		5	4

**Gauley Coal Land Company Prospect No. 164—
No. 62 on Map II.**

On the east side of Mill Creek, 1.1 miles northeast of Big Branch School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3263' L.

			Ft.	In.
Coal	5'	1"		
Coal and slate	0	10	5	11

**Gauley Coal Land Company Prospect No. 163—
No. 63 on Map II.**

On the east side of Mill Creek, 1 mile northeast of Big Branch School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3281' L.

			Ft.	In.
Coal	4'	6"		
Coal and slate, laminated	1	6	6	0

Gauley Coal Land Company Prospect No. 162—
No. 64 on Map II.

On the east side of Mill Creek, 1.35 miles northeast of Big Branch School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3320' L.

	Ft.	In.
Coal	4	0

Gauley Coal Land Company Prospect No. 159—
No. 65 on Map II.

On the west side of Rich Knob, 0.9 mile east of Big Branch School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3345' L.

	Ft.	In.
Coal	4'	1"
Coal and slate	1	9
	5	10

Gauley Coal Land Company (?) Prospect No. 158—
No. 66 on Map II.

On the south end of Rich Knob, 1.2 miles southeast of Big Branch School; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3395' L.

	Ft.	In.
Coal	4'	0"
Slate	1	6
Coal	2	0
	7	6

Gauley Coal Land Company Prospect No. 157—
No. 67 on Map II.

On the southeast side of Rich Knob, 0.7 mile west of the mouth of Brown Creek; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3422' L.

	Ft.	In.
Coal	3	4

Gauley Coal Land Company Prospect No. 156—
No. 68 on Map II.

On the east side of Rich Knob, 0.75 mile northwest of the mouth of Brown Creek; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3387' L.

	Ft.	In.
Coal	3	9

The total output from mines and openings Nos. 69, 92, 93, 94, 95, and 96 are given in the tables of production statistics at the beginning of this Chapter under the "Leekie" mine.

**Leekie Smokeless Coal Company "Big Mountain" Mine—
No. 69 on Map II.**

On the west side of Brown Creek, 1.2 miles northwest of mouth; section as shown on mine map 600' W. of mouth of main entry; Sewell Coal; elevation, 3357' L.

			Ft.	In.
Coal	3'	4"		
Bone	0	5		
Coal	2	2	5	11

**Gauley Coal Land Company Prospect No. 154—
No. 70 on Map II.**

On the west side of Brown Creek, 1.4 miles north of mouth; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3374' B.

			Ft.	In.
Coal	3'	4"		
Coal and slate	0	1		
Coal	1	9	5	2

**Gauley Coal Land Company Prospect No. 153—
No. 71 on Map II.**

On the west side of Brown Creek, 1.8 miles northwest of mouth and 0.75 mile southeast of Sumac Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3335' L.

			Ft.	In.
Coal	0'	11"		
Sandstone	0	4½		
Coal	4	1		
Coal and slate	1	0	6	4½

**Gauley Coal Land Company Prospect No. 152—
No. 72 on Map II.**

On the west side of Brown Creek, 2.1 miles northwest of mouth and 0.5 mile southeast of Sumac Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3335' B.

			Ft.	In.
Coal	0'	8"		
Sandstone	0	1½		
Coal	4	0		
Coal and slate	1	0	5	9½

**Gauley Coal Land Company Prospect No. 151—
No. 73 on Map II.**

On the west side of Brown Creek, 2.4 miles northwest of mouth and 0.25 mile southeast of Sumac Knob; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3331' B.

			Ft.	In.
Coal	4'	0"		
Coal and slate	1	0	5	0

**Leckie Smokeless Coal Company Mine No. ?—
No. 74 on Map II.**

On the west side of Brown Creek, 2.7 miles north of mouth and 0.3 mile southwest of Brier Knob; Sewell Coal; elevation, 3353' L.

		Ft.	In.
Coal, reported 3' 0" to.....		4	0

**Gauley Coal Land Company Prospect No. 145—
No. 75 on Map II.**

On the west side of Brown Creek, 2.7 miles north of mouth; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3420' B.

			Ft.	In.
Coal	3'	6"		
Coal and slate	1	2	4	8

**Gauley Coal Land Company Prospect No. 144—
No. 76 on Map II.**

On the west side of Brown Creek, 3.1 miles north of mouth; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3420' L.

	Ft.	In.
Coal	4	4

**Gauley Coal Land Company Prospect No. 143—
No. 77 on Map II.**

On the west side of Brown Creek, 3.4 miles north of mouth; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3355' B.

	Ft.	In.
Coal	4	5

**Gauley Coal Land Company Prospect No. 142—
No. 78 on Map II.**

On the west side of Brown Creek, 3.6 miles north of mouth; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3349' B.

	Ft.	In.
Coal	3	1

**Gauley Coal Land Company Prospect No. 140—
No. 79 on Map II.**

On the west side of Brown Creek, 4 miles north of mouth; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3352' B.

		Ft.	In.
Coal	1' 11"		
Slate	3 0		
Coal	2 3	7	2

**Gauley Coal Land Company Prospect No. 138—
No. 80 on Map II.**

On the west side of Huggins Ridge near the head of Brown Creek, 2.6 miles northeast of the mouth of Sam Creek; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3348' L.

	Ft.	In.
Coal	3	1

The topography as shown on the topographic map in the vicinity of Brown Creek does not conform to conditions found there. While topographic revision of this and adjoining areas has now been completed it was not possible to obtain the corrected editions in time for this report. An attempt was made to map the outcrop of the Sewell Coal in this region with regard to its correct areal position. Attention is called to the fact that the elevations of this outcrop line and of the prospect points do not conform to the elevations shown on the base map.

**Gauley Coal Land Company Prospect No. 136—
No. 81 on Map II.**

On the west side of Huggins Ridge, 2.45 miles northeast of the mouth of Sam Creek; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3406' L.

	Ft.	In.
Coal	2	9

**Gauley Coal Land Company Prospect No. 135—
No. 82 on Map II.**

On the west side of Huggins Ridge, 2.35 miles northeast of the mouth of Sam Creek; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3427' L.

	Ft.	In.
Coal	3	2

**Gauley Coal Land Company Prospect No. 134—
No. 83 on Map II.**

On the west side of Huggins Ridge, 2.25 miles northeast of the mouth of Sam Creek; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3448' L.

	Ft.	In.
Coal	3	2

**Gauley Coal Land Company Prospect No. 133—
No. 84 on Map II.**

On the east side of Huggins Ridge, 2.05 miles northeast of the mouth of Sam Creek; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3503' L.

		Ft.	In.
Slate	0' 1½"		
Coal	1 10½	2	0

**Gauley Coal Land Company Prospect No. 132—
No. 85 on Map II.**

On the east side of Huggins Ridge, 2 miles northeast of the mouth of Sam Creek; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3475' L.

	Ft.	In.
Coal	3	8

**Gauley Coal Land Company Prospect No. 131—
No. 86 on Map II.**

On the west side of Pollock Mountain, 3.9 miles north of Anjean; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3496' L.

		Ft.	In.
Bone	0' 7"		
Coal	4 2	4	9

**Gauley Coal Land Company Prospect No. 130—
No. 87 on Map II.**

On the west side of Pollock Mountain, 3.75 miles north of Anjean; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3540' L.

	Ft.	In.
Coal	3	1

**Gauley Coal Land Company Prospect No. 129—
No. 88 on Map II.**

On the west side of Pollock Mountain, 3.65 miles north of Anjean; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3576' L.

		Ft.	In.
Bone	1' 6"		
Coal	4 1	5	7

**Gauley Coal Land Company Prospect No. 128—
No. 89 on Map II.**

On the west side of Pollock Mountain, 3.3 miles north of Anjean; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3555' L.

	Ft.	In.
Coal	3	10

**Gauley Coal Land Company Prospect No. 127—
No. 90 on Map II.**

On the west side of Pollock Mountain, 3.2 miles north of Anjean; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3548' L.

	Ft.	In.
Coal	3	10

**Gauley Coal Land Company Prospect No. 118—
No. 91 on Map II.**

On Pollock Mountain, 1.3 miles northwest of the mouth of Sam Creek and 2.8 miles north of Anjean; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3540' L.

	Ft.	In.
Coal	4	0

Correspondence with Mr. W. W. Coleman, Chief Engineer of the Leekie Smokeless Coal Company, indicates that the elevations for mine openings Nos. 92, 93, 94, 95, and 96 as used in

making the geologic structure map are 17.57 feet too high. The correct elevations are given below.

The net result of this error is to shift the 3500-contour northeast until it goes between mines 92 and 93 instead of between 93 and 94. The 3450-contour should be moved east with a rather sharp bend to include, only, mine No. 95. The other contours are not materially affected.

**Leekie Smokeless Coal Company Mine No. 5—
No. 92 on Map II.**

On the west side of Pollock Mountain, 2.5 miles north-northwest of Anjean; section as shown on mine map; Sewell Coal; elevation, 3519.3', less 17.57'=3501.73' L.

	Ft.	In.
Coal	5	1

**Leekie Smokeless Coal Company Mine No. 4—
No. 93 on Map II.**

On the northwest side of Pollock Knob, 2.25 miles north-northwest of Anjean; section as shown on mine map; Sewell Coal; elevation, 3511.5', less 17.57'=3493.93' L.

	Ft.	In.
Coal	4	8

**Leekie Smokeless Coal Company Mine No. 3—
No. 94 on Map II.**

On the west side of Pollock Knob, 2.1 miles north-northwest of Anjean; section as shown on mine map; Sewell Coal; elevation, 3482.3', less 17.57'=3464.73' L.

	Ft.	In.
Coal	4	6

**Leekie Smokeless Coal Company Mine No. 2—
No. 95 on Map II.**

On the west side of Pollock Knob, 1.95 miles north-northwest of Anjean; Sewell Coal; elevation, 3467.2', less 17.57'=3449.63' L.
Location of sample; room number 11 off air-course.

		Ft.	In.
1. Coal, hard, dull	0' 4"		
2. Coal, columnar, soft.....	1 3		
3. Coal, medium-hard (lumps well)	1 7		
4. Coal, soft	2 3		
5. Coal, jet-black bands with thin streaks of bone (not mined)	1 1	6	6

A sample (No. 80PH) was taken from Nos. 1, 2, 3, and 4, of the above section, and its analysis is published under No. 95 in the Table of Coal Analyses at the end of this Chapter.

Mine office and shipping point, Anjean; Chief Engineer, W. W. Coleman; mine superintendent, C. C. Wilburn; on Nicholas, Fayette, and Greenhrier Railroad.

**Leckie Smokeless Coal Company Mine No. 1—
No. 96 on Map II.**

On the southwest side of Pollock Knob, 1.7 miles north-northwest of Anjean; section as shown on mine map; Sewell Coal; elevation, 3478.5', less 17.57' = 3460.93' L.

	Ft.	In.
Coal	5	3

**Gauley Coal Land Company (?) Prospect—
No. 97 on Map II.**

On the east side of Pollock Mountain, 2.35 miles north of Anjean; section as shown on Gauley Coal Land Company map; Sewell Coal; elevation, 3515' L.

	Ft.	In.
Coal	4	5

**Gauley Coal Land Company (?) Prospect—
No. 98 on Map II.**

On the east side of Pollock Mountain, 2.5 miles north of Anjean; section as shown on Gauley Coal Land Company map; Sewell Coal; elevation, 3523' L.

	Ft.	In.
Coal	4	0

**Gauley Coal Land Company Prospect No. 117—
No. 99 on Map II.**

On the east side of Pollock Mountain, 1.3 miles northwest of the mouth of Sam Creek; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3551' L.

	Ft.	In.
Coal	4	6

**Gauley Coal Land Company Prospect No. 116—
No. 100 on Map II.**

On the east side of Pollock Mountain, 1.4 miles northwest of the mouth of Sam Creek; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3574' L.

	Ft.	In.
Coal	2	9

**Gauley Coal Land Company Prospect No. 115—
No. 101 on Map II.**

On the east side of Pollock Mountain, 1.7 miles north of the mouth of Sam Creek; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3562' L.

			Ft.	In.
Coal	4'	0"		
Coal and bone	1	0	5	0

**Gauley Coal Land Company Prospect No. 113—
No. 102 on Map II.**

On the west bank of Sam Creek, 2.1 miles north of its mouth and 1.85 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3514' L.

			Ft.	In.
Coal			3	0

**Gauley Coal Land Company Prospect No. 112—
No. 103 on Map II.**

On the west bank of Sam Creek, 1.8 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3497' L.

			Ft.	In.
Coal	3'	6"		
Slnto	0	6		
Coal	0	2	4	2

**Gauley Coal Land Company Prospect No. 111—
No. 104 on Map II.**

On the west bank of Sam Creek, 1.9 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3483' L.

			Ft.	In.
Coal	3'	0"		
Bone	0	2	3	2

**Gauley Coal Land Company Prospect No. 110—
No. 105 on Map II.**

On the west bank of Sam Creek, 1.9 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3487' L.

			Ft.	In.
Bone	0'	6"		
Coal	3	4½	3	10½

**Gauley Coal Land Company Prospect No. 109—
No. 106 on Map II.**

On west side of Sam Ridge, 1.75 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3492' L.

			Ft.	In.
Bone	1'	6½"		
Coal	3	2½	4	9

**Gauley Coal Land Company Prospect No. 108—
No. 107 on Map II.**

On the west side of Sam Ridge, 1.7 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3429' L? (3492').

			Ft.	In.
Coal	3'	2"		
Slate	0	3		
Coal	0	6	3	11

**Gauley Coal Land Company Prospect No. 107—
No. 108 on Map II.**

On the west side of Sam Ridge, 1.5 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3458' L.

			Ft.	In.
Coal	0'	2"		
Slate	0	2		
Coal	3	4		
Bone	1	10	5	6

**Gauley Coal Land Company Prospect No. 106—
No. 109 on Map II.**

On the west side of Sam Ridge, 1.4 miles northwest of Duo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3448' L.

			Ft.	In.
Slate	0'	9"		
Coal	3	6		
Bone	2	5	6	8

			Thlekness.		Total.		
			Ft.	ln.	Ft.	In.	
Shale, gray, sandy			19	0	183	0	
Sandstone, light-gray, hard			10	0	193	0	
Shale, dark, soft			1	8	194	8	
Coal, 3" bone near top	4'	9"	No. 6 Poca- hontas (2951')	7	4	202	0
Shale, dark, soft	1	7					
Coal	1	0					
Fire elay, small seams of coal and elay.....			4	9	206	9	

**Bellwood Coal Company Coal Test Boring No. 3—
No. 150 on Map II.**

Fayette County, Quinlmont District; 2.7 miles south of Bellwood and 2.2 miles northwest of Springdale; drilled in March, 1928; elevation, 3196.97' L.

tion, 3196.97' L.	Thickness.	Total.
Pottsville Series (201'+)	Feet.	Feet.
Surface	7.50	7.50
Sandstone, brown	21.50	29.00
Sandstone, very hard, light-gray	34.00	63.00
Sandstone, light-gray	37.00	100.00
Sandstone, dark-gray, coarse	1.05	101.05
Coal, No. 7 Pocahontas	1.03	102.08
Shale, fire clay	1.70	103.78
Sandstone, light-gray, coarse	13.82	117.60
Fire clay and shale	1.50	119.10
Slate	0.67	119.77
Coal and fire clay.....	1.58	121.35
Shale, gray, sandy	12.75	134.10
Fire clay and shale	5.50	139.60
Sandstone, light, coarse	4.50	144.10
Shale, gray, sandy	4.08	148.18
Slate and fire clay	1.25	149.43
Bone and fire clay	0.67	150.10
Fire clay and shale	5.08	155.18
Shale, gray, sandy	3.42	158.60
Bone and coal	0.50	159.10
Shale, gray, sandy	12.48	171.58
Coal	0.97	172.55
Bone	0.04	172.59
Coal	0.42	173.01
Bone	0.08	173.09
Fire clay, crumbly shale	0.12	173.21
Fire clay, light slaty shale	12.21	185.42
Shale, dark, slaty	2.00	187.42

		Thickness. Feet.	Total. Feet.
Coal	0.46'		
Bone	0.14		
Coal	3.04		
Bone	0.17		
Coal	0.10		
Bone	0.17		
Shale, fire clay, slaty ...	2.00		
Coal	0.08		
Bone and slate	0.08	No. 6 Poca- hontas (3000') ...	10.33
Coal	0.17		
Bone	0.08		
Slate	1.17		
Coal	1.29		
Shale, slaty	0.39		
Coal	0.14		
Bone and slate	0.21		
Fire clay, slaty.....	0.58		
Coal	0.06		
Fire clay, shaly.....		4.35	201.10

**Bellwood Coal Company Coal Test Boring No. 11—
No. 151 on Map II.**

Fayette County, Quinnimont District; 2.1 miles northwest of Springdale, and 0.95 mile south of Quinton School; started, June 29, 1935; completed, July 8, 1935; elevation, 3144.2' L.

Pottsville Series—Pocahontas Group (145.5'+)			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Sand boulders and yellow clay			5	0	5	0
Sandstone, yellow ...	4'	0"	Flattop	0	61	0
Sandstone	3	0				
Sandstone, hard, gray	15	0				
Sandstone, hard	5	0				
Sandstone, broken ...	14	0				
Sandstone	15	0				
Slate, blue			7	6	68	6
Coal, No. 7 Pocahontas?			4	2	72	8
Shale, dark, sandy			3	3	75	11
Coal			0	1	76	0
Sandstone, conglom- erate	5'	0"	Pierpont	0	92	0
Shale, light, sandy....	2	0				
Sandstone, gray	9	0				
Coal			0	1	92	1
Fire clay			1	5	93	6
Coal			2	0	95	6
Fire clay and shale			5	6	101	0
Shale, dark			4	0	105	0
Slate, black, with coal partings.....			2	0	107	0
Shale, dark, sandy			9	0	116	0
Shale, hard, sandy			6	6	122	6
Coal			2	0	124	6
Shale, gray, sandy			7	6	132	0

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, dark, slick	2	6	134	6
Coal and bone, No. 6 Pocahontas (3005')	4	9	139	3
Shale, fire clay, and coal	6	3	145	6

**Bellwood Coal Company Coal Test Boring No. 2—
No. 152 on Map II.**

Fayette County, Quinnimont District; 2.6 miles south of Bellwood and 1.8 miles northwest of Springdale; drilled in March, 1928; elevation, 3195.96' L.

		Thickness.	Total.
Pottsville Series (184'+)		Feet.	Feet.
Surface		16.00	16.00
Coal, Fire Creek		1.08	17.08
Sandstone, hard	35.42'	P'neville?	69.50
Sandstone	17.00'		
Shale, dark		0.42	69.92
Coal, No. 8 Pocahontas?		0.58	70.50
Fire clay and shale		3.00	73.50
Shale, sandy		4.50	78.00
Sandstone		7.58	85.58
Coal		0.34	85.91
Fire clay and soft shale		0.42	86.33
Fire clay		0.42	86.75
Fire clay and soft shale		0.58	87.33
Fire clay and shale		7.67	95.00
Shale, sandy		12.42	107.42
Shale, soft, sandy, and fire clay.....		1.00	108.42
Coal		0.91	109.33
Shale, soft, and fire clay		0.50	109.83
Shale, sandy		3.42	113.25
Shale and fire clay, soft		4.67	117.92
Shale, sandy		4.18	122.10
Shale, dark, sandy		1.25	123.35
Shale, soft, dark		1.25	124.60
Sandstone, light		1.75	125.35
Shale, dark, soft, crumbly		6.79	132.14
Sandstone		0.50	132.64
Shale, soft, dark, crumbly		3.48	136.12
Shale, light, sandy		5.50	141.62
Sandstone		7.25	148.87
Shale, sandy		16.76	165.63
Shale, soft, dark, slaty, and crumbly		2.17	167.80
Coal	0.25'	No. 6 Poca- hontas (3019')	176.72
Bone	0.29		
Coal	3.71		
Shale, light, slaty	2.27		
Coal	1.29		
Shale, slaty, and fire clay	0.91		
Coal	0.20		
Fire clay, slaty		1.55	178.27
Shale, slaty		5.45	183.72

MINABLE COALS, NEW RIVER GROUP OF
POTTSVILLE SERIES.

SEWELL COAL.

The Sewell Coal previously discussed in Chapter VI, pages 229-234, is the uppermost minable coal bed in the New River Group in Greenbrier County. It is usually multiple-bedded, soft and columnar, with a thickness varying from 2 to 9 feet. The coal lumps fairly well when mined and its very low ash and low sulphur content make it an excellent domestic fuel. The volatile matter ranges from 23 to 28 per cent. In B. T. U. it usually is above 14,500 and may exceed 15,000. The fusion temperature of the ash appears to vary somewhat but is generally between 2,100° F. and 2,700° F. Commercial production from this seam began in 1910 and has continued to date.

The Sewell bed is by far the most continuous member of the Pottsville Series of this county and its position with respect to sea-level is indicated on Map II by the green structure contours. The outcrop of this seam, outlined in blue, and the location of numerous prospect openings and mines are also given on Map II. Figure 17 shows the probable area of minable Sewell Coal.

FIGURE 17

MAP


OF

GREENBRIER COUNTY

SHOWING

PROBABLE AREA OF MINABLE

SEWELL COAL



Sewell Coal, Meadow Bluff District.

In this district, the Sewell Coal was noted in the Duo Section as 3½ feet thick, in the Charmco Section as 6.1 feet thick, and in Coal Test Borings Nos. 5E, 5J, 5K, 5L, 5M, 6, 7, 8, 9, 10, and 11, the details of which have been given on preceding pages.

The Summarized Records of Borings, on pages 386-389, as also the detailed core test records for Fayette and Nicholas Counties, will give additional information regarding this coal in adjacent areas.

All of the Sewell Coal now produced in the county is mined in Meadow Bluff District. The large commercial mines on Meadow Creek and Big Clear Creek produced a total of 1,768,016 tons from this seam in 1936.

In the extreme western part of the county the Sewell Coal is of doubtful value. Little or no prospecting has been done in that area but judging from cores and exposures in Nicholas and Fayette Counties the Sewell Coal will probably be found to be thin and split with partings. The westernmost openings are as follows:

Abraham Nutter Mine—No. 10 on Map II.

Farm mine, located 0.35 mile west of Nutter School and 0.9 mile east of Nutterville; **Sewell Coal**; elevation, 2735' B.

	Ft.	In.
Coal, reported, 1' 9" to.....	1	10

The following mine was previously reported by Reger on page 709 in the Nicholas County Report:

Johnson Nutter Mine—No. 11 on Map II.

Farm mine, on a branch of Anglins Creek, 1.8 miles S. 87° E. of Nutterville; **Sewell Coal**; elevation, 2705' B.

	Ft.	In.
Fallen shut, coal, reported, 1' 7" to.....	1	9

Pascual and James Nutter Mine—No. 12 on Map II.

Farm mine, 0.65 mile east of Nutter School and 1.8 miles east of Nutterville; **Sewell Coal?**; elevation, 2770' B.

	Ft.	In.
Coal, reported good, with thickness of.....	1	10

Two and one-half miles southeast on Burdette Creek the Gauley Coal Land Company's map shows the following:

**Gauley Coal Land Company Prospect No. 252—
No. 13 on Map II.**

On the head of Burdette Creek, 3.9 miles north-northeast of Charmco; Sewell Coal; elevation, 2863' B.

			Ft.	In.
Coal	2'	7"		
Coal and slate.....	0	8		
Coal	1	0	4	3

**Gauley Coal Land Company Prospect No. 251—
No. 14 on Map II.**

On the south side of Burdette Creek, 2.2 miles northwest of Charmco; Sewell Coal; elevation, 2820' B.

			Ft.	In.
Coal	2'	4"		
Coal and slate.....	2	0		
Coal	1	1	5	5

**Gauley Coal Land Company Prospect No. 245—
No. 15 on Map II.**

On the waters of Burdette Creek, 1.8 miles northwest of Charmco; Sewell Coal; elevation, 2896' B.

			Ft.	In.
Coal	2'	4"		
Coal and slate.....	0	4		
Slate	1	5		
Coal	1	2	5	3

One and one-half miles southwest the following three sections were measured by Price:

Haines Mine—No. 16 on Map II.

On the west side of Beegarden Knob, 3 miles north-

			Ft.	In.
3. Coal, bony, bright, laminated with bone.....	3	0		
4. Coal, good (slate floor).....	0	6	5	1

A sample (No. 158PH) was collected from No. 2 of section, the analysis of which is given under **No. 16** in the Table of Coal Analyses at the end of this Chapter.

E. M. Boyer Mine—No. 17 on Map II.

Farm mine, 1.2 miles southeast of Bingham and 2.6 miles northwest of Charmeo; Sewell Coal; elevation, 2915' B.

			Ft.	In.
Bone, dull (shale roof; fossil collection 146)	0'	4"		
Coal, good, laminated.....	1	6	1	10
Shale and bony coal.....				

H. J. and W. A. Pitzenbarger Mine—No. 18 on Map II.

Farm mine, on the east side of Beargarden Knob 2.4 miles northwest of Charmeo; Sewell Coal; elevation, 2885' B.

			Ft.	In.
1. Shale roof, (fossil collection 145).....		
2. Coal, hard, dull, bony, grading downwards into clean coal	0'	4"		
3. Coal, blocky	2	0	2	4
4. Coal, and shale floor, thickness undetermined.....				

A sample (No. 159PH) was collected from Nos. 2 and 3 of section, the analysis of which is published under **No. 18** in the Table of Coal Analyses at the end of this Chapter.

The following sections show the eastward thinning of the parting:

Gauley Coal Land Company Prospect No. 225— No. 19 on Map II.

On the north side of Meadow Creek, 0.8 mile west of Bellburn and 2.85 miles north-northwest of Charmeo; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2917' B.

			Ft.	In.
Coal	2'	11"		
Slate	0	1½		
Coal	1	2	4	2½

**Gauley Coal Land Company Prospect No. 224—
No. 20 on Map II.**

On the north side of Meadow Creek, 0.85 mile west of Bellburn and 2.6 miles north-northwest of Charmico; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2931' B.

			Ft.	In.
Coal	2'	6"		
Bone and slate	0	6		
Coal	1	2	4	2

**Gauley Coal Land Company Prospect No. 223—
No. 21 on Map II.**

On the north side of Meadow Creek, 1.05 miles southwest of Bellburn and 2.25 miles north-northwest of Charmico; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2946' B.

			Ft.	In.
Coal	2'	7"		
Bone and slate	0	5		
Coal	1	5	4	5

**Gauley Coal Land Company Prospect No. 222—
No. 22 on Map II.**

On the north side of Meadow Creek, 1.1 miles southwest of Bellburn and 2 miles north-northwest of Charmico; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2963' B.

			Ft.	In.
Coal	1'	5"		
Bone and slate	0	6		
Coal	1	5	3	4

**Gauley Coal Land Company Prospect No. 221—
No. 23 on Map II.**

On the northwest side of Meadow Creek, 0.5 mile southwest of Bellburn; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2941' B.

	Ft.	In.
Coal	4	1

**Gauley Coal Land Company Prospect No. 220—
No. 24 on Map II.**

On the northwest side of Meadow Creek 0.2 mile southwest of Bellburn; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2959' B.

	Ft.	In.
Coal	4	0

Greenbrier Smokeless Coal Company "Crichton No. 2" Mine— No. 25 on Map II.

Formerly known as "Bellburn" and prior to 1927 as the "Greenbrier" mine; on the northwest side of Meadow Creek, 0.4 mile north-northwest of Beliburn; Sewell Coal; elevation of mine entry, 2921' L.

Location of section; No. 1 Entry, 1st left, 2nd panel, room No. 9; elevation at point of sampling, 2913' L.

		Ft.	In.
1. Coal, hard, columnar (slate roof)	0' 10½"		
2. Coal, soft, laminated with fusain (mineral charcoal)	1 6½		
3. Coal, laminated with bone, (discarded in mining).....	0 8		
4. Coal, soft, columnar	1 1½	4	2½

A sample (No. 90PH) was taken from Nos. 1, 2, and 4 of section, the analysis of which is given under **No. 25** in the Table of Coal Analyses at the end of this Chapter.

Location of section; No. 2 Entry, 4th right, 6th panel, room No. 17; elevation at point of sampling, 2924' L.

		Ft.	In.
1. Coal, columnar, hard (slate roof)	1' 1"		
2. Coal, soft, laminated with fusain (mineral charcoal)	1 10		
3. Slate, (discarded in mining)	0 4½		
4. Coal soft, laminated with "mother of coal".....	0 11	4	2½

A sample (No. 91PH) was taken from Nos. 1, 2, and 4 of section, the analysis of which is given under **No. 25** in the Table of Coal Analyses at the end of this Chapter.

Post-office address, Crichton; shipping point, Beliburn; superintendent of mine, J. B. Penman; on Nicholas, Fayette, and Greenbrier Railroad.

Gauley Coal Land Company Prospect No. 217— No. 26 on Map II.

On the northwest side of Meadow Creek, 0.1 mile north of Beliburn; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 2941' B.

	Ft.	In.
Coal	4	5

Gauley Coal Land Company Prospect No. 216—
No. 27 on Map II.

On the northwest side of Meadow Creek, 0.4 mile northeast of
Bellburn; Gauley Coal Land Company authority for this section; Sewell
Coal; elevation, 2959' B.

	Ft.	In.
Coal	3	6

Johnstown Coal & Coke Company "Crichton No. 1" Mine—
No. 28 on Map II.

Prior to 1927, known as the Meadow Creek Coal Company; on
the north side of Meadow Creek at Crichton; Sewell Coal; elevation
of mine opening, 2965' L.

Location of sample; 10th right off air-coarse, 1st parallel, 3600' 5"
E. of N. of mine entry; elevation at point of sampling, 2875' L.

		Ft.	In.
1. Coal, medium-hard, colum- nar (slate roof)	1' 3"		
2. Coal, soft, laminated with fusain (mineral charecoal)	2 3		
3. Bone parting	0 11		
4. Coal, soft	0 9	5	2

A sample (No. 88PH) was taken from Nos. 1, 2, and 4
of section, the analysis of which is given under No. 28 in the
Table of Coal Analyses at the end of this Chapter.

Location of sample; 9th right off main heading, room No. 11; 700'
S. W. of No. 88PH; elevation at point of sampling, 2864.39' L.

		Ft.	In.
Coal, medium-hard, columnar (slate roof)	1' 10"		
Coal, soft, laminated with fusain (mineral charecoal)	1 0		
Coal, soft, columnar.....	0 7		
Coal, hard, some bone at base (1" to 2")	1 0	4	5

A sample (No. 89PH) was taken from the above section
and its analysis is published under No. 28 in the Table of Coal
Analyses at the end of this Chapter.

Shipping point and post-office address, Crichton; superintendent
of mine, J. B. Penman; on Nicholas, Fayette, and Greenbrier Railroad.

**Gauley Coal Land Company Prospect No. 212—
No. 29 on Map II.**

On the north side of Meadow Creek, 0.35 mlle west of Quinwood;
Gauley Coal Land Company authority for this section; Sewell Coal;
elevation, 2990' B.

			Ft.	In.
Coal	0'	7"		
Slate	0	3		
Coal	0	3		
Slate	0	11		
Coal	5	4	7	4

The output from the following openings of the Imperial
Smokeless Coal Company is given in the tables of coal pro-
duction under the "Quinwood" mine.

**Imperial Smokeless Coal Company Mine No. 1 (Pony)—
No. 30 on Map II.**

On the north side of Meadow Creek, 0.25 mlle east of Quinwood;
section from mine map; Sewell Coal; elevation, 3016' L.

	Ft.	In.
Coal, with a little bony coal at top and bottom	7	0

**Imperial Smokeless Coal Company Mine No. 1—
No. 31 on Map II.**

On the north side of Meadow Creek, 0.25 mlle east of Quinwood;
Sewell Coal; elevation, of mine mouth, 3012.3' L.

Location of sample; right main; between 14th and 15th right;
elevation of sample, 2866' L.

			Ft.	In.
Coal, soft, with thin bony part- ings (slate roof)	0'	6"		
Coal, hard	0	10		
Coal, soft, columnar.....	1	8		
Coal, medium-hard	1	1		
Coal, hard, columnar	1	8		
Coal, hard	0	7	6	4

A sample (No. 81PH) was taken from the above section
and its analysis is published under **No. 31** in the Table of
Coal Analyses at the end of this Chapter.

**Imperial Smokeless Coal Company Mine No. 2 (Pony)—
No. 32 on Map II.**

On the north side of Meadow Creek, 0.6 mlie east of Quinwood;
section from mino map; Sewell Coal; elevation, 3030' L.

	Ft.	In.
Coal	6	0

**Imperial Smokeless Coal Company Mine No. 2—
No. 33 on Map II.**

On the north side of Meadow Creek, 0.6 mile east of Quinwood;
Sewell Coal; elevation of mine opening, 3030' L.

Location of section; head of 9th right off main entry.

	Ft.	In.
Coal, hard, good (slate roof)....	1'	2"
Coal, medium-hard, columnar....	1	11
Coal, hard, laminated with fusain (mineral charcoal) (lumps well)	1	0
Coal, soft, columnar	2	6
	6	7

A sample (No. 82PH) was taken from the above section
and its analysis is published under **No. 33** in the Table of
Coal Analyses at the end of this Chapter.

Post-office address and shipping point, Quinwood; superintendent
of mine, V. A. Summerfield; on Nicholas, Fayette, and Greenbrier
Railroad.

**Frances Coal Company "Frances" Mine No. 1—
No. 34 on Map II.**

On the north side of Meadow Creek, 1.35 miles northeast of Quin-
wood; Sewell Coal; elevation at mine opening, 3160' B.

Location of section; 2nd north, 3 panel, room 9.

	Ft.	In.
Coal, hard, columnar (slate roof)	1'	3"
Coal, soft, laminated with fusain (mineral charcoal) (lumps well)	1	8
Coal, soft, columnar	0	7
Coal, medium-hard (slate floor)	1	5
	4	11

A sample (No. 86PH) was taken from the above section
and its analysis is published under **No. 34** in the Table of
Coal Analyses at the end of this Chapter.

The Frances Coal Company has ceased operation (1936) and the mine has reverted to the owner, the Gauley Coal Land Company. The coal is largely exhausted but it is reported that there is still some recoverable coal on the property.

**Gauley Coal Land Company Prospect No. 206—
No. 35 on Map II.**

On the north side of the head of Meadow Creek, 1.7 miles northeast of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3221' B.

	Ft.	In.
Coal	3	10

**Gauley Coal Land Company Prospect No. 205—
No. 36 on Map II.**

On the head of Meadow Creek, on the west side of Big Clear Creek Mountain, 1.75 miles east of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3284' B.

	Ft.	In.
Coal	4	0

**Gauley Coal Land Company Prospect No. 204—
No. 37 on Map II.**

On the south side of the head of Meadow Creek, 1.4 miles east of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3197' B.

	Ft.	In.
Coal	1'	2"
Slate	0	5
Coal	5	1
	6	8

About 3000 feet in from the mouth of the following mine a parting was noted about 14 inches from the top of the coal. This parting thickens to such an extent that the upper bench of coal is unrecoverable in some parts of the mine:

**Margarette Coal Corporation "Margarette" Mine No. 2—
No. 38 on Map II.**

Successor to the Margarette Coal Company; located on the south side of Meadow Creek, 0.85 mile east of Quinwood; Sewell Coal; elevation of mine opening, 3125' B.(?)

Location of sample; 2nd east at 3rd south.

	Ft.	In.
1. Coal, bony, laminated (slate roof, poor)	1'	2"

			Ft.	In.
2. Slate	4	0		
3. Coal, medium-hard, columnar	1	3		
4. Coal, soft, laminated (lumps well)	1	5		
5. Coal, soft, columnar	1	11		
6. Coal, bony, laminated (slate floor)	0	10	10	7

A sample (No. 84PH) was taken from Nos. 3, 4, and 5 of the above section, and its analysis is published under No. 38 in the Table of Coal Analyses at the end of this Chapter.

Post-office address and shipping point, Marfrance; mine superintendent, G. B. Staley; on Nicholas, Fayette, and Greenbrier Railroad.

Gauley Coal Land Company Prospect No. 201— No. 39 on Map II.

On the south side of Meadow Creek, 0.65 mile southwest of Quinwood; Gauley Coal Land Company authority for this section; Sewell Coal; elevation, 3133' B.

	Ft.	In.
Coal	6	0

It is reported that the Burley Coal Company is operating the following mine under a sub-lease from the Margarette Coal Corporation and that the production in 1936 is credited under the Margarette Coal Corporation. In 1934 and 1935, however, its production was separately reported by the Department of Mines under the Burley Coal Company, Burley Mine:

Margarette Coal Corporation "Margarette" Mine No. 1— No. 40 on Map II.

On the south side of Meadow Creek, 0.55 mile southeast of Quinwood; Sewell Coal; elevation, 3120' B.

Location of section; 1st panel, room 6.

			Ft.	In.
1. Coal, bony	0'	1"		
2. Coal, soft, laminated with fusain (mineral charcoal)	0	10		
3. Coal, hard, columnar	1	11		
4. Coal, soft, laminated (lumps well)	1	0		
5. Coal, soft, columnar	1	7		
6. Coal, bony (not mined).....	0	4	5	9

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Bone and coal, No. 3 Pocahontas? (2469').....	0	2	508	10
Sandstone	2	2	511	0

The partial record of boring No. 128 may be found in the table of Summarized Records at the beginning of this chapter. The complete record was not secured.

New River & Pocahontas Consolidated Coal Company Coal Test Boring No. 12—No. 129 on Map II.

Fayette County, Quinnimont District; just east of Rock of Ages School on Laurel Creek; elevation, 2872' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (439' +)				
Surface	16	0	16	0
Shale, dark	6	0	22	0
Shale, dark, sandy	0	3	22	3
Sandstone	0	1	22	4
Shale, dark	0	5	22	9
Sandstone	0	2	22	11
Shale, dark, sandy	1	5	24	4
Sandstone	3	4	27	8
Shale, dark, sandy	36	4	64	0
Shale, dark	16	4	80	4
Shale, dark, sandy	1	4	81	8
Shale, dark	1	0	82	8
Shale, dark, sandy	0	3	82	11
Shale, dark	0	2	83	1
Bone and coal	0	2	83	3
Shale, light, sandy	6	6	89	9
Shale, dark, sandy	4	6	94	3
Sandstone	6	6	100	9
Shale, light, sandy	1	0	101	9
Sandstone	1	4	103	1
Shale, sandy, light and dark	10	0	113	1
Shale, dark	16	6	129	7
Bone and coal, Beckley? (2742').....	0	1	129	8
Shale, dark	32	0	161	8
Bone	0	1	161	9
Shale, dark	0	3	162	0
Bone	0	2	162	2
Soapstone and shale, light	3	10	166	0
Shale, sandy, light and dark	36	8	202	8
Sandstone	1	0	203	8
Shale, dark	50	3	253	11
Shale, dark, with coal streaks	1'	0"	No. 8 Pocahontas (2616')	
Soapstone and shale, dark	0	8		
Shale, dark, with coal streaks	0	6		
Soapstone and shale, light	8	2	264	3

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Sandstone	25'	11"	44	2	308	5
Sandstone, with coal seams	4	2				
Sandstone	14	1				
Flat top						
Bone and coal, No. 7 Pocahontas (2563').....			0	5	308	10
Soapstone			0	3	309	1
Sandstone			0	5	309	6
Shale, dark and light.....			2	1	311	7
Sandstone			8	0	319	7
Shale, light			1	1	320	8
Sandstone			5	7	326	3
Shale, sandy, light and dark			23	2	349	5
Bone and coal.....			0	3	349	8
Sandstone and coal, mixed.....			0	4	350	0
Soapstone			0	4	350	4
Shale, light			0	6	350	10
Shale, sandy, light and dark			10	9	361	7
Slate, dark			0	4	361	11
Bone and coal			0	3	362	2
Shale, dark			13	3	375	5
Bone and coal, No. 6 Pocahontas? (2496').....			0	10	376	3
Shale, light and dark			5	1	381	4
Shale, dark			2	1	383	5
Bone and coal			0	4	383	9
Soapstone and light shale			7	0	390	9
Shale, dark			1	3	392	0
Bone and coal, No. 6 Pocahontas? (2474').....			5	9	397	9
Soapstone and light shale			11	6	409	3
Shale, dark			4	6	413	9
Bone and coal.....			0	3	414	0
Shale, dark, and coal			0	5	414	5
Shale, dark, sandy			11	6	425	11
Bone and coal			0	6	426	5
Shale, dark			0	6	426	11
Soapstone and dark shale.....			4	1	431	0
Sandstone			8	0	439	0

The partial records of borings Nos. 130 and 131 may be found in the table of Summarized Records at the beginning of this chapter. The complete records were not secured.

New River & Pocahontas Consolidated Coal Company Coal Test Boring No. 8—No. 132 on Map II.

Fayette County, Quinnimont District; on Bear Branch of Lanrel Creek, 0.9 mile west of Walnut Flat School; elevation, 2741' L.

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Pottsville Series (233'+)						
Surface			7	6	7	6
Shale, light			1	0	8	6
Shale, dark			11	6	20	0
Shale, light, sandy			9	0	29	0
Shale, dark			15	7	44	7
Coal, Fire Creek? (2697').....			0	2	44	9

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Fire clay	1	3	46	0
Shale, light	3	0	49	0
Shale, light, sandy	2	4	51	4
Shale, dark	24	0	75	4
Sandstone	0	4	75	8
Shale, dark	0	1	75	9
Sandstone 18' 6" } Pineville	38	4	114	1
Shale, dark 0 6				
Sandstone 19 4				
Sandstone and coal streaks	1	5	115	6
Coal, No. 9 Pocahontas	0	4	115	10
Sandstone	1	0	116	10
Coal and slate.....	0	4	117	2
Sandstone	0	7	117	9
Coal, No. 8 Pocahontas? (2623')	0	5	118	2
Sandstone	18	8	136	10
Shale, dark, sandy	0	8	137	6
Sandstone and coal.....	1	3	138	9
Sandstone	4	10	143	7
Coal	0	3	143	10
Soapstone	1	8	145	6
Shale, light	7	1	152	7
Sandstone	30	2	182	9
Coal	0	3	183	0
Soapstone	1	3	184	3
Shale, dark	3	1	187	4
Shale, light	2	0	189	4
Sandstone	12	4	201	8
Shale, dark, sandy	6	10	208	6
Shale, dark	2	5	210	11
Coal 0' 4" } No. 6 Poca-				
Bone 0 1				
Coal 3 4				
Bone 0 4½				
Coal 0 7½				
Shale, dark.....	0	6	216	2
Sandstone	13	4	229	6
Shale, dark	0	6	230	0
Coal and bone.....	0	1	230	1
Shale, dark	0	2	230	3
Coal and bone	0	7	230	10
Soapstone	1	11	232	9

New River & Pocahontas Consolidated Coal Company Coal Test Boring No. 9—No. 133 on Map II.

Fayette County, Quinnimont District; on Bear Branch of Laurel Creek, 1 mile northeast of Red Spring; elevation, 2719' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (335' +)				
Surface	15	0	15	0
Shale, dark, sandy	24	9	39	9
Sandstone	8	6	48	3
Shale, dark	6	10	55	1
Coal and bone, Fire Creek? (2663').....	0	8	55	9

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, dark	4	6	60	3
Shale, dark, sandy	11	0	71	3
Shale, dark	9	0	80	3
Coal and bone, Little Fire Creek? (2638').....	0	2	80	5
Shale, dark and light	17	10	98	3
Shale, dark, sandy	20	6	118	9
Coal and bone, No. 8 Pocahontas (2600').....	1	0	119	9
Clay and soapstone	0	5	120	2
Shale, dark	2	3	122	5
Shale, dark, sandy	5	5	127	10
Sandstone, Flat top and Pierpont	72	0	199	10
Coal, No. 6 Pocahontas (2518').....	1	4	201	2
Soapstone and shale, dark	9	3	210	5
Sandstone, Eckman	35	7	246	0
Sandstone and shale, mixed	0	10	246	10
Sandstone	0	6	247	4
Shale, dark	0	1	247	5
Coal, No. 5 Pocahontas? (2471').....	0	10	248	3
Shale, dark	7	9	256	0
Sandstone	5	5	261	5
Shale, dark	0	11	262	4
Sandstone	0	4	262	8
Shale, dark, sandy	1	4	264	0
Shale, dark	41	1	305	1
Bone coal, No. 3 Pocahontas	0	2	305	3
Shale, dark	4	0	309	3
Shale, light	1	2	310	5
Shale, dark	3	6	313	11
Shale, dark, sandy	2	8	316	7
Sandstone	1	8	318	3
Shale, dark	0	8	318	11
Sandstone	13	9	332	8
Coal and sandstone	0	2	332	10
Sandstone	2	2	335	0

New River & Pocahontas Consolidated Coal Company Coal Test Boring No. 13—No. 134 on Map II.

Fayette County, Quinnimont District; on Red Spring Creek, 0.2 mile east from Red Spring; elevation, 2844' L.

Pottsville Series (421'+)	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Surface	18	2	18	2
Sandstone	33	8	51	10
Shale, dark, sandy	3	9	55	7
Sandstone	1	0	56	7
Shale, dark, sandy	18	0	74	7
Shale, light	5	5	80	0
Shale, dark, sandy	4	2	84	2
Shale, dark	34	0	118	2
Bone and coal.....	0	2	118	4
Soapstone and light shale	2	5	120	9
Shale, dark, sandy	22	4	143	1

			Thiekness.		Total.	
			Ft.	In.	Ft.	In.
Shale, dark			6	6	149	7
Sandstone			0	3	149	10
Shale, dark, sandy			4	2	154	0
Bone and coal			0	5	154	5
Soapstone			1	0	155	5
Shale, dark			12	10	168	3
Sandstone			0	2	168	5
Shale, dark			0	10	169	3
Sandstone			0	4	169	7
Shale, dark			0	4	169	11
Bone and coal			0	2	170	1
Soapstone and light shale			4	0	174	1
Shale, dark			16	7	190	8
Coal	1'	1"	2	0	192	8
Bone	0	1				
Slate	0	8				
Bone and coal	0	2				
Shale, dark			2	9	195	5
Shale, dark, sandy			0	9	196	2
Shale, dark			3	6	199	8
Shale, dark, sandy			7	7	207	3
Shale, dark			9	7	216	10
Shale, dark, sandy			0	6	217	4
Shale, dark			1	10	219	2
Sandstone			0	5	219	7
Shale, dark			8	2	227	9
Shale, dark, sandy			2	7	230	4
Sandstone			35	9	266	1
Shale, dark, sandy			0	3	266	4
Shale, dark and coal	0'	1"	1	8	268	0
Sandstone and coal..	1	7				
Sandstone						
Shale, dark			33	4	301	4
Shale, dark			0	4	301	8
Sandstone			5	4	307	0
Sandstone and coal..	6'	1"	6	10	313	10
Bone and coal	0	9				
Shale, dark						
Shale, dark, sandy			6	2	320	0
Shale, dark			6	1	326	1
Shale, dark			1	1	327	2
Bone and coal			0	3	327	5
Shale, dark			0	6	327	11
Soapstone			0	7	328	6
Shale, dark			6	0	334	6
Shale, dark, sandy			7	2	341	8
Sandstone			12	4	354	0
Shale, dark			3	2	357	2
Shale, dark, sandy			1	1	358	3
Sandstone			3	9	362	0
Shale, dark, sandy			3	0	365	0
Sandstone			16	10	381	10
Sandstone and coal			0	1	381	11
Sandstone			16	2	398	1
Shale, dark			12	0	410	1
Bone and coal, No. 3 Pocahontas (2434')			0	4	410	5
Shale, light			3	2	413	7
Shale, dark, sandy			7	7	421	2

New River & Pocahontas Consolidated Coal Company Coal Test **Boring No. 7—No. 135 on Map II.**

Fayette County, Quinnimont District; on Beellek Branch, 1.5 miles east-northeast from Red Spring; elevation, 2774' L.

Pottsville Series (332'+)	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Surface	7	0	7	0
Sandstone	2	6	9	6
Shale	2	6	12	0
Fire clay	3	6	15	6
Sandstone	3	6	19	0
Shale, dark, sandy	3	6	22	6
Sandstone	25	0	47	6
Shale, dark	0	5	47	11
Coal and bone	0	5	48	4
Fire clay	3	0	51	4
Sandstone	59	10	111	2
Shale	0	5	111	7
Coal 1' 8" } No. 7 Poca-				
Coal and slate 0 4 } hontas (2660')	2	0	113	7
Shale	7	0	120	7
Shale, sandy	18	4	138	11
Shale	3	2	142	1
Shale, sandy	2	10	144	11
Sandstone	7	2	152	1
Shale, dark, sandy	20	0	172	1
Limestone (?)	7	0	179	1
Shale	2	6	181	7
Shale, dark, sandy	6	1	187	8
Sandstone	21	1	208	9
Shale	7	5	216	2
Coal and bone No. 4 Pocahontas? (2557')	0	10	217	0
Shale	9	7	226	7
Sandstone	0	10	227	5
Shale, dark, sandy	14	6	241	11
Shale	18	7	260	6
Sandstone	19	4	279	10
Shale, sandy	3	0	282	10
Coal and bone	0	7	283	5
Shale, dark	22	4	305	9
Shale, light and dark	4	0	309	9
Shale, dark, sandy	6	0	315	9
Shale, dark	12	0	327	9
Coal and bone	0	6	328	3
Shale, light	4	3	332	6

New River & Pocahontas Consolidated Coal Company Coal Test **Boring No. 16—No. 136 on Map II.**

Fayette County, Quinnimont District; 0.6 mile south of Walnut Flat School; elevation, 2844.62' L.

Pottsville Series 142'+)	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Surface	7	4	7	4

		Thickness.		Total.	
		Ft.	In.	Ft.	In.
Sandstone		9	2	16	6
Shale, dark		0	5	16	11
Sandstone		9	7	26	6
Shale, dark		1	4	27	10
Sandstone, Flattop?		28	0	55	10
Coal, No. 7 Pocahontas? (2788')		0	10	56	8
Soapstone and light shale		13	0	69	8
Sandstone, Pierpont		62	4	132	0
Shale, dark		0	2	132	2
Coal	2'	8"	} No. 6 Poca- hontas (2708')	4	8
Bone	0	9			
Coal	1	3			
Soapstone and shale		4	10	141	8

The partial record of boring No. 137 may be found in the table of Summarized Records at the beginning of this chapter. The complete record was not secured.

New River & Pocahontas Consolidated Coal Company Coal Test Boring No. 6—No. 138 on Map II.

Fayette County, Quinnimont District; on Beellek Branch, 1.5 miles east from Red Spring; elevation, 2741' L.

		Thickness.		Total.	
		Ft.	In.	Ft.	In.
Pottsville Series (232'+)					
Surface		7	0	7	0
Sandstone		49	0	56	0
Shale, gray, sandy		10	0	66	0
Sandstone, conglomerate		52	0	118	0
Bony, No. 6 Pocahontas Coal (2622')		1	2	119	2
Shale, soft, gray		4	0	123	2
Sandstone		42	0	165	2
Shale, gray		5	0	170	2
Coal and bone		0	6	170	8
Slate, gray		30	0	200	8
Slate, dark		5	0	205	8
Slate, gray		2	0	207	8
Coal and bone		0	7	208	3
Shale, gray		5	0	213	3
Shale, sandy		10	0	223	3
Slate, gray		2	0	225	3
Coal	0'	6"	} No. 3 Poca- hontas (2513')	2	8
Bone	0	4			
Coal	1	10			
Shale, gray		4	1	232	0

The partial record of boring No. 139 may be found in the table of Summarized Records at the beginning of this chapter. The complete record was not secured.

New River & Pocahontas Consolidated Coal Company Coal Test Boring No. 24—No. 140 on Map II.

Fayette County, Quinimont District; 1.05 miles east of Red Spring and 0.25 mile northwest of Eburnean School; elevation, 2746' L.

			Thickness.		Total.		
Pottsville Series (205'+)			Ft.	In.	Ft.	In.	
Surface			4	0	4	0	
Shale, dark			28	9	32	9	
Bone and coal			0	6	33	3	
Soapstone			1	4	34	7	
Shale, dark, sandy			10	4	44	11	
Coal and dirt, No. 7 Pocahontas (2701').....			0	3	45	2	
Shale, dark, sandy			2	8	47	10	
Sandstone, Plerpont			62	4	110	2	
Coal	0'	6½"	1	8	111	10	
Coal and slate	0	5½"					
Coal	0	3					
Bone	0	1½"					
Coal	0	3½"					
			No. 6 Poca-				
			hontas (2634')				
Soapstone			3	3	115	1	
Shale, light			5	6	120	7	
Sandstone, Eckman			49	5	170	0	
Shale, dark			31	2	201	2	
Coal	0'	8"	1	5	202	7	
Bone	0	1					
Coal	0	8					
							No. 3 Poca-
			hontas (2543')				
Soapstone			2		5	205	0

The record of boring No. 141 of the Bellwood Coal Company was not secured.

The records of borings Nos. 142-152 inclusive, drilled on the property of the Bellwood Coal Company, were furnished the Survey by Mr. M. F. Peltier, Vice-President of the Peabody Coal Company, Chicago, Illinois.

Bellwood Coal Company Coal Test Boring No. 4— No. 142 on Map II.

Fayette County, Quinimont District; 1.6 miles southwest of Bellwood and 0.35 mile northwest of Quilton School; drilled in April, 1928; elevation, 3285' L.

			Thickness.		Total.
Pottsville Series—New River Group (445'+)			Feet.		Feet.
Surface			4.00		4.00
Sandstone, coarse, brown	2.00'	Lower Guyandot	34.10		38.10
Sandstone, hard, gray, light	7.10				
Sandstone, dark, soft, crumbly	23.33				
Sandstone, hard, light-gray	1.67				

		Thickness. Feet.	Total. Feet.
Shale, dark, gray, sandy		18.25	56.35
Coal, Sewell (3228')		0.25	56.60
Shale, fire clay		0.25	56.85
Shale, fire clay, light, sandy		1.33	58.18
Sandstone, light, fine, Welch		19.83	78.01
Shale, light, sandy		10.58	88.59
Coal, Welch		0.17	88.76
Bone and slato		0.50	89.26
Shale, dark, sandy		11.33	100.59
Fire clay, light, and shale		1.67	102.26
Sandstone, light, fine, Upper Raleigh		43.33	145.59
Shale, dark, crumbly, fire clay		2.20	147.79
Shale, dark, sandy, fire clay		15.33	163.12
Sandstone, light, coal spars		1.75	164.87
Shale, blue, gray		38.39	203.26
Slate, dark-blue		1.71	204.97
Shale, dark-blue		6.83	211.80
Shale, dark, black, sandy		2.00	213.80
Sandstone, very hard, quartzy, light	4.25'	38.29	252.09
Sandstone, very hard, dark, shaly	3.50		
Sandstone, very hard, gray, quartzy	16.54		
Sandstone, light-gray, coarse	14.00		
Shale, light-gray, sandy		66.12	318.21
Shale, dark, sandy		7.96	326.17
Coal, bony, Beckley?		0.50	326.67
Fire clay, shaly		0.12	326.79
Shale, light, sandy		66.79	393.58
Fire clay shale, Fire Creek Coal horizon?		0.29	393.87
Sandstone, light, coarse	23.94'	46.31	440.18
Shale, dark, sandy	0.70		
Sandstone, light, coarse	17.45		
Sandstone, light-gray, (coal spars)	4.22		
Shale, dark, sandy		3.17	443.35
Coal	1.08'	2.27	445.62
Bone	0.07		
Coal	0.25		
Sulphur	0.02		
Coal	0.19		
"Mother coal"	0.10		
Coal	0.44		
Shaly bone	0.08		
Coal	0.04		
Shale, sandy		0.12	445.74
Slato, dark		0.04	445.78
Pottsville Series—Pocahontas Group (138' +)			
Shale, sandy	17.66'	28.95	474.73
Sandstone, light, coarse	11.29		
Coal		0.25	474.98
Bone and shale		0.62	475.60
Slaty shale		0.52	476.12
Coal		0.08	476.20

	Thickness.	Total.
	Feet.	Feet.
Sulphur	0.01	476.21
Coal	0.58	476.79
Shale, sandy	16.72	493.51
Sandstone, light-gray, Plerpont	50.50	544.01
Shale, dark, sandy.....	16.55	560.56
Shale, dark, sandy, fire clay	0.19	560.75
Coal, No. 6 Pocahontas (2720').....	3.94	564.69
Shale, sandy, fire clay	1.50	566.19
Sandstone, light, coarso	9.50	575.69
Shale, light, sandy	5.02	580.71
Coal, with $\frac{1}{2}$ bone in center.....	0.58	581.29
Shale, dark, sandy, fire clay	2.33	583.62

**Bellwood Coal Company Coal Test Boring No. 6—
No. 143 on Map II.**

Fayette County, Quinnsimont District; 1.95 miles south of Bellwood and 0.55 mile southeast of Quinnton School; started, April 26, 1935; completed, May 15, 1935; elevation, 3006.32' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (200'+)				
Sand boulders and yellow clay	9	0	9	0
Boulders and sandy clay	12	0	21	0
Sand boulders and yellow clay	15	0	36	0
Coal, No. 6 Pocahontas? (2970').....	0	6	36	6
Shale, gray, sandy.....	5	6	42	0
Coal, bone, and slate, No. 6 Pocahontas?	4	0	46	0
Shale, gray, sandy	7	0	53	0
Coal and fire clay	3	0	56	0
Shale, dark, sandy	9	4	65	4
Shale, dark	4	8	70	0
Coal	1	0	71	0
Shale, dark	3	0	74	0
Sandstone, hard	1	0	75	0
Shale, dark	0	8	75	8
Shale, sandy, hard	6	4	82	0
Shale, dark	14	6	96	6
Shale, black	4	6	101	0
Shale, sandy	6	0	107	0
Sandstone, hard	13	0	120	0
Sandstone, Upper Pocahontas	20	6	140	6
Shale, dark	3	6	144	0
Sandstone, hard	2	9	146	9
Coal, No. 3 Pocahontas	1	6	148	3
Shale and fire clay	1	8	149	11
Coal and bone	0	9	150	8
Shale, dark	8	4	159	0
Shale, light, sandy	4	6	163	6
Shale, dark, sandy	5	6	169	0
Sandstone, hard, Lower Pocahontas	26	7	195	7
Slate, black	0	5	196	0
Coal, No. 2 Pocahontas	0	9	196	9
Shale, blue, sandy	3	3	200	0

Bellwood Coal Company Coal Test Boring No. 1— No. 144 on Map II.

Fayette County, Quinnimont District; 2 miles south of Bellwood and 2.1 miles northwest of Springdale; drilled in February, 1928; elevation, 3155.35' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—New River Group (64'±)				
Clay	8	0	8	0
Sandstone, hard, light-gray, Pineville	35	0	43	0
Shale, dark-gray, sandy	3	0	46	0
Sandstone, dark-gray	18	0	64	0
Coal, very soft, No. 8 Pocahontas	0	1½	64	1½
Pottsville Series—Pocahontas Group (364')				
Sandstone, dark, hard	0	3½	64	5
Shale, dark, sandy	4	5½	68	10½
Sandstone, dark-gray, coarse, Flattop	33	8	102	6½
Coal	0	8½	103	2¾
Slate	0	2½	103	4½
Coal	0	2½	103	6¾
Shale, dark, fire clay seams	3	0	106	6¾
Coal, bony, dark	0	2	106	8¾
Shale, dark	0	7½	107	4
Shale, sandy	9	5½	116	9½
Slate, with fire clay	1	0	117	9½
Coal	1	1	118	10½
Fire clay	0	1½	118	11¾
Shale, dark, sandy	38	1¾	157	1 (?)
Coal	0	3	157	4
Coal and bone	0	5	157	9
Shale, light	1	9¾	159	7 (?)
Shale, dark, oily	0	6	160	1
Shale, dark	2	0	162	10¾ (?)
Shale, sandy	2	10	165	8¾
Shale, light	11	1½	176	10½
Slate, draw	0	1½	176	11¾
Coal, clean, No. 6 Pocahontas (2975')	3	1½	180	1½
Shale, sandy	8	6½	188	7½ (?)
Coal	1'	0½"	No. 6 Pocahontas	5
Bone	0	1		
Coal	0	2½		
Bone	0	2½		
Coal	0	8		
Slate and fire clay....	1	10		
Coal	0	3¾		
Slate	0	2¾		
Coal	0	1¾		
Slate	0	2		
Coal	0	10		
Shale, sandy	23	6¾	217	10
Bone	0	0½	217	10½
Shale, dark	1	5	219	3½
Bone	0	2½	219	6
Coal, No. 5 Pocahontas	0	5	219	11
Shale, dark	3	9	223	8
Sandstone	5	8	229	4
Shale, sandy	5	2	234	6

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Sandstone	4	0	238	6
Shale, sandy	9	4	247	10
Shale, dark	9	0½	256	10½
Coal, No. 4 Pocahontas	2	1	258	11½
Shale, dark	3	3	262	2½
Sandstone, Upper Pocahontas	18	4	280	6½
Shale, slaty	0	1½	280	7½
Coal	0'	2"	No. 3 Poca- hontas (2783')....	
Bone	0	1		
Coal	0	11½		
Shale, dark	0	1½		
Coal	0	3		
Bone and coal	0	5½	282	8½
Shale, light	4	4½	287	1
Shale, dark	5	11	293	0
Slate and fire clay	0	8½	293	8½
Shale, dark	4	7½	298	4
Shale, gray, sandy	6	2	304	6
Sandstone, Lower Pocahontas	25	1½	329	7½
Shale, dark, slaty	0	6½	330	1½
Coal, No. 2 Pocahontas	0	6½	330	7½
Shale, gray, sandy	46	7½	377	3
Shale, dark, slaty	0	11	378	2
Coal, No. 1 Pocahontas	0	7	378	9
Shale, sandy	45	3	424	0
Fire clay and shale	0	6	424	6
Shale and fire clay, dark, sandy	3	9	428	3
Mauch Chunk Series (14'+)				
Shale, red and gray	10	6	438	9
Shale, and fire clay, soft, gray	4	0	442	9

**Bellwood Coal Company Coal Test Boring No. 7—
No. 145 on Map II.**

Fayette County, Quinnimont District; 2 miles south of Bellwood and 0.65 mile southeast of Quilton School; started, May 18, 1935; completed, May 23, 1935; elevation, 3104.85' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (143'+-)				
Surface, sand boulders, and clay	12	6	12	6
Coal, No. 8 Pocahontas	2	0	14	6
Fire clay	1	6	16	0
Sandstone, Flattop	29	7	45	7
Coal	0	5	46	0
Fire clay	1	6	47	6
Shale, dark	7	0	54	6
Coal	0	1	54	7
Shale, dark, soft	0	5	55	0
Coal	0	2	55	2
Fire clay	1	0	56	2
Shale, dark	7	6	63	8
Coal	0	2	63	10
Fire clay	0	10	64	8
Coal	0	8	65	4
Shale, dark, hard	15	8	81	0

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, dark	5	0	86	0
Coal	0	1	86	1
Shale, dark	1	3	87	4
Bone	0	4	87	8
Shale, dark	5	7	93	3
Shale, dark, soft	1	0	94	3
Shale, dark	3	0	97	3
Shale, sandy	1	5	98	8
Shale, black	0	8	99	4
Coal	1	0	100	4
Fire clay	4	2	104	6
Shale, light, sandy	9	6	114	0
Shale, with streaks of sand	4	0	118	0
Shale, with streaks of dark-gray	16	2	134	2
Coal	0'	5"	No. 6 Pocahontas (2962.5')	8
Bone	0	3		
Coal	4	2		
Shale, dark	1	0		
Bone coal	0	3		
Coal	0	2		
Shale, dark	0	3		
Coal	0	1		
Shale, dark	0	2		
Coal	0	6		
Shale, dark	0	7		
Coal	0	4		
Shale, dark	0	8	143	0

**Bellwood Coal Company Coal Test Boring No. 9—
No. 146 on Map II.**

Fayette County, Quilmont District; 2.1 miles northwest of Springdale and 0.75 mile southeast of Quiuton School; started, June 8, 1935; completed, June 17, 1935; elevation, 3155.45' L.

Pottsville Series (180'+)	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Clay, yellow	12	0	12	0
Shale, dark	3	0	15	0
Sandstone	9'	0"	Pineville	26
Shale, dark	1	0		
Sandstone	16	9		
Shale, dark, soft	5	0	46	9
Coal, No. 9 Pocahontas	0	3	47	0
Shale, sandy	8	0	55	0
Shale, dark	4	0	59	0
Coal, No. 8 Pocahontas	1	6	60	6
Sandstone, Flattop	26	2	86	8
Coal	0	4	87	0
Fire clay	3	6	90	6
Shale, dark, sandy	7	6	98	0
Shale, dark, soft	1	0	99	0
Shale, sandy	9	0	108	0
Shale, black	0	6	108	6
Shale, dark	1	6	110	0

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Coal			0	8	110	8
Bone			0	3	110	11
Coal			0	3	111	2
Fire clay			1	10	113	0
Shale, dark			7	7	120	7
Coal			0	5	121	0
Shale, dark			5	0	126	0
Coal			0	1	126	1
Shale, dark			10	11	137	0
Shale, dark, sandy			3	0	140	0
Shale, hard, streaks of sand			15	0	155	0
Coal			2	0	157	0
Fire clay			1	0	158	0
Shale, light, sandy			4	0	162	0
Shale, dark			4	9	166	9
Coal	0'	3"	No. 6 Pocahontas (2976')	4	177	1
Bone	0	3				
Coal	4	0				
Fire clay	1	3				
Coal	0	1				
Shale	0	2				
Coal	1	0				
Fire clay	0	10				
Coal	0	1				
Fire clay or soft gray shale	1	2				
Coal	0	2				
Fire clay	1	0				
Coal	0	1				
Shale, gray			2	11	180	0

**Bellwood Coal Company Coal Test Boring No. 10—
No. 147 on Map II.**

Fayette County, Quinnimont District; 1.9 miles northwest of Springdale and 0.9 mile southeast of Quintou School; started, June 20, 1935; completed, June 27, 1935; elevation, 3192' L.

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Pottsville Series (208'+)						
Clay, yellow			4	0	4	0
Shale, soft, yellow			8	0	12	0
Shale, dark			17	0	29	0
Sandstone, hard	16'	0"	Pineville	0	60	0
Sandstone, hard, broken	15	0				
Shale, dark			5	4	65	4
Coal, No. 9 Pocahontas			1	0	66	4
Fire clay			0	8	67	0
Shale, light, sandy			2	4	69	4
Coal			0	2	69	6
Sandstone			0	6	70	0
Sandstone, hard			5	6	75	6
Shale, dark, sandy			9	6	85	0
Coal, No. 8 Pocahontas			1	3	86	3

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, dark, sandy	7	9	94	0
Shale, sandy	2	0	96	0
Sandstone, light-gray, Flattop	16	0	112	0
Coal	0	9	112	9
Shale, dark	10	3	123	0
Shale, sandy	3	0	126	0
Shale, dark	2	6	128	6
Shale, black	1	0	129	6
Coal	0	10	130	4
Shale, dark	7	8	138	0
Coal	0	6	138	6
Shale, sandy	3	6	142	0
Shale, dark	10	0	152	0
Shale, sandy	5	10	157	10
Coal	0	8	158	6
Fire clay	1	6	160	0
Shale, dark	4	6	164	6
Shale, sandy	2	6	167	0
Sandstone, Pierpont	11	6	178	6
Shale, dark-blue	19	9	198	3
Coal	0'	5"	No. 6 Pocahontas (2984')	
Bone	0	3		
Coal	4	2		
Shale, dark	1	2		
Coal, bone, and slate	1	2		
Shale, dark	0	8		
Bone, coal	0	4		
Shale, dark	0	8		
Coal	0	2		
Shale, dark	0	7		
Coal	0	2		
Shale, dark			0	6
			208	6

**Bellwood Coal Company Coal Test Boring No. 8—
No. 148 on Map II.**

Fayette County, Quinnimont District; 2.1 miles northwest of Springdale and 0.75 mile south-southwest of Quilton School; started, May 27, 1935; completed, June 6, 1935; elevation, 3158.7' L.

				Thickness.		Total.	
				Ft.	In.	Ft.	In.
Pottsville Series (195' +)							
Boulders and elay				5	0	5	0
Sandstone, broken..... 7' 0"				26	0	31	0
Sandstone 19 0							
Shale, dark, with sandy streaks				23	0	54	0
Shale, dark				6	0	60	0
Coal, No. 9 Pocahontas				0	6	60	6
Shale, dark				14	6	75	0
Coal, No. 8 Pocahontas				2	0	77	0
Sandstone 7' 0"				39	6	116	6
Sandstone, light-							
gray 12 0							
Sandstone, light 14 0							
Sandstone, broken ... 6 6							

		Thickness.		Total.	
		Ft.	In.	Ft.	In.
Shale, black, soft		2	0	118	6
Fire clay		1	6	120	0
Shale, dark		17	4	137	4
Shale, sandy		1	6	138	10
Sandstone		3	6	142	4
Shale, black		0	2	142	6
Coal		0	3	142	9
Fire clay		5	3	148	0
Shale, dark		12	9	160	9
Shale, with streaks of sand		3	0	163	9
Sandstone, light-gray, Pierpont		4	0	167	9
Shale, black		2	3	170	0
Shale, dark-gray		11	0	181	0
Shale, gray, "slippery"		2	0	183	0
Coal	5' 0"	No. 6 Pocahontas (2966')	9 10	192	10
Fire clay	1 4				
Coal	0 5				
Fire clay	0 8				
Coal	0 2				
Fire clay	1 2				
Coal	0 1				
Fire clay	0 10				
Coal	0 2	1 9	194	7	
Fire clay					

**Bellwood Coal Company Coal Test Boring No. 12—
No. 149 on Map II.**

Fayette County, Quinlinton District; 2.5 miles northwest of Springdale and 1.05 miles south of Quinlinton School; started, July 17, 1935; completed, July 25, 1935; elevation, 3152.9' L.

		Thickness.		Total.	
		Ft.	In.	Ft.	In.
Pottsville Series (206'+)					
Clay, yellow		5	0	5	0
Sandstone, yellow	18' 0"	Plneville	66 0	71	0
Sandstone, light-gray,					
yellow streaks	25 0				
Sandstone, light-gray	23 0	Flat top	15 0	115	0
Shale, dark, soft					
Shale, gray, sandy		16	0	87	0
Coal, No. 8 Pocahontas		3	9	90	9
Fire clay		2	1	92	10
Shale, sandy		0	8	93	6
Sandstone	5' 0"	6	6	100	0
Sandstone, light-gray	10 0	15 0	115	0	
Shale, dark					
Coal		0	3	115	3
Shale, dark, very soft		0	3	115	6
Shale, blue, sticky		6	6	122	0
Coal		15	9	137	9
Fire clay and dark shale		1	5	139	2
Sandstone, gray		12	10	152	0
Coal		5	10	157	10
Fire clay		0	6	158	4
		5	8	164	0

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Shale			10	6	84	0
Sandstone			2	1	86	1
Shale, dark			19	3	105	4
Coal, Castle			0	2	105	6
Clay			1	3	106	9
Sandstone	6'	0"	38	9	145	6
Shale, sandy	4	0				
Sandstone	0	10				
Shale, sandy	3	7				
Sandstone	1	0				
Shale, dark	1	0				
Sandstone	3	9				
Shale	0	5				
Sandstone	18	2				
Coal and bone, Sewell "B"			0	6	146	0
Fire clay			2	0	148	0
Sandstone			4	0	152	0
Shale, sandy			17	2	169	2
Sandstone			4	0	173	2
Shale, dark			26	6	199	8
Sandstone			7	6	207	2
Shale, dark			16	10	224	0
Coal	0'	11½"	3	6½	227	6½
Parting	0	8½				
Coal	1	10½				
Slate, to bottom			1	5½	229	0

The record of boring No. 94, drilled on the property of Charles White, was not obtained.

The records of borings Nos. 95 and 96, drilled on the property of the Nuttall Heirs, were not obtained.

The record of boring No. 97, drilled on the property of Jno. Jordan-Amick, was not obtained.

The following record is reprinted from pages 446-447 of the Fayette County Report:

Beury Coal Test Boring No. 3—No. 111 on Map II.

Fayette County, Sewell Mountain District; on east bank of Laurel Creek, southeast of Pine Grove Schoolhouse, 2.7 miles S. 75° E. of Landsburg; by New River & Poehontas Consolidated Coal Company; authority; J. S. Cunningham; elevation, 2545' L.

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Pottsville Series (300'+)						
Surface			10	0	10	0
Shale			10	0	20	0
Sandstone, Lower Raleigh			90	0	110	0
Slate			0	10	110	10
Coal, Beckley (2434' L.)			0	5	111	3

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Slate			0	9	112	0
Sandstone	16'	0"	47	0	159	0
Shale	20	6				
Sandstone	10	6				
Sandstone and shale			16	0	175	0
Shale, Quinnimont			68	6	243	6
Coal, Fire Creek(?) (2297' L.)			3	10	247	4
Sandstone			6	0	253	4
Shale			8	0	261	4
Coal, Little Fire Creek(?)			1	0	262	4
Slate			2	0	264	4
Sandstone, Pineville			35	8	300	0

The records of borings Nos. 112, 113, 115, 116, 117, 118, 121, 122, 123, 124, 124A, 125, 126, 127, 129, 132, 133, 134, 135, 136, 138, and 140 are published with the permission of Mr. S. M. Wolfe, Superintendent of Lands, New River and Pocahontas Consolidated Coal Company.

New River & Pocahontas Consolidated Coal Company Coal Test Boring No. 19—No. 112 on Map II.

Fayette County, Sewell Mountain District; located on Glade Creek $\frac{1}{4}$ mile southeast of Sims School; elevation, 2641' L.

				Thickness.		Total.	
				Ft.	In.	Ft.	In.
Pottsville Series (282'+)							
Surface				4	6	4	6
• Sandstone				0	8	5	2
Shale, dark				14	5	19	7
Shale, dark, sandy				1	5	21	0
Shale, dark				15	6	36	6
Sandstone				6	0	42	6
Shale, dark				6	6	49	0
Shale, light				5	0	54	0
Shale, light, sandy				3	1	57	1
Sandstone 24' 0"				45	6	102	7
Shale, dark, sandy... 0 3							
Sandstone 21 3							
Shale, dark				24	9	127	4
Bone and coal 0' 5"				1	6	128	10
Coal 0 2½							
Bone 0 1½							
Coal 0 9							
Shale, dark				7	9	136	7
Shale, dark, sandy				11	0	147	7
Bone and coal 0' 5"				3	3	150	10
Shale, dark 2 4							
Slate and coal..... 0 6							
Shale, dark				1	1	151	11
Shale, dark, sandy				1	10	153	9
Shale, dark				6	3	160	0
Soapstone and light shale				2	0	162	0

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, sandy, light and dark	14	7	176	7
Sandstone, Pineville	10	5	187	0
Shale, dark	0	5	187	5
Sandstone	6	2	193	7
Shale, dark	0	2	193	9
Sandstone, Flattop	32	5	226	2
Shale, dark	0	7	226	9
Sandstone	8	4	235	1
Shale, dark	1	1	236	2
Sandstone, Pierpont	32	2	268	4
Shale, dark, sandy, Royal	7	10	276	2
Coal, dirty	1'	11"	No. 6 Poca- hontas (2361')	
Coal and slate	0	9		
Slate	0	6		
Coal and slate	0	5		
Shale, dark, and soapstone	2	3	279	9
			282	0

**New River & Pocahontas Consolidated Coal Company Coal Test
Boring No. 20—No. 113 on Map II.**

Fayette County, Sewell Mountain District; on Glade Creek, 1.3
miles southwest of Sims School; elevation, 2595' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (495' +)				
Surface	22	6	22	6
Shale, dark, sandy	13	2	35	8
Shale, dark	10	8	46	4
Sandstone	4	5	50	9
Shale, dark, sandy	1	6	52	3
Sandstone	16	8	68	11
Shale, dark	31	6	100	5
Sandstone	3	0	103	5
Shale, dark	0	4	103	9
Sandstone	2	1	105	10
Shale	2	8	108	6
Sandstone	0	10	109	4
Shale, dark	1	3	110	7
Sandstone, Quinnimont	14	0	124	7
Shale, dark	15	5	140	0
Bone	0	4	140	4
Coal, Fire Creek (2454')	1	0	141	4
Slate, dark	0	6	141	10
Shale, dark	5	0	146	10
Shale, dark, sandy	1	5	148	3
Shale, dark	23	0	171	3
Shale, dark, sandy	10	11	182	2
Sandstone, Pineville?	8	0	190	2
Shale, dark	0	6	190	8
Slate and coal	0	4	191	0
Soapstone and dark shale	7	1	198	1
Bone and coal, No. 8 Pocahontas	0	2	198	3
Slate, dark	2	5	200	8
Soapstone and dark shale	10	0	210	8
Sandstone	1	8	212	4

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, dark	5	4	217	8
Sandstone, Flattop and Pierpont	78	3	295	11
Shale, dark, and coal, No. 6 Pocahontas?.....	1	9	297	8
Shale, dark	7	1	304	9
Slate and coal, No. 6 Pocahontas?	1	2	305	11
Soapstone and light shale	4	7	310	6
Shale, dark	32	9	343	3
Sandstone	3	10	347	1
Shale, dark	0	11	348	0
Sandstone	3	3	351	3
Coal and slate	0'	10"	} No. 4 Poca-	hontas (2240')
Bone	0	2		
Coal	2	9		
Soapstone and dark shale.....	6	4	355	0
Shale, dark	8	8	361	4
Sandstone	1	0	370	0
Shale, dark, sandy	0	9	371	0
Sandstone	4	1	371	9
Shale, dark, sandy	16	3	375	10
Shale, dark, and soapstone	11	0	392	1
Shale, dark	3	3	403	1
Bone and coal	0'	9"	406	4
Shale, dark	3	0	} No. 3 Poca-	hontas (2184')
Bone and coal	0	8		
Shale, dark	3	10		
Shale, dark, sandy	4	5	410	9
Shale, dark	3	0	414	7
Shale, dark, sandy	2	0	419	0
Shale, dark	9	2	422	0
Shale, dark, sandy	1	0	424	0
Shale, dark	0	7	433	2
Slate and coal	0	3	434	2
Soapstone	0	3	434	9
Bone and coal	0	3	435	0
Slate, dark	0	3	435	3
Bone and coal	0	3	435	6
Shale, dark	2	0	437	6
Bone	0	1	437	7
Coal	2'	0"	} No. 2 Poca-	hontas (2155')
Bone	0	1		
Coal	0	6		
Shale, dark, and soapstone	7	7	440	2
Shale, dark	2	8	447	9
Shale, light	8	10	450	5
Shale, sandy, light, and dark	11	8	459	3
Slate	0	2	470	11
Coal, No. 1 Pocahontas	0	7	471	1
Soapstone and light shale	0	7	471	8
Shale, sandy, dark	2	4	474	0
	21	0	495	0

The partial record of boring No. 114 may be found in the table of Summarized Records at the beginning of this chapter. The complete record was not secured.

New River & Pocahontas Consolidated Coal Company Coal Test **Boring No. 22—No. 115 on Map II.**

Fayette County, Sewell Mountain District; 1.95 miles east of Danese and 2 miles west of Bellwood; elevation, 2801' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (277'+)				
Surface	24	0	24	0
Sandstone	1	2	25	2
Shale, dark	1	3	26	5
Bone and coal, Beckley	1	0	27	5
Soapstone	1	0	28	5
Shale, light	1	11	30	4
Shale, dark	7	4	37	8
Bone	0	1	37	9
Shale, light	1	11	39	8
Sandstone	59	8	99	4
Sandstone and coal mixed	10	3	109	7
Shale, dark	13	8	123	3
Coal, Fire Creek (2678')	0	3	123	6
Soapstone and light shale	6	11	130	5
Shale, dark, sandy	48	0	178	5
Shale, dark	0	7	179	0
Soapstone and light shale	3	2	182	2
Sandstone, Pineville	26	10	209	0
Shale, dark, sandy	1	0	210	0
Sandstone	1	1	211	1
Shale, dark	52	0	263	1
Bone	0	2	263	3
Coal 2' 8" } No. 7 Poca-				
Bone and coal 0 4 } hontas?	3	0	266	3
Soapstone	0	2	266	5
Shale, dark	0	6	266	11
Shale, dark, sandy	5	6	272	5
Coal 0' 5" } No. 6 Poca-				
Bone and coal 0 6 } hontas?	0	11	273	4
Soapstone	0	6	273	10
Shale, light	3	7	277	5

New River & Pocahontas Consolidated Coal Company Coal Test **Boring No. 18—No. 116 on Map II.**

Fayette County, Sewell Mountain District; on Glade Creek, 1.7 miles northeast of Danese; elevation, 2571' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (336'+)				
Surface	14	0	14	0
Sandstone, Lower Raleigh	18	0	32	0
Shale, dark, sandy	12	7	44	7
Coal, Beckley? (2526')	0	3	44	10
Soapstone and shale5	7	50	5
Shale, dark	1	0	51	5
Sandstone	0	7	52	0
Shale, dark, sandy	0	1	52	1
Sandstone	0	2	52	3

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Shale, dark			0	7	52	10
Shale, dark, sandy			34	0	86	10
Shale, dark			42	10	129	8
Sandstone, Quinlmont			35	0	164	8
Shale, dark			6	11	171	7
Coal	0'	11"	5	4	176	11
Bone	0	1				
Soapstone and shale	1	1				
Bone and coal	1	7				
Coal	1	5				
Slate and coal.....	0	3	} Fire Creek (2394')			
Soapstone			1	3	178	2
Shale, dark			7	0	185	2
Shale, dark, sandy			16	10	202	0
Coal			0	2	202	2
Slate and coal			1	1	203	3
Soapstone			0	4	203	7
Shale, dark			32	7	236	2
Bone coal and slate..	2'	2"	3	10	240	0
Slate and coal.....	1	8				
} No. 8 Pocahontas (2331')						
Soapstone			2	1	242	1
Shale, dark			30	11	273	0
Shale, dark, and sandstone			0	9	273	9
Sandstone, Flattop			10	1	283	10
Sandstone and coal			0	1	283	11
Sandstone			5	4	289	3
Shale, dark			0	9	290	0
Sandstone, Pierpont			34	10	324	10
Bone and coal, No. 6 Pocahontas (2243').....			2	10	327	8
Soapstone and light shale			8	7	336	3

**New River & Pocahontas Consolidated Coal Company Coal Test
Boring No. 2—No. 117 on Map II.**

Fayette County, Sewell Mountain District; located on the south branch of Pole Creek, 1.5 miles southeast of Danese; elevation, 2863' L.

Pottsville Series (419'+)	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Surface	17	0	17	0
Sandstone	19	0	36	0
Sandstone, red	0	10	36	10
Shale, dark	12	0	48	10
Shale, dark, sandy	9	5	58	3
Sandstone	28	0	86	3
Shale, dark	4	0	90	3
Slate, black	2	0	92	3
Shale, gray	10	0	102	3
Slate, gray	6	5	108	8
Coal, Little Raleigh	0	4	109	0
Shale, dark	20	0	129	0
Shale, sandy	3	0	132	0
Sandstone	35	6	167	6
Shale, sandy	4	6	172	0

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, dark	6	6	178	6
Slate, black, Beckley Coal horizon?.....	5	0	183	6
Shale, varlegated	1	6	185	0
Shale, gray	3	0	188	0
Sandstone	15	0	203	0
Shale, dark	12	0	215	0
Shale, sandy	10	8	225	8
Slate, gray	15	4	241	0
Shale, sandy	6	0	247	0
Sandstone	28	0	275	0
Slate, gray	16	0	291	0
Shale, gray	9	0	300	0
Slate, gray	6	0	306	0
Shale, sandy	9	0	315	0
Slate, gray	53	0	368	0
Shale, sandy	2	0	370	0
Slate, black	0	6	370	6
Bone	0	6	371	0
Slate, black	2	0	373	0
Slate, gray	12	0	385	0
Shale, sandy	14	6	399	6
Shale, dark	4	6	404	0
Shale, sandy	7	0	411	0
Sandstone, conglomerate	4	0	415	0
Shale, dark	1	2	416	2
Sandstone, conglomerate	3	6	419	8

New River & Pocahontas Consolidated Coal Company Coal Test Boring No. 1—No. 118 on Map II.

Fayette County, Sewell Mountain District; located on Smoky Branch one-half mile east of Danese; elevation, 2621' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (513'+)				
Surface	12	0	12	0
Shale, dark	36	0	48	0
Slate, gray	4	0	52	0
Shale, gray	4	0	56	0
Sandstone	2	0	58	0
Shale, gray	3	8	61	8
Coal and bone	0	4	62	0
Shale, sandy	1	0	63	0
Sandstone	91	0	154	0
Coal, Beckley (2467').....	0	6	154	6
Shale, varlegated	2	0	156	6
Sandstone	5	0	161	6
Shale, gray, sandy	10	0	171	6
Sandstone	2	0	173	6
Shale, gray, sandy	12	0	185	6
Sandstone	2	0	187	6
Shale, dark	23	0	210	6
Shale, sandy	7	0	217	6
Shale, gray	3	0	220	6
Slate, black	0	6	221	0

				Thickness.		Total.	
				Ft.	In.	Ft.	In.
Coal and bone, Fire Creek (2400').....				0	6	221	6
Shale, gray				1	6	223	0
Sandstone				15	0	238	0
Slate, gray				55	0	293	0
Slate, black				12	9	305	9
Bone	0'	4"	} No. 8? No. 9? Pocahontas (2311')	4	7	310	4
Coal	2	3					
Slate	0	10					
Coal	1	1					
Bone	0	1					
Slate, dark				10	0	320	4
Coal				0	5	320	9
Slate				0	6	321	3
Coal				1	3	322	6
Shale, gray				6	0	328	6
Bone				0	4	328	10
Shale, gray				12	0	340	10
Coal and bone				1	0	341	10
Shale, variegated				3	10	345	8
Sandstone				2	0	347	8
Shale, sandy				10	0	357	8
Sandstone, conglomerate				28	0	385	8
Bone				0	1	385	9
Sandstone, conglomerate				15	0	400	9
Shale				0	4	401	1
Coal	1'	7"	} No. 6 Poca- hontas (2218')	1	10	402	11
Bone	0	3					
Shale, gray				5	0	407	11
Shale and clay				3	6	411	5
Shale, gray				6	8	418	1
Slate, dark				32	0	450	1
Slate, gray				37	0	487	1
Sandstone, hard				26	0	513	1

The two following records were previously published on pages 447 and 448 of the Fayette County Report:

New River & Pocahontas Consolidated Coal Company Coal Test Boring No. 5—No. 119 on Map II.

Fayette County, Sewell Mountain District; on south bank of Glade Creek, 0.3 mile southeast of Pittman Schoolhouse and 1.0 mile north of Danese; authority, J. S. Cunningham; elevation, 2554' L.

				Thickness.		Total.	
				Ft.	In.	Ft.	In.
Pottsville Series (291'+)							
Surface				9	6	9	6
Shale, gray, sandy				18	6	28	0
Sandstone				14	9	42	9
Shale, gray				3	4	46	1
Slate				1	0	47	1
Bone, Little Raleigh Coal				0	7	47	8
Shale, gray				14	0	61	8

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Sandstone	17'	5"	64	6	126	2
Shale	0	9				
Sandstone, conglomerate	42	4				
Shale, soft, gray	4	0				
Lower Raleigh						
Sandstone			4	0	130	2
Shale, sandy			11	6	141	8
Shale, gray, with iron and sulphur			5	6	147	2
Slate			1	7	148	9
Coal, Beckley (2405' L.)			1	1	149	10
Shale			0	6	150	4
Shale, variegated			2	0	152	4
Shale, sandy			3	0	155	4
Shale, dark			6	0	161	4
Shale, sandy			3	0	164	4
Slate, gray			14	5	178	9
Shale, gray, sandy			18	6	197	3
Shale, gray	25'	0"	73	6	270	9
Slate	0	6				
Bone	0	8				
Slate, gray	47	4				
Quinnimont						
Slate			15	4	286	1
Coal, bone	0'	1"	3	5	289	6
Coal	3	4				
Fire Creek						
Shale, gray, to bottom			2	0	291	6

New River & Pocahontas Consolidated Coal Company Coal Test Boring No. 21—No. 120 on Map II.

Fayette County, Sewell Mountain District; on south bank of Smoky Branch at mouth of Sandy Creek, 1.0 mile northwest of Danese; location on Map II in error—belongs 0.83 mile south; authority, J. S. Cunningham; elevation, 2560.42' L.

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Pottsville Series (215' +)						
Surface			7	0	7	0
Sandstone, Lower Raleigh			24	0	31	0
Shale, light and dark			4	6	35	6
Shale, dark, sandy			12	6	48	0
Shale, dark			5	10	53	10
Coal and bone, Beckley (2506')			0	7	54	5
Shale, light, and soapstone			2	7	57	0
Sandstone	24'	0"	91	10	148	10
Shale, dark, sandy	11	8				
Sandstone	11	2				
Shale, dark, sandy	14	2				
Sandstone	30	10	61	6	210	4
Shale, dark, Quinnimont						
Coal and bone	0'	1½"	4	5½	214	9½
Coal	3	7½				
Bone	0	4½				
Coal	0	4				
(2346' L.)						
Soapstone, to bottom			0	2½	215	0

New River & Pocahontas Consolidated Coal Company Coal Test

Boring No. 17—No. 121 on Map II.

Fayette County, Sewell Mountain District; 0.35 mile east of Ebenezer School and 1.05 miles south of Danese; elevation, 2739' L.

Pottsville Series (386'±)	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Surface	3	0	3	0
Sandstone, Upper Raleigh	66	6	69	6
Shale, dark	1	6	71	0
Sandstone	1	6	72	6
Shale, dark, sandy	1	7	74	1
Sandstone	2	5	76	6
Shale, dark, sandy	1	7	78	1
Sandstone	11	6	89	7
Shale, dark	0	11	90	6
Bone and coal 0' 3" } Little				
Coal 0 10 } Raleigh "A" ...	1	1	91	7
Soapstone and light shale.....	5	3	96	10
Shale, dark, sandy	5	4	102	2
Shale, dark	2	7	104	9
Shale and coal	0	4	105	1
Soapstone and light shale	1	2	106	3
Shale, dark	1	0	107	3
Bone	0	3	107	6
Coal, Little Raleigh	0	6	108	0
Soapstone	2	2	110	2
Sandstone, Lower Raleigh	61	0	171	2
Shale, dark, sandy	21	10	193	0
Shale, dark	1	0	194	0
Shale, dark, sandy	7	6	201	6
Shale, dark	17	7	219	1
Shale, dark, sandy	9	7	228	8
Shale, dark	4	0	232	8
Sandstone	7	2	239	10
Shale, dark	6	4	246	2
Sandstone	2	10	249	0
Shale, dark, sandy	7	7	256	7
Sandstone, Quinnimont	22	2	278	9
Shale, dark	32	0	310	9
Bone and coal 0' 7" } Fire Creek				
Shale, dark 3 9 } Coal (2422')....	5	1	315	10
Bone and coal 0 9 }				
Shale, dark	18	0	333	10
Shale, dark, sandy	9	6	343	4
Shale, dark	17	7	360	11
Coal 0' 2" } No. 8 Poca-				
Bone and coal 1 0 } hontas Coal ...	3	0	363	11
Shale, dark 1 5 } (2374')				
Bone and coal 0 5 }				
Shale, dark and light	14	8	378	7
Sandstone, Flattop	7	5	386	0

448 COMMERCIAL COAL

New River & Pocahontas Consolidated Coal Company Coal Test

Boring No. 15—No. 122 on Map II.

Fayette County, Quinnimont District: 1.6 miles north-northwest from Red Spring; elevation. 2781' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (343' +)				
Surface	5	10	5	10
Sandstone	44	0	49	10
Shale, dark	2	4	52	2
Coal and dirt	1	0	53	2
Fire clay	2	0	55	2
Shale, dark	1	3	56	5
Bone and coal	0	2	56	7
Slate	0	2	56	9
Shale, light and dark	5	0	61	9
Shale, dark, sandy	12	2	73	11
Shale, dark	3	3	77	2
Coal and bone, Little Raleigh?	1	1	78	3
Slate, dark	14	0	92	3
Bone and coal	0	5	92	8
Shale, dark	0	3	92	11
Sandstone	37	6	130	5
Shale, dark, sandy	2	6	132	11
Sandstone	8	9	141	8
Shale, dark, sandy	1	3	142	11
Sandstone	1	6	144	5
Shale, dark, sandy	5	0	149	5
Sandstone	0	3	149	8
Shale, dark, sandy	18	10	168	6
Shale, dark	4	0	172	6
Soapstone and shale, light	5	6	178	0
Sandstone	32	10	210	10
Shale, dark, sandy	0	6	211	4
Sandstone	37	0	248	4
Shale, light and dark	6	2	254	6
Shale, dark, sandy	11	6	266	0
Sandstone	4	6	270	6
Shale, dark	51	2	321	8
Coal	0'	5"	No. 8 Pocahontas? (2455')	
Slate, dark	0	10		
Coal	1	11		
Bone	0	3		
Coal	0	2		
Slate and coal	0	6		
Coal	0	4		
Slate and coal	0	1	No. 8 Pocahontas? (2438')	
Soapstone and dark shale	15	4		
Coal	0'	8"		
Bone	0	1		
Coal	0	5		
Bone	0	1		
Soapstone	0	6		

New River & Pocahontas Consolidated Coal Company Coal Test **Boring No. 14—No. 123 on Map II.**

Fayette County, Quinlmont District; on Pond Branch, 1.1 miles northwest of Red Spring; elevation, 2584' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (219'+)				
Surface	10	0	10	0
Sandstone	0	9	10	9
Shale, dark	12	6	23	3
Bone and coal, Fire Creek? (2550')	0	4	23	7
Fire clay	0	1	23	8
Soapstone and shale, light and dark	3	2	26	10
Shale, dark	5	8	32	6
Shale, dark, sandy	24	0	56	6
Shale, dark	4	5	60	11
Coal 2' 8" } No. 8 Poca-				
Slate and coal 1 0 } hontas (2519')	3	8	64	7
Soapstone and shale	5	3	69	10
Shale, dark	42	1	111	11
Sandstone	35	0	146	11
Shale, dark	0	10	147	9
Sandstone	10	7	158	4
Shale, dark	0	2	158	6
Coal 0' 7" } No. 6 Poca-				
Bone and coal 0 1 } hontas (2425')	0	8	159	2
Shale, dark, sandy	12	10	172	0
Sandstone	2	7	174	7
Shale, dark	0	8	175	3
Shale, dark, and coal	0	4	175	7
Shale, dark	14	0	189	7
Shale, dark, sandy	16	2	205	9
Sandstone	13	3	219	0

New River & Pocahontas Consolidated Coal Company Coal Test **Boring No. 10—No. 124 on Map II.**

Fayette County, Quinlmont District; at mouth of Red Spring Creek, 0.95 mile north of Red Spring; elevation, 2564' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (150'+)				
Surface	10	0	10	0
Shale, dark	14	3	24	3
Shale, dark, sandy	10	9	35	0
Sandstone	2	7	37	7
Shale, dark	1	4	38	11
Sandstone	3	4	42	3
Shale, dark	3	0	45	3
Sandstone	4	1	49	4
Shale, light	12	7	61	11
Sandstone	0	11	62	10
Coal, No. 8 Pocahontas? (2500')	0	8	63	6
Sandstone, Flattop	41	0	104	6
Sandstone and coal	1	6	106	0
Sandstone	4	2	110	2
Shale	0	7	110	9

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Bone and coal	0'	9"	1	10	112	7
Slate	0	4				
Coal	0	9				
No. 7 Pocahontas (2451')						
Soapstone and shale, light			9	0	121	7
Shale, sandy, light and dark			5	0	126	7
Sandstone, Pierpont			18	0	144	7
Bone			0	2	144	9
Bone and coal	0'	6"	2	2	146	11
Slate and coal	1	8				
Shale, light						
No. 6 Pocahontas (2417')			3	1	150	0

New River & Pocahontas Consolidated Coal Company Coal Test Boring No. 11—No. 124A on Map II.

Fayette County, Quinnimont District; 0.85 mile west of Crlekmer:
elevation, 2722' L.

Pottsville Series (328'+)	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Surface	28	0	28	0
Shale, dark	11	0	39	0
Sandstone, Lower Raleigh	58	6	97	6
Shale, dark, sandy	9	6	107	0
Soapstone and fire clay	4	0	111	0
Sandstone	2	0	113	0
Shale, dark, sandy	15	0	128	0
Shale, dark	29	0	157	0
Shale, dark, sandy	22	3	179	3
Shale, dark	5	10	185	1
Coal, Fire Creek? (2536')	0	8	185	9
Shale, dark	20	4	206	1
Coal	0	8	206	9
Shale, dark, sandy	11	8	218	5
Slate	1	0	219	5
Bone and coal	0	8	220	1
Shale, dark	4	8	224	9
Shale, dark, sandy	4	8	229	5
Shale, light, sandy	4	0	233	5
Shale, dark, sandy	2	4	235	9
Sandstone	0	6	236	3
Shale, dark, sandy	4	10	241	1
Sandstone	0	6	241	7
Shale, dark	0	3	241	10
Sandstone	1	4	243	2
Shale, dark	0	6	243	8
Sandstone, Flattop	46	6	290	2
Shale, dark	6	3	296	5
Coal, No. 6 Pocahontas? (2424')	1	7	298	0
Soapstone and shale	4	8	302	8
Bone and coal	0	9	303	5
Shale, very dark	1	0	304	5
Shale, dark	10	8	315	1
Shale, dark, sandy	2	2	317	3
Shale, light, sandy	7	0	324	3
Sandstone	3	9	328	0

New River & Pocahontas Consolidated Coal Company Coal Test **Boring No. 4—No. 125 on Map II.**

Fayette County, Quinnimont District; near Crlekmer; elevation, 2865' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (459'+)				
Surface	6	6	6	6
Sandstone, conglomerate	62	0	68	6
Shale, gray	36	6	105	0
Sandstone	47	0	152	0
Shale, gray	22	0	174	0
Sandstone	56	0	230	0
Slate, gray	13	0	243	0
Sandstone	26	0	269	0
Slate, dark	58	0	327	0
Bone	0	4	327	4
Slate, gray	30	8	358	0
Shale, gray	11	0	369	0
Slate, gray	24	0	393	0
Sandstone	34	10	427	10
Slate, dark	7	6	435	4
Bone and coal	0'	2"	No. 6 Poca- hontas (2428')	
Coal	1	1		
Bone and coal	0	4		
Shale, light, sandy.....	5	0	441	11
Sandstone	17	4	459	3

New River & Pocahontas Consolidated Coal Company Coal Test **Boring No. 3—No. 126 on Map II.**

Fayette County, Quinnimont District; headwaters of Laurel Creek, one mile northeast from Crlekmer; elevation, 2904' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (532'+)				
Surface	10	6	10	6
Shale, gray	25	6	36	0
Slate, black	2	0	38	0
Coal, Little Raleigh	0	4	38	4
Shale, gray	6	6	44	10
Shale, sandy	2	4	47	2
Sandstone	2	6	49	8
Shale, gray	8	6	58	2
Sandstone	25	0	83	2
Shale, sandy	17	0	100	2
Shale, gray	14	0	114	2
Shale, sandy	19	0	133	2
Slate, black	4	0	137	2
Sandstone	41	0	178	2
Shale, dark	16	0	194	2
Shale, gray	54	6	248	8
Slate, dark	18	0	266	8
Shale, gray	5	0	271	8
Shale, variegated	1	6	273	2
Sandstone	6	6	279	8

	Thickness.		Tot.	
	Ft.	In.	Ft.	In.
Shale, dark	1	6	281	2
Sandstone	0	8	281	10
Shale, gray	4	0	285	10
Sandstone	1	0	286	10
Slate, gray	3	6	290	4
Coal, Fire Creek? (2613').....	0	4	290	8
Slate, gray	22	0	312	8
Shale, sandy	8	0	320	8
Shale, dark	5	0	325	8
Slate, dark	1	0	326	8
Bone	1	0	327	8
Slate, black	2	6	330	2
Coal	0	6	330	8
Slate	1	0	331	8
Shale, sandy	8	0	339	8
Slate, black	4	0	343	8
Bone and coal	0	9	344	5
Shale	1	3	345	8
Sandstone	2	0	347	8
Shale, sandy	1	0	348	8
Sandstone, conglomerate	74	6	423	2
Bone	0	4	423	6
Coal, No. 6 Pocahontas (2479').....	1	9	425	3
Shale	0	4	425	7
Shale, variegated	2	4	427	11
Shale, gray	4	0	431	11
Shale, sandy, dark	29	9	461	8
Shale, gray	16	6	478	2
Slate, gray	41	10	520	0
Bone	0	2	520	2
Coal, No. 3 Pocahontas (2383').....	1	1	521	3
Shale, sandy, variegated	5	3	526	6
Sandstone, conglomerate	6	4	532	10

**New River & Pocahontas Consolidated Coal Company Coal Test
Boring No. 23—No. 127 on Map II.**

Fayette County, Quinnimont District; 1.65 miles west of Quinton School and 1.2 miles northeast of Rock of Ages School; elevation, 2978' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (511'+)				
Surface	27	8	27	8
Sandstone	40	4	68	0
Shale, dark	3	3	71	3
Sandstone	4	10	76	1
Shale, dark, sandy	11	4	87	5
Sbale, dark	3	3	90	8
Sandstone	64	0	154	8
Shale, dark	3	0	157	8
Shale, light and dark	15	8	173	4
Sandstone	11	0	184	4
Shale, dark	34	6	218	10
Coal, Fire Creek? (2759').....	0	7	219	5
Soapstone	0	8	220	1

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, dark	23	4	243	5
Shale, dark, sandy	22	11	266	4
Coal	0	5	266	9
Bone	0	1	266	10
Coal (2710')	0	7	267	5
Shale, dark	15	11	283	4
Coal	0	1	283	5
Soapstone and dark shale	2	11	286	4
Coal, No. 8 Pocahontas?	0	9	287	1
Soapstone	0	8	287	9
Shale, dark	11	6	299	3
Bone and coal	0	10	300	1
Shale, dark	3	4	303	5
Bone and coal	0	4	303	9
Coal, No. 7 Pocahontas?	0	5	304	2
Shale and coal	1	2	305	4
Bone and coal	0	7	305	11
Shale	0	9	306	8
Shale and coal	1	1	307	9
Shale, dark	7	10	315	7
Shale, dark, sandy	4	4	319	11
Sandstone	1	1	321	0
Shale	0	3	321	3
Sandstone	3	9	325	0
Sandstone and coal	0	11	325	11
Sandstone	5	1	331	0
Shale, light	1	4	332	4
Shale, dark	2	9	335	1
Sandstone	45	0	380	1
Sandstone and coal	2	3	382	4
Bone	0	1	382	5
Coal and dlrt, No. 6 Pocahontas? (2595')	0	8	383	1
Soapstone and dark shale	13	0	396	1
Shale, dark	4	7	400	8
Sandstone	7	5	408	1
Shale, dark	1	3	409	4
Coal, No. 6 Pocahontas? (2569')	0	6	409	10
Soapstone and dark shale	4	6	414	4
Sandstone	0	4	414	8
Shale, dark	2	0	416	8
Shale, dark, sandy	9	1	425	9
Sandstone	2	0	427	9
Shale, dark	0	2	427	11
Sandstone	1	4	429	3
Shale, dark, sandy	16	0	445	3
Shale, dark	14	5	459	8
Coal	0	3	459	11
Soapstone	3	2	463	1
Shale, dark	3	11	467	0
Sandstone	7	9	474	9
Coal and dlrt, No. 3 Pocahontas? (2503')	0	6	475	3
Soapstone and dark shale	5	5	480	8
Sandstone	13	8	494	4
Shale, dark, sandy	3	0	497	4
Shale, dark	11	4	508	8

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, light	5	3	380	2
Sandstone	5	7	385	9
Shale, dark	10	6	396	3
Shale, light	9	7	405	10
Slate, dark	2	0	407	10
Coal, Sewell "A"	0	6	408	4
Fire clay	2	4	410	8
Shale and sandstone	6	2	416	10
Shale, dark	23	0	439	10
Coal, Sewell (1613' B.).....	2	2	442	0
Slate, streaked with coal	2	7	444	7
Fire clay	0	10	445	5

**Gauley Coal Land Co. Coal Test Boring No. 27—No. 44
on Map II.**

Nicholas County, Kentucky District; on Hughey Branch of Deer Creek, 3.15 miles northeast of Deepwell, and 3.05 miles northwest of Nettle; elevation, 2225' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—Kanawha Group (38'+)				
Surface	11	0	11	0
Sandstone	2	0	13	0
Shale and clay	8	0	21	0
Sandstone, with coal spars.....	3	8	24	8
Coal, Lower Douglas	0	1	24	9
Fire clay, sandy	2	3	27	0
Shale, dark, sandy	11	0	38	0
Pottsville Series—New River Group (412'+)				
Sandstone, Upper Nuttall	67	2	105	2
Slate, soft, black	0	4½	105	6½
Coal, dirty, laeger "B"	0	3½	105	10
Fire clay	2	10	108	8
Shale, gray	3	4	112	0
Shale, sandy	26	0	138	0
Shale, dark, and slate	38	2	176	2
Slate, with coal spars, laeger "A"	1	9	177	11
Fire clay	0	11	178	10
Shale, sandy	6	4	185	2
Sandstone	13	0	198	2
Shale, sandy	2	0	200	2
Shale, gray	12	8	212	10
Shale, dark	0	1	212	11
Shale, gray	1	5	214	4
Shale, dark	0	7	214	11
Shale, gray	6	0	220	11
Sandstone and shale	3	10	224	9
Sandstone	47	6	272	3
Coal, Lower laeger?	1	1	273	4
Sandstone	7	3	280	7
Shale, dark, with sand streaks.....	8	6	289	1
Shale, dark	13	7	302	8
Coal, Lower laeger?	1	3	303	11
Shale, dark	5	0	308	11

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, dark, sandy	3	6	312	5
Sandstone	5	0	317	5
Shale, dark	14	3	331	8
Sandstone	14	6	346	2
Shale and sandstone, dark	42	7	388	9
Sandstone	1	0	389	9
Coal, clean	1'	11"	} Castle (1833' B.)
"Mother coal"	0	1		
Coal, clean	0	3		
Fire clay	8	0	400	0
Shale, dark, sandy	4	0	404	0
Sandstone	18	4	422	4
Shale, dark	27	8	450	0

**Gauley Coal Land Co. Coal Test Boring No. 16—No. 45
on Map II.**

Nicholas County, Kentucky District; 0.5 mile west-southwest of Odell School; elevation, 2400' B.

Pottsville Series—New River Group (392'+)	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Surface	27	0	27	0
Sandstone, hard	0	10	27	10
Shale, soft, light	9	2	37	0
Shale, light, sandy	16	4	53	4
Coal, bony, laeger "A"	0	7	53	11
Shale, light	1	3	55	2
Sandstone, hard	7	2	62	4
Shale, light	18	2	80	6
Sandstone, hard	15	0	95	6
Sandstone, mottled	1	0	96	6
Sandstone, dark	0	3	96	9
Coal, Hughes Ferry	0	4	97	1
Shale, dark	0	3	97	4
Shale, light	1	6	98	10
Sandstone, hard	44	2	143	0
Shale, dark, with white streaks	15	9	158	9
Shale, dark, sandy	2	4	161	1
Coal, and slate, laminated, Lower laeger.....	10	7	171	8
Shale, dark, with white streaks	9	2	180	10
Shale, dark	33	8	214	6
Coal, dirty	0	8	215	2
Shale, gray	15	7	230	9
Fire clay, hard	2	0	232	9
Slate, black	1	3	234	0
Shale, gray, sandy	20	0	254	0
Sandstone, hard, Guyandot	67	0	321	0
Shale, gray	30	6	351	6
Slate, black	2	0	353	6
Coal, Sewell "A"	1	1	354	7
Fire clay, dark	0	6	355	1
Shale, gray	23	1	378	2
Sandstone	1	5	379	7
Shale, gray	1	8	381	3

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Slate, black	0	1	381	4
Coal, clean	2'	7½"	} Sewell (2016' B.)	
Coal, laminated.....	0	2½"		
Slate, laminated			384	2
Shale, dark	1	2	385	4
	6	8	392	0

**Gauley Coal Land Co. Coal Test Boring No. 14—No. 46
on Map II.**

Nicholas County, Kentucky District; 1.6 miles northwest of Hominy Mill and 0.75 mile southeast of Odell School; completed, June 28, 1917; elevation, 2300'±.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—New River Group (196'+)				
Surface	6	6	6	6
Sandstone, hard, Guyandot	31	0	37	6
Shale, hard, dark	33	6	71	0
Sandstone, hard, mottled	2	0	73	0
Slate, black	0	4	73	4
Coal, Sewell "A"	1	1	74	5
Shaly fire clay	1	1	75	6
Shale, dark	2	0	77	6
Sandstone, Lower Guyandot	6	0	83	6
Shale, dark, sandy	12	6	96	0
Coal	2'	3"	} Sewell (2201' ±)	
Slate parting	0	1½"		
Coal, bony	0	8½"		
Shale, dark			99	1
Shale, dark	0	4	99	5
Fire clay	3	4	102	9
Shale, dark, soft	12	3	115	0
Slate, black	7	0	122	0
Coal, dirty, Welch	1	1	123	1
Fire clay, shaly	1	5	124	6
Shale, dark	10	0	134	6
Shale, dark, sandy	45	0	179	6
Shale, dark	16	6	196	0

**Gauley Coal Land Co. Coal Test Boring No. 15—No. 46A
on Map II.**

Nicholas County, Kentucky District; 1 mile northeast of Hominy Mill and 1 mile southwest of Tolbert; completed, Aug. 2, 1917; elevation, 2600'±.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—New River Group (419'+)				
Surface	5	0	5	0
Shale, soft	6	0	11	0
Shale, soft, dark	11	6	22	6
Sandstone	26	4	48	10
Shale, variegated	28	8	77	6
Sandstone, hard	30	6	108	0
Shale, gray	42	9	150	9
Coal, Hughes Ferry	0	7	151	4

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, gray	2	4	153	8
Sandy shale	4	10	158	6
Slate, black	0	4	158	10
Shale, dark, gray	2	0	160	10
Fire clay, shaly	2	8	163	6
Shale, light, sandy	6	0	169	6
Shale, dark, sandy	20	4	189	10
Sandstone	3	0	192	10
Shale, dark	13	8	206	6
Fire clay, shaly	6	4	212	10
Shale, dark, hard, with hard sandstone streaks	18	2	231	0
Sandstone, hard, mottled	29	0	260	0
Shale, dark, sandy	2	0	262	0
Sandstone, hard	12	6	274	6
Shale, gray	14	6	289	0
Sandstone, hard, with coal streaks	26	0	315	0
Shale, gray	3	0	318	0
Sandstone, hard	16	3	334	3
Shale, light, sandy	1	6	335	9
Sandstone, hard, with coal streaks	45	3	381	0
Sandstone, with shale	27	7	408	7
Coal, clean, Sewell "A"?	1	6	410	1
Fire clay, sandy	0	6	410	7
Shale, sandy	3	10	414	5
Slate	0	3	414	8
Coal, clean, Sewell (2183'±)	2	13	416	93
Fire clay, sandy	2	24	419	0

The following record of boring No. 46B appears to be the same as the record of the depth and thickness of the Sewell Coal given for boring No. 48 as published in the Nicholas County Report, page 432. However, the location of No. 48 given in that report and shown on Map II in this report is one mile southwest of the location for No. 46B:

**Gauley Coal Land Co. Coal Test Boring No. 8—No. 46B
on Map II.**

Nicholas County, Kentucky District; on Grassy Creek, on John Collison, near Grassy Creek School; authority, Gauley Coal Land Company; elevation, 2375'±.

Pottsville Series (665'±)	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Sand, gravel, and boulders	15	0	15	0
Sandstone	19	0	34	0
Shale, dark	9	6	43	6
Sandstone	34	6	78	0
Coal	3'	9"	Sewell (2293'±)	
Bony parting	0	1		
Coal	0	6		
Fire clay	6	9	82	4
			89	1

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Slate, black	2	2	91	3
Fire clay	3	2	94	5
Shale, gray, sandy	8	2	102	7
Shale, dark	16	3	118	10
Slate, black	0	8	119	6
Coal, Welch	0	7	120	1
Slate, black	0	8	120	9
Sandstone	18	0	138	9
Shale, dark, sand streaks	20	0	158	9
Shale, dark	39	5	198	2
Coal, Little Raleigh?	0	3	198	5
Fire clay	8	0	206	5
Shale, gray	22	10	229	3
Sandstone	7	0	236	3
Shale, dark	2	6	238	9
Sandstone	9	4	248	1
Shale, dark	1	7	249	8
Sandstone	3	0	252	8
Shale, dark	3	0	255	8
Shale, gray	6	6	262	2
Sandstone	5	6	267	8
Shale, gray	9	0	276	8
Fire clay, impure	3	0	279	8
Shale, dark	15	5	295	1
Slate, black	13	11	309	0
Coal	0	6	309	6
Slate, gray	2	0	311	6
Bone	0	3	311	9
Coal, Beckley?	0	6	312	3
Fire clay	6	3	318	6
Shale, gray	3	0	321	6
Shale, dark	12	2	333	8
Sandstone	10	9	344	5
Shale, dark	1	10	346	3
Sandstone	6	5	352	8
Shale, dark	2	10	355	6
Fire clay	3	0	358	6
Shale, dark	2	6	361	0
Coal, bony, Fire Creek? (2014'±)	0	7	361	7
Fire clay	8	0	369	7
Shale, gray	3	5	373	0
Shale, dark	11	10	384	10
Shale, gray	20	0	404	10
Fire clay	6	6	411	4
Shale, gray	3	4	414	8
Shale, dark	1	6	416	2
Fire clay	7	4	423	6
Sandstone	8	0	431	6
Shale, dark	2	10	434	4
Shale, dark	1	10	435	9.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Slate, dark	3	6	484	10
Fire clay	2	8	487	6
Sandstone	6	0	493	6
Shale, gray	7	3	500	9
Slate, gray	6	3	507	0
Coal, bony	1	7	508	7
Fire clay	1	8	510	3
Shale, gray	6	0	516	3
Slate, dark	3	3	519	6
Coal, bony, No. 6 Pocahontas? (1755'±).....	0	5	519	11
Fire clay	14	0	533	11
Sandstone	30	10	564	9
Shale, dark	1	9	566	6
Sandstone	3	6	570	0
Slate, dark	16	0	586	0
Coal, bony	0	10	586	10
Fire clay, dark	1	6	588	4
Coal, bony	0	5	588	9
Fire clay	5	9	594	6
Sandstone	6	3	600	9
Shale, dark	0	9	601	6
Sandstone	0	6	602	0
Shale, dark	0	9	602	9
Sandstone	1	0	603	9
Shale, dark	37	3	641	0
Sandstone, conglomerate	0	10	641	10
Fire clay, impure.....	6	0	647	10
Shale, gray	4	0	651	10
Sandstone, conglomerate	13	2	665	0

**Gauley Coal Land Co. Coal Test Boring No. 22A—No. 46C
on Map II.**

Nicholas County, Kentucky District; 1.1 miles east of Tolbert;
elevation, 2480'±.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—New River Group (289'±)				
Surface	20	0	20	0
Sandstone	6	0	26	0
Shale, soft, light	10	0	36	0
Shale, light, sandy	39	3	75	3
Sandstone, hard, Guyandot	68	4	143	7
Shale, hard, dark, with sandstone streaks.....	37	10	181	5
Coal, clean	1'	1"	Sewell (2295'±)	0
Slate	0	0 $\frac{1}{2}$		
"Mother coal"	0	0 $\frac{1}{2}$		
Coal and thin layers of "mother coal"	1	9		
"Mother coal"	0	0 $\frac{1}{2}$		
Coal	0	0 $\frac{1}{2}$		
"Mother coal"	0	0 $\frac{1}{2}$		
Coal, clean	0	6 $\frac{1}{2}$		
Clay, sandy	4	0	189	0
Sandstone, Welch	10	5	199	5

	Thlekness.		Total.	
	Ft.	In.	Ft.	In.
Slate, black	2	2	91	3
Fire clay	3	2	94	5
Shale, gray, sandy	8	2	102	7
Shale, dark	16	3	118	10
Slate, black	0	8	119	6
Coal, Welch	0	7	120	1
Slate, black	0	8	120	9
Sandstone	18	0	138	9
Shale, dark, sand streaks	20	0	158	9
Shale, dark	39	5	198	2
Coal, Little Raleigh?	0	3	198	5
Fire clay	8	0	206	5
Shale, gray	22	10	229	3
Sandstone	7	0	236	3
Shale, dark	2	6	238	9
Sandstone	9	4	248	1
Shale, dark	1	7	249	8
Sandstone	3	0	252	8
Shale, dark	3	0	255	8
Shale, gray	6	6	262	2
Sandstone	5	6	267	8
Shale, gray	9	0	276	8
Fire clay, impure	3	0	279	8
Shale, dark	15	5	295	1
Slate, black	13	11	309	0
Coal	0	6	309	6
Slate, gray	2	0	311	6
Bone	0	3	311	9
Coal, Beckley?	0	6	312	3
Fire clay	6	3	318	6
Shale, gray	3	0	321	6
Shale, dark	12	2	333	8
Sandstone	10	9	344	5
Shale, dark	1	10	346	3
Sandstone	6	5	352	8
Shale, dark	2	10	355	6
Fire clay	3	0	358	6
Shale, dark	2	6	361	0
Coal, bony, Fire Creek? (2014'±)	0	7	361	7
Fire clay	8	0	369	7
Shale, gray	3	5	373	0
Shale, dark	11	10	384	10
Shale, gray	20	0	404	10
Fire clay	6	6	411	4
Shale, gray	3	4	414	8
Shale, dark	1	6	416	2
Fire clay	7	4	423	6
Sandstone	8	0	431	6
Shale, dark	2	10	434	4
Sandstone	1	10	436	2
Shale, gray	15	10	452	0
Slate, black	0	6	452	6
Coal	0	6	453	0
Fire clay	4	0	457	0
Shale, gray	8	4	465	4
Sandstone	16	0	481	4

Thickness.	Ft.	In.	Total.
Shale, soft	0	5	199 11
Shale, dark	0	6	200 5
Shale, sandy	2	0	202 5
Clay, sandy	11	4	213 9
Shale, dark, hard	18	3	232 0
Shale, gray	0	7	232 7
Coal, Welch	2	0	234 7
Clay, dark, soft	22	6	257 1
Sandstone, hard, Upper Raleigh	31	11	289 0
Shale, dark	31	11	289 0

Gauley Coal Land Co. Coal Test Boring No. 29—No. 47
on Map II.

Nicholas County, Kentucky District; 2.2 miles east-southeast of
Lolvasy; elevation, 2740' B.

Thickness.	Ft.	In.	Total.
Surface	27	6	27 6
Shale, dark	18	4	45 10
Shale, soft, gray	12	9	58 7
Shale, dark, sandy	8	8	67 3
Shale, soft, gray	16	1	83 4
Shale, gray, sandy	3	8	87 0
Shale, gray, sandy	3	8	87 0
Sandstone	1	1	88 1
Shale, sandy	4	0	92 1
Shale	0	2	92 3
Coal, bony, Castle	0	1	92 4
Fire clay, shaly	1	2	93 6
Shale, gray, sandy	9	2	102 8
Sandstone and shale	8	3	110 11
Shale, soft, gray	2	0	112 11
Shale, dark	5	0	117 11
Shale, black	0	4	118 3
Slate and coal mixed	0	6	118 9
Shale, dark, sandy	2	11	121 8
Sandstone, hard	28	6	150 2
Shale, dark, sandy	6	4	156 6
Sandstone	4	9	161 3
Shale, dark, sandy	16	6	177 9
Shale, dark, sandy	4	0	181 9
Shale, light, sandy	36	3	218 0
Shale, dark, sandy	0	7	218 7
Slate, black	3	2	221 9
Coal, Sewell (2518' B.)	3	2	221 9
Fire clay, sandy	5	3	227 0

Pottsville Series—New River Group (227'+)

Gauley Coal Land Co. Coal Test Boring No. 17—No. 48
on Map II.

Nicholas County, Kentucky District; 1.5 miles east of Hominy
Mill and 0.45 mile southeast of Grassy Falls; completed, Aug. 17, 1917;
elevation, 2420' B.

Pottsville Series—New River Group (142'+)

Thickness.	Ft.	In.	Total.
Surface	20	0	20 0

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, soft	0	5	199	11
Shale, dark	0	6	200	5
Clay, sandy	2	0	202	5
Shale, dark, hard	11	4	213	9
Slate, gray	18	3	232	0
Coal, Welch	0	7	232	7
Clay, dark, soft	2	0	234	7
Sandstone, hard, Upper Raleigh	22	6	257	1
Shale, dark	31	11	289	0

**Gauley Coal Land Co. Coal Test Boring No. 29—No. 47
on Map II.**

Nicholas County, Kentucky District; 2.2 miles east-southeast of Leivasy; elevation, 2740' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—New River Group (227'+)				
Surface	27	6	27	6
Shale, dark	18	4	45	10
Shale, soft, gray	12	9	58	7
Shale, dark, sandy	8	8	67	3
Shale, soft, gray	16	1	83	4
Shale, gray, sandy	3	8	87	0
Sandstone	1	1	88	1
Shale, sandy	4	0	92	1
Slate	0	2	92	3
Coal, bony, Castle	0	1	92	4
Fire clay, shaly	1	2	93	6
Shale, gray, sandy	9	2	102	8
Sandstone and shale	8	3	110	11
Shale, soft, gray	2	0	112	11
Shale, dark	5	0	117	11
Slate, black	0	4	118	3
Slate and coal mixed	0	6	118	9
Shale, dark, sandy	2	11	121	8
Sandstone, hard	28	6	150	2
Shale, dark, sandy	6	4	156	6
Sandstone	4	9	161	3
Shale, dark, sandy	16	6	177	9
Shale, light, sandy	4	0	181	9
Shale, dark, sandy	36	3	218	0
Slate, black	0	7	218	7
Coal, Sewell (2518' B.)	3	2	221	9
Fire clay, sandy	5	3	227	0

**Gauley Coal Land Co. Coal Test Boring No. 17—No. 48
on Map II.**

Nicholas County, Kentucky District; 1.5 miles east of Hominy Mill and 0.45 mile southeast of Grassy Falls; completed, Aug. 17, 1917; elevation, 2420' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—New River Group (142'+)				
Surface	20	0	20	0

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Clay and shale	5	0	25	0
Sandstone and gray shale streaks	12	0	37	0
Sandstone, hard	39	2	76	2
Shale, gray	1	5	77	7
Sandstone, gray, mottled	0	4	77	11
Shale, gray	1	0	78	11
Sandstone, hard	15	1	94	0
Shale, gray	1	0	95	0
Sandstone, hard	9	4	104	4
Shale, gray	13	6	117	10
Sandstone, light	7	1	124	11
Shale, gray	9	1	134	0
Slate, black	0	3	134	3
Coal	1'	1½"	Sewell (2283' B.)	
Slate, gray	0	2½"		
Coal	1	4½"		
"Mother coal"	0	0½"		
Coal	0	5		
Fire clay, dark	1	5		
Fire clay, light	2	8	138	10
			141	6

**Gauley Coal Land Co. Coal Test Boring No. 18—No. 49
on Map II.**

Nicholas County, Kentucky District; 1.3 miles northeast of Grassy Creek School, and 1.9 miles northwest of Eureka School; completed, Aug. 13, 1917; elevation, 2510' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—New River Group (151'+)				
Surface	14	0	14	0
Shale, gray	20	0	34	0
Sandstone	1	0	35	0
Shale, gray	4	6	39	6
Fire clay, shaly	4	3	43	9
Slate	0	11	44	8
Coal, bony	0	4	45	0
Fire clay, shaly	2	0	47	0
Slate, black	0	7	47	7
Fire clay, shaly	6	9	54	4
Coal, bony, Castle	0	6	54	10
Clay, shaly	4	0	58	10
Shale, gray, sandy	6	0	64	10
Shale, gray, soft	17	8	82	6
Shale, dark, sandy	9	2	91	8
Shale, light, sandy	19	6	111	2
Shale, dark	20	3	131	5
Sandstone	2	8	134	1
Shale, dark, sandy	10	2	144	3
Coal	1'	6"	Sewell (2362' B.)	
Fire clay binder	0	4		
Coal	0	4½"		
"Mother coal"	0	1		
Coal	1	8½"		
Fire clay	2	9		
			151	0

**Gauley Coal Land Co. Coal Test Boring No. 6—No. 50
on Map II.**

Nicholas County, Wilderness District; on Hominy Creek, 1.8 miles
south from Hominy Falls; completed, May 26, 1916; elevation, 2375' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (384'+)				
Surface	7	0	7	0
Shale, dark	4	3	11	3
Sandstone	14	3	25	6
Shale, black	4	0	29	6
Coal, Welch	1	4	30	10
Fire clay, impure	0	8	31	6
Sandstone	14	0	46	6
Shale, gray	17	0	63	6
Shale, dark	39	0	102	6
Slate, black	5	0	107	6
Fire clay, impure	4	9	112	3
Shale, dark	3	0	115	3
Sandstone	1	6	116	9
Slate, gray	1	5	118	2
Coal, Little Raleigh	0	8	118	10
Fire clay, impure	13	0	131	10
Slate, gray	25	6	157	4
Slate, black	16	4	173	8
Slate, gray	10	4	184	0
Slate, black	1	0	185	0
Coal, bony	0	2	185	2
Sulphur	0	1	185	3
Coal	0	1	185	7
Slate, dark	61	9	247	4
Coal, bony	0	1	247	5
Shale, dark	0	8	248	1
Sandstone	16	11	265	0
Slate, black	1	6	266	6
Coal, Fire Creek?	1	11	268	5
Slate parting	0	3	268	8
Coal, bony	0	7	269	3
Fire clay	3	7	272	10
Slate, dark	10	3	283	1
Fire clay, impure	1	0	287	1
Shale, gray	8	0	295	1
Sandstone	8	4	303	5
Shale, dark	1	6	304	11
Fire clay	6	2	311	1
Slate, black	1	5	312	6
Sandstone	0	6	313	0
Coal	0	3	313	3
Slate, black	0	3	313	6
Fire clay	1	8	315	2
Shale, dark	30	3	345	5
Fire clay	17	7	363	0
Sandstone	21	0	384	0

Gauley Coal Land Co. Coal Test Boring No. 7—No. 51 on Map II.

Nicholas County, Wilderness District; on Hominy Creek near the mouth of Price Fork; completed, June 23, 1916; elevation, 2435' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (385'+)				
Surface	27	0	27	0
Shale, dark	1	0	28	0
Sandstone	25	4	53	4
Shale, dark	72	1	125	5
Slate, black	2	0	127	5
Fire clay	2	0	129	5
Shale, sandy, gray	25	0	154	5
Sandstone, crystallized, very hard	34	6	188	11
Shale, dark	5	2	194	1
Sandstone, hard	6	11	201	0
Coal, Fire Creek?	0	5	201	5
Fire clay	3	0	204	5
Shale, gray	10	0	214	5
Slate, dark	3	0	217	5
Coal	0	7	218	0
Slate	0	9	218	9
Coal	0	4	219	1
Fire clay	8	6	227	7
Sandstone	4	6	232	1
Shale, gray	3	5	235	6
Slate, black	1	8	237	2
Shale, dark	10	6	247	8
Fire clay	4	0	251	8
Shale, gray	4	9	256	5
Sandstone	25	1	281	6
Coal, No. 8 Pocahontas?	0	4	281	10
Fire clay	1	0	282	10
Sandstone	5	6	288	4
Shale, dark	1	6	289	10
Sandstone	1	0	290	10
Slate, black	2	10	293	8
Coal	0	4	294	0
Fire clay	3	0	297	0
Shale, gray	2	0	299	0
Slate, gray	2	6	301	6
Coal, bony	1	4	302	10
Fire clay	2	10	305	8
Coal	0	4	306	0
Fire clay	1	8	307	8
Sandstone	6	8	314	4
Slate, gray	4	2	318	6
Coal	0	4	318	10
Sulphur band	0	1	318	11
Coal	0	7	319	6
Fire clay	4	3	323	9
Shale, gray	10	0	333	9
Sandstone	3	10	337	7
Slate, dark	0	8	338	3
Coal	0	2	338	5
Slate	0	1	338	6

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Coal, bony	0	5	338	11
Fire clay	4	7	343	6
Shale, dark	2	8	346	2
Sandstone (6' 9" marked with coal spar).....	30	4	376	6
Shale, gray	1	11	378	5
Sandstone	0	2	378	7
Shale, light	6	9	385	4

**Gauley Coal Land Co. Coal Test Boring No. 5—No. 52
on Map II.**

Nicholas County, Wilderness District; in Anglin's Creek, 1.1 miles
north-northeast from Sugargrove School; elevation, 2090' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (137'+)				
Surface	12	0	12	0
Shale, light	19	10	31	10
Shale, dark	1	0	32	10
Fire clay	4	0	36	10
Shale, light	17	10	54	8
Slate, dark	0	8	55	4
Coal, Fire Creek?	0	9	56	1
Fire clay	2	0	58	1
Shale, light	3	0	61	1
Sandstone	5	0	66	1
Shale, light	2	0	68	1
Sandstone	0	9	68	10
Coal, Little Fire Creek?	0	3	69	1
Shale, dark	11	0	80	1
Sandstone	27	5	107	6
Sandstone	12	0	119	6
Slate, soft	3	2	122	8
Fire clay	15	0	137	8
Mauch Chunk Series—Bluestone Group (37'+)				
Shale, red	11	0	148	8
Sandstone	14	0	162	8
Shale, red	2	6	165	2
Shale, light	2	0	167	2
Shale, red	3	0	170	2
Shale, light	1	0	171	2
Shale, red	1	0	172	2
Shale, light	1	0	173	2
Shale, red	1	10	175	0

**DETAILED COAL TEST RECORDS,
FAYETTE COUNTY.**

There have been drilled some 50 core tests for coal in eastern Fayette County, the complete records of most of which were not available for use in the State Survey's report on that county. The complete records of 39 of these tests are now available for publication. Mr. Ray V. Hennen, author of the

Fayette County Report, had access to 28 of these records for study, but only 7 of which were available for publication. (See comments preceeding Nos. 93, 93B, 93C, 93D, 111, 119, and 120.) The elevations and some details of the other 21 tests were given in that report in the table of "Summarized Records of Borings," pages 388B and 388C.

Since, therefore, many of these records are now available in full detail and since they are important in correlating the Greenbrier County coals, it is considered advisable to publish them in this report.

The junior author is responsible for the correlation of all records not previously published by the Survey.

The records of borings Nos. 93, 93B, 93C, and 93D with comments by Ray V. Hennen are reprinted from pages 443-446 of the Fayette County Report.

"The four following records of coal test borings were kindly furnished the Survey by C. E. Krebs of Charleston, W. Va., the correlation of the coal beds being determined by the author (Ray V. Hennen):"

Braekens Creek Coal & Land Company Coal Test Boring No. 2
—No. 93 on Fayette County Map II. (Not Shown on Map II.)

Fayette County, Sewell Mountain District; on waters of Braekens Creek, on hillside just southeast of road fork, 1.4 miles N. 75° W. of Shelton Schoolhouse; 1.25 miles west of S0° 55' and 2.5 miles north of 38° 00'; by Braekens Creek Coal & Land Company, authority, with C. E. Krebs; completed in 1913; elevation, 2357' L.

Pottsville Series (365'+)			Thlekoess.		Total.	
			Ft.	In.	Ft.	In.
Surface			5	6	5	6
Sandstone	6'	7"	51	0	56	6
Shale, sandy	0	8				
Sandstone	27	7				
Sand, shaly	0	11				
Sandstone	15	3	67	4	123	10
Shale, blue, sandy						
Shale, soft						
Slate, coal partings, Sewell "A"						
Sandstone, shale partings						
Shale, sandy						
Fire clay						
Shale, sandy, blue						
Sandstone						
Shale, sandy						
Sandstone						
Shale, sandy						
Sandstone						
Shale, sandy						
Sandstone						
Shale, sandy						
Shale, sandy						

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Sandstone			0	5	157	8
Shale, soft			0	2	157	10
Coal	1'	1"	1	6½	159	4½
Slate and coal part- ing	0	5½				
Fire clay			1	10	161	2½
Shale, sandy			24	10½	186	1
Shale, dark-blue			38	1	224	2
Shale, sandy			0	4	224	6
Sandstone and coal partings	6'	7"	85	8	310	2
Shale, sandy	6	6				
Shale, black, and coal partings	5	11				
Shale, sandy	28	0				
Sandstone	6	4				
Sandstone, coal part- ings	14	8				
Shale, sandy	17	8	0	7	310	9
Coal and slate, Little Raleigh			20	0	330	9
Shale, sandy			3	6	334	3
Shale, blue			0	2	334	5
Slate and coal partings			1	3	335	8
Fire clay			29	4	365	0
Sandstone, Lower Raleigh, to bottom						

Brackens Creek Coal & Land Company Coal Test Boring No. 4—No. 93B on Map II.

Fayette County, Sewell Mountain District; on south branch of Brackens Creek, 1.8 miles S. 35° W. of Russellville; by Brackens Creek Coal & Land Company, authority, with C. E. Krebs; elevation, 2124' L.

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Pottsville Series (247'+)			14	9	14	9
Surface			24	3	39	0
Sandstone, soft, and shale			13	3	52	3
Sandstone, Guyandot			0	6	52	9
Coal, Sewell "B"			1	6	54	3
Fire clay			20	9	75	0
Shale, sandy			17	7	92	7
Shale, blue			1	4	93	11
Fire clay			5	7	99	6
Sandstone			16	4	115	10
Shale, blue			0	8	116	6
Shale, black			0	8	117	2
Shale, blue			1	3	118	5
Shale, black			0	4	118	9
Coal, Sewell "A"			6	0	124	9
Shale, sandy			16	9	141	6
Shale, dark						
Coal	0'	1"	2	5½	143	11½
Slate	0	1				
Coal	1	5				
Coal	0	10½				

		Thickness.		Total.	
		Ft.	In.	Ft.	In.
Shale, black		0	7	144	6½
Sandstone, shaly, Welch		50	5½	195	0
Coal and slate, Welch		0	8	195	8
Shale, sandy		29	6	225	2
Sandstone, with shale	} Upper Raleigh	22	4	247	6
partings					
Sandstone					

Brackens Creek Coal & Land Company Coal Test Boring No. 1
—No. 93D on Fayette County Map II. (Not Shown on Map II.)

Fayette County, Sewell Mountain District; on hillside 0.28 mile S. 45° W. from Shelton Schoolhouse and 1.75 miles northeast of Clifftop; 0.1 mile west of 80° 55' and 1.93 miles north of 38° 00'; by Brackens Creek Coal & Land Company, authority, with C. E. Krebs; completed, Dec. 24, 1912; elevation, 2431' L.

		Thickness.		Total.	
		Ft.	In.	Ft.	In.
Pottsville Series (300'+)					
Surface		12	11	12	11
Shale		14	6	27	5
Shale, sandy		36	0	63	5
Sandstone		10	6	73	11
Shale, dark		23	0	96	11
Sandstone		14	6	111	5
Shale, sandy		5	6	116	11
Shale, dark		35	0	151	11
Coal, bone, Sewell "A"		0	3	152	2
Clay		5	0	157	2
Sandstone, Lower Guyandot		24	5	181	7
Coal, bone	} Sewell	1	1	182	8
Coal					
Fire clay		2	6	185	2
Sandstone		12	0	197	2
Shale		32	10	230	0
Sandstone		20	0	250	0
Shale, sandy		10	0	260	0
Sand and shale		26	1	286	1
Bone		0	1	286	2
Sandstone and coal partings.....		4	6	290	8
Shale, sandy		9	4	300	0

Brackens Creek Coal & Land Company Coal Test Boring No. 3
—No. 93C on Fayette County Map II. (Not Shown on Map II.)

Fayette County, Sewell Mountain District; on a branch of Brackens Creek, southeast of Hogpen Ridge, 0.9 mile N. 15° W. of Shelton Schoolhouse, and 2.6 miles N. 15° E. of Clifftop; 0.35 mile west of 80° 55' and 2.82 miles north of 38° 00'; by Brackens Creek Coal & Land Company, authority, with C. E. Krebs; completed, Jan. 9, 1913; elevation, 2345' L.

		Thickness.		Total.	
		Ft.	In.	Ft.	In.
Pottsville Series (229'+)					
Surface		12	4	12	4
Sandstone, Harvey Conglomerate		61	2	73	6

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, dark	6	2	182	0
Fire clay	4	4	186	4
Shale, gray, sandy, hard	15	2	201	6
Sandrock, hard	31	6	233	0
Shale, dark, sandy	11	6	244	6
Slate, black	4	2	248	8
Fire clay, soft	7	4	256	0
Mauch Chunk Series (15'+)				
Shale and fire clay, hard	9	0	265	0
Shale, red	6	0	271	0

DETAILED COAL TEST RECORDS, IRISH CORNER DISTRICT.

In Irish Corner District three test holes have been drilled for coal. Another was drilled just across the county line in Monroe County. These holes all start in the Pocono Series of lower Mississippian age. As previously stated in Chapter VII, there appears to be little chance of finding coal of commercial thickness and purity in this series in Greenbrier County.

The following record with comments by Reger is reprinted from the Mercer, Monroe, and Summers Report, pages 671-672:

"... The Merrimac Coal horizon appears to belong about the level of the shale at 118-121 feet, the elevation of which is 1659' B., as compared to 1650' B., at Coal Prospect No. 797 on Map IV (No. 503 on Map II of present report), located slightly to the northwest. The black shale at 170-171 feet is plainly too low for the Merrimac Coal since the dip is northwestward, and it evidently belongs at or near the Langhorne horizon:"

Hunter Moore Coal Test Boring No. 1—No. 16 on Map II.

Irish Corner District; along road east of Second Creek and 0.5 mile north of Hokes Mill; authority, Homer Hoke et al.; elevation, 1780' B.

	Thickness.	Total.	
Pocono Series (329'+)	Feet.	Feet.	
Sand and clay	8	8	
Sandstone, Squaw	107	115	
Shale	2	117	
Sandstone	1	118	
Shale, Merrimac Coal horizon?.....	3	121	
Sandstone and shale, laminated	13	134	
Sandstone	10'	} Lindsie Sandstone.. 36	170
Shale	15		
Sandstone, light-gray	4		
Sandstone	7		
Shale, black, "nearly coal," Langhorne Coal?.....	1		171

		Thickness. Feet.	Total. Feet.
Sandstone, gray, fine-grained	7'		
Sandstone, with layers of shale	24		
Sandstone and shale	86		
Sandstone, dark, close-grained	7	Broad Ford	158
Sandstone, with quartz partings	21		
Sandstone, hadly broken, to bottom	13		
			329

The following record with comments by Reger is reprinted from the Mercer, Monroe, and Summers Report, page 671:

" . . . the coal partings found at 160-161 feet and having an elevation of 1734' B., apparently belong about the proper level for the Merrimac Coal and indicate its unreliable nature in this vicinity . . ."

Mary E. Morris Heirs Coal Test Boring No. 2— No. 17 on Map II.

Irish Corner District; near road fork east of Second Creek and 0.3 mile northeast of Hokes Mill; authority Homer Hoke et al.; elevation, 1895' B.

	Thickness. Feet.	Total. Feet.
Pocono Series (223'+)		
Sandstone and clay	19	19
Sandstone (water at 60')..126'		
Sandstone shale, "conglomerated"	2	
Sandstone	13	
Squaw Sand	141	160
Coal partings, Merrimac?.....	1	161
Sandstone	16	177
Sandstone and shale	18	195
Shale	5	200
Shale and sandstone, to bottom	23	223

The following record with comments by Reger is reprinted from the Mercer, Monroe, and Summers Report, pages 670-671:

"On the evidence of surface outcrops the following hole starts below the level of the Merrimac Coal which should have an elevation of 1750 feet or more at this point, and it probably starts nearly 200 feet below the top of the Pocono Series . . ."

A. W. Smith Coal Test Boring No. 3—No. 18 on Map II.

Irish Corner District; on east side of Second Creek just south of Hokes Mill; authority, Homer Hoke et al.; completed, May 11, 1922; elevation, 1705' B.

		Thickness.	Total.
		Feet.	Feet.
Pocono Series (216'+)			
Clay and boulders		6	6
Sandstone, Broad Ford		45	51
Shale, black, Sunbury		49	100
Shale		35	135
Sandstone	52'	81	216
Sandstone and shale	16		
Sandstone, to bottom	13		

The following record is of a boring drilled in Monroe County 0.55 mile south of coal test boring No. 18. The record with comments by Reger is reprinted from the Mereer, Monroe, and Summers Report, page 670:

"... the Merrimac Coal horizon appears to have been penetrated at 63-64 feet, its elevation being 1791' B."

Harry Ellis Coal Test Boring No. 4—No. 19 on Map II.

Monroe County, Second Creek District; on short branch of Second Creek 0.7 mile south of Hokes Mill; authority, Homer Hoke et al.; completed, May 25, 1922; elevation, 1855' B.

		Thickness.	Total.
		Feet.	Feet.
Pocono Series (156'+)			
Shale, blue		16	16
Shale, dark and bluish-gray		34	50
Shale, carbonaceous		13	63
Shale, black, with coal seams, Merrimac Coal?		1	64
Shale, black		1	65
Shale and sandstone		26	91
Sandstone		14	105
Sandstone and shale, to bottom		51	156

DETAILED COAL TEST RECORDS, NICHOLAS COUNTY.

The records of the following borings in Nicholas County are published with the permission of Mr. J. S. McWhorter, Resident Attorney, Gauley Coal Land Company, Rupert, W. Va. The location and barometric surface elevation of most of these borings were given by Reger in the Nicholas County Report published in 1921. With three exceptions, however, the records of these borings were not available to Mr. Reger. For these exceptions see the comments immediately preceding the records of Nos. 25, 38, and 46B.

No elevations for these borings were found in the Gauley Coal Land Company's files, and it has been necessary to use the elevation shown on the topographic map for those bore

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Coal, Sewell (2269' B.)	1	3	261	3
Fire clay, soft	2	9	264	0
Fire clay, hard	2	0	266	0
Shale, sandy	2	2	268	2
Sandstone and shale	11	0	279	2
Sandstone	14	11	294	1
Shale, dark	15	5	309	6
Coal, Welch	0	7	310	1
Fire clay	1	8	311	9
Shale, dark, sandy	2	0	313	9
Sandstone and shale	23	11	337	8
Shale, dark, sandy	34	8	372	4
Shale, light, sandy	4	8	377	0
Sandstone, hard	19	3	396	3
Shale, variegated	1	0	397	3
Shale, light, sandy	7	4	404	7
Sandstone and shale	2	7	407	2
Shale, dark, sandy	14	4	421	6
Shale, soft, dark	1	8	423	2
Shale, light, sandy	7	3	430	5
Shale, dark, sandy	2	7	433	0
Fire clay, hard	0	2	433	2
Shale, gray	0	9	433	11
Shale and clay	12	1	446	0
Shale, dark, sandy	19	9	465	9
Slate, black	0	10	466	7
Slate, black, with coal	0'	10"	Beckley	11
Coal, dirty	0	6		
Fire clay, sandy	2	2	470	1
Shale, dark, sandy	24	7	494	8
Sandstone and shale	5	4	500	0
Sandstone, with shale streaks	8	3	508	3
Sandstone and shale	4	3	512	6
Shale, dark, sandy	9	6	522	0
Slate, black	0	2	522	2
Fire clay and shale	4	0	526	2
Slate, black, Fire Creek Coal horizon (2001' B.)	2	5	528	7
Shale, soft	0	3	528	10
Shale, sandy	7	2	536	0
Sandstone	12	2	548	2
Coal, bony	0	1	548	3
Fire clay	0	10	549	1
Coal, bony, Little Fire Creek	0	3	549	4
Fire clay	0	8	550	0

Gauley Coal Land Co. Coal Test Boring No. 26—No. 26 on Map II.

Nicholas County, Kentucky District; on Taylor Run 1.3 miles south of mouth, 2.9 miles northeast of Lowland and 3.9 miles northwest of Fenwick; elevation, 2125' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—New River Group (230'+)				
Surface	6	0	6	0

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Sandstone	40	8	46	8
Sandstone and shale	20	0	66	8
Shale, sandy	6	2	72	10
Fire clay	2	7	75	5
Shale, sandy	3	0	78	5
Shale, dark	1	0	79	5
Coal, Castle	1	0½	80	5½
Fire clay	1	0½	81	6
Shale, sandy	4	0	85	6
Sandstone	6	0	91	6
Shale, sandy	46	4	137	10
Shale, light, sandy	3	4	141	2
Sandstone	4	0	145	2
Shale, dark, sandy	1	4	146	6
Sandstone	0	8	147	2
Shale, dark, sandy	4	5	151	7
Sandstone and shale	24	6	176	1
Sandstone, with slate streaks	7	1	183	2
Sandstone	9	0	192	2
Coal, bony	0'	6½"	Sewell (1928' B.) ...	5
Coal	0	5½		
Coal, bony	0	1		
Coal	3	2		
Coal, with knife- edge streaks of slate	0	9		
Fire clay, sandy	2	10	200	0

**Gauley Coal Land Co. Coal Test Boring No. 24—No. 27 on
on Map II.**

Nicholas County, Kentucky District; 2.6 miles west of Feuerick;
elevation, 2600' B.

Pottsville Series—New River Group (476' +)	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Surface	43	6	43	6
Sandstone, hard	31	8	75	2
Slate, black	0	1	75	3
Coal, dirty	0	4	75	7
Shale, dark	0	1	75	8
Shale, gray	9	4	85	0
Shale, dark	1	0	86	0
Shale, gray	3	7	89	7
Shale, dark, sandy	10	6	100	1
Shale, gray	3	5	103	6
Slate, black	0	4	103	10
Coal, dirty, Lower laeger	1	1	104	11
Slate, black	0	3	105	2
Shale, gray, sandy	6	2	111	4
Shale, dark, sandy	10	6	121	10
Sandstone, hard, Lower laeger	42	6	164	4
Shale, dark	4	6	168	10
Coal, dirty	0	2	169	0

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, dark	3	0	172	0
Sandstone, hard, Harvey	7	0	179	0
Shale, dark, gray	28	6	207	6
Slate, black, with coal spars, Castle?	2	9	210	3
Clay, shaly	0	9	211	0
Shale, gray	12	6	223	6
Shale, dark, sandy	21	3	244	9
Sandstone, hard, Guyandot?	32	3	277	0
Shale, dark, sandy	29	6	306	6
Coal, dirty, Sewell? (2293' B.)	0	1	306	7
Shale, dark, sandy, with sand streaks	6	2	312	9
Sandstone, hard, with shalo streaks	47	3	360	0
Sandstone, with coal spars	20	6	380	6
Coal	0	1	380	7
Sandstone, with coal streaks	2	11	383	6
Coal	0	2	383	8
Sandstone	0	1½	383	9½
Coal	0	1½	383	11
Sandstone, with coal spars	2	1	386	0
Coal	0	1	386	1
Sandstone, with coal spars	1	8½	387	9½
Coal	0	0½	387	10
Sandstone	0	0½	387	10½
Coal	0	3½	388	2
Sandstone, hard, with coal spars	5	0	393	2
Coal	0	3	393	5
Sandstone,	0	0½	393	5½
Coal	0	1½	393	7
Sandstone	0	5½	394	0½
Coal and sandstone mixed	0	3½	394	4
Sandstone	0	4½	394	8½
Coal	0	1	394	9½
Sandstone	0	10	395	7½
Coal, dirty	0	11½	396	7
Sandstone, with coal spars	0	1½	396	8½
Coal, clean	0	2	396	10½
Sandstone	0	3	397	1½
Coal	0	1	397	2½
Sandstone	0	2	397	4½
Coal	0	2½	397	7
Sandstone, with coal spars	0	6	398	1
Coal	0	1½	398	2½
Sandstone	0	0½	398	3
Coal	0	0½	398	3½
Sandstone, with coal spars.....	0	1½	398	5
Coal, dirty	0	3	398	8
Sandstone, with coal spars	0	6	399	2
Fire clay, sandy	4	1	403	3
Shale, light, sandy	11	7	414	10
Shale, dark, sandy	10	5	425	3
Sandstone	0	8	425	11
Shale, dark, sandy	0	3	426	2
Sandstone	3	5	429	7
Shale, dark, sandy	4	0	433	7
Sandstone	36	8	470	3
Shale, dark	5	9	476	0

Gauley Coal Land Co. Coal Test Boring No. 21—No. 27A on Map II.

Nicholas County, Kentucky District; 0.15 mile south of Lowland;
elevation, 2190' \pm .

				Thickness.		Total.	
Pottsville Series—New River Group (137' +)				Ft.	In.	Ft.	In.
Surface				22	0	22	0
Shale, soft				2	0	24	0
Shale, dark, variegated				33	0	57	0
Shale, light, sandy				8	0	65	0
Slate, soft				0	1½	65	1½
Coal	0'	3"	Sewell "B"....	1	3	66	4½
Slate, dark, with coal streaks	1	0		2	0	68	4½
Shale, dark, sandy				6	5½	74	10
Shale, dark, variegated				1	11	76	9
Sandstone, light				10	1	86	10
Shale, dark, sandy				1	7	88	5
Shale, dark				1	3	89	8
Shale, dark, soft				2	0	91	8
Shale, dark				4	7	96	3
Shale, dark, sandy				1	9	98	0
Shale, gray, sandy				1	3	99	3
Sandstone				0	11	100	2
Shale, dark, sandy				5	10	106	0
Sandstone, hard, with coal spars				8	8	114	8
Sandstone				14	10	129	6
Shale, light and dark				0	0½	129	6½
Slate, black							
Coal, bony	0'	1½"	Sewell (2057' \pm)	3	5½	132	11½
Coal, clean	1	4					
Coal, knife-edge layers of slate	1	11½"		4	0½	137	0
Fire clay, light							

The record of boring No. 29 could not be found in the files of the Gauley Coal Land Company.

Gauley Coal Land Co. Coal Test Boring No. 23A—No. 30 on Map II.

Nicholas County, Kentucky District; 1.65 miles southwest of Saxman; elevation, 2580' B.

				Thickness.		Total.	
Pottsville Series—New River Group (177' +)				Ft.	In.	Ft.	In.
Surface				21	6	21	6
Shale, gray				39	2	60	8
Shale, soft, light				3	4	64	0
Shale, soft, gray				9	3	73	3
Shale, gray, sandy				16	0	89	3
Shale, soft, light				3	4	92	7
Shale, light, sandy				2	6	95	1
Sandstone, hard, Guyandot				56	9	151	10

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Coal, Sewell "A".....	0	11	152	9
Fire clay, shaly.....	1	6	154	3
Shale, hard, dark	1	2	155	5
Sandstone, hard, with shale streaks	13	2	168	7
Sandstone and shale, dark	2	1	170	8
Shale, soft, gray	0	7½	171	3½
Slate	0	0½	171	4
Coal, Sewell (2407' B.)	1	10	173	2
Shale, soft, dark.....	0	6½	173	8½
Fire clay, sandy	3	3½	177	0

Gauley Coal Land Co. Coal Test Boring No. 11—No. 31 on Map II.

Nicholas County, Kentucky District; on west bank of Jims Branch, 1.55 miles northwest of Tolbert and 1.4 miles south of Lowland; elevation, 2315' B.

		Thickness.		Total.	
		Ft.	In.	Ft.	In.
Pottsville Series—New River Group (138'+)					
Surface		12	0	12	0
Sandstone, hard		1	5	13	5
Shale, sandy, dark.....		1	0	14	5
Sandstone, Guyandot		12	1	26	6
Shale, sandy, dark.....		14	0	40	6
Sandstone		2	4	42	10
Shale, sandy, gray.....		8	2	51	0
Slate		1	6	52	6
Coal, bony	0'	5"	Sewell "B"....	4	1
Slate, with coal streaks	0	6			
Coal, bony	0	2			
Slate	0	1			
Slate and coal, bony	2	1			
Coal	0	10	Lower Guyandot	43	2
Slate and clay		0			
Clay, soft		2			
Shale, sandy, gray.....		5			
Sandstone and shale 11' 1".....		32			
Sandstone, hard.....	32	1		107	6
Shale, gray, with hard streaks.....		23	4	130	10
Coal, Sewell (2181' B.).....		3	6½	134	4½
Fire clay		3	7½	138	0

Gauley Land Co. Coal Test Boring No. 2—No. 32 on Map II.

Nicholas County, Kentucky District; on Jims Branch of Panther Creek, 2.4 miles northeast from Tolbert; elevation, 2325' B.

		Thickness.		Total.	
		Ft.	In.	Ft.	In.
Pottsville Series (285'+)					
Surface		15	0	15	0
Shale, dark		37	4	52	4
Sandstone		2	6	54	0

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Slate, dark	4	2	60	0
Bone	0	5	60	5
Coal and slate.....	2	5	62	10
Coal, Sewell "B"?	1	4	64	2
Fire clay	0	10	65	0
Shale, light	9	6	74	6
Sandstone, hard, Lower Guyandot?	29	10	104	4
Shale, dark	6	0	110	4
Coal	1	4	111	8
Slate, dark	0	3	111	11
Coal	0	3	112	2
Slate, dark	0	4	112	6
Coal	0	1	112	7
Fire clay	3	4	115	11
Shale, dark	11	6	127	5
Slate, dark	0	4	127	9
Coal, Sewell (2194' B.).....	3	8	131	5
Fire clay	3	4	134	9
Sandstone	8	0	142	9
Shale, dark	8	2	150	11
Coal, Welch?	0	1	151	0
Fire clay	2	0	153	0
Shale, light	10	0	163	0
Shale, dark	14	0	177	0
Slate, dark	5	2	182	2
Coal, dirty, Welch?.....	0	8	182	10
Fire clay	0	10	183	8
Sandstone, Upper Raleigh.....	11	8	195	4
Shale, dark	51	5	246	9
Slate, dark	1	3	248	0
Fire clay	7	0	255	0
Sandstone, Lower Raleigh?	30	0	285	0

Gauley Coal Land Co. Coal Test Boring No. 25—No. 33 on Map II of Nicholas County Report.

Nicholas County, Kentucky District; on Little Laurel Creek, 2.1 miles northwest of Lowland and 2.15 miles north of Nettle; elevation, 2045' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (150'+)				
Surface	20	6	20	6
Sandstone	23	6	44	0
Shale, soft, dark.....	10	8	54	8
Slate and coal, Castle?.....	0	6	55	2
Shale, dark	3	8	58	10
Slate, with coal spars.....	1	4	60	2
Shale, sandy	10	0	70	2
Slate, soft, with coal spars.....	1	10½	72	0½
Slate and coal.....	0	1½	72	2
Slate, soft	0	2	72	4
Coal streaks and "mother coal".....	0	8	73	0
Fire clay, soft	5	2	78	2

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, gray	4	0	82	2
Sandstone	6	0	88	2
Shale, sandy	10	6	98	8
Slate	27	2	125	10
Shale, light, sandy.....	8	2	134	0
Shale, sandy	9	3	143	3
Shale, soft	0	1	143	4
Coal, clean, with thin streaks of "mother coal," Sewell (1898').....	3	4	146	8
Fire clay, sandy	3	4	150	0

Gauley Coal Land Co. Coal Test Boring No. 4—No. 37 on Map II.

Nicholas County, Kentucky District; on Hominy Creek, 0.9 mile northeast from Blacks Chapel School; elevation, 1660' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (236'+)				
Surface	13	0	13	0
Shale, dark	10	0	23	0
Slate and coal.....	0	6	23	6
Sandstone and shale.....	11	7	35	1
Shale, dark	12	3	47	4
Coal	0	6	47	10
Fire clay	3	8	51	6
Sandstone	3	3	54	9
Sandstone and shale.....	13	4	68	1
Sandstone	4	0	72	1
Shale, dark	3	0	75	1
Sandstone	27	6	102	7
Slate, dark	6	1	108	8
Bone	0	4	109	0
Coal, dirty	0	8	109	8
Fire clay	1	6	111	2
Shale, dark	8	0	119	2
Coal, dirty	0	3	119	5
Shale, dark	11	3	130	8
Shale and sandstone	24	5	155	1
Fire clay, Sewell Coal horizon?.....	6	0	161	1
Shale, light	9	6	170	7
Sandstone, hard	30	5	201	0
Shale, dark	35	0	236	0

The record of coal test No. 38 was published in connection with the Fury Knob Section by Reger in the Nicholas County Report, pages 170-171. The Sewell Coal in that publication was reported as 4 feet 6 inches. However, it will be noted from this record that the bottom 2' 2" is given here as slate, streaked with coal, leaving only 2' 4" of clean coal at the top. The record of this hole as given below differs slightly in other particulars from that previously published:

Gauley Coal Land Co. Coal Test Boring No. 1—No. 38 on Map II.

Nicholas County, Kentucky District; at Deepwell P. O.; drilled by E. F. Saxman; authority, Gauley Coal Land Co.; elevation, 1800' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (640'+)				
Surface	23	0	23	0
Shale and soapstone.....	21	9	44	9
Sandstone	4	6	49	3
Shale, light	1	8	50	11
Sandstone	3	4	54	3
Shale, light	17	0	71	3
Slate, black	2	9	74	0
Coal	0	9	74	9
Fire clay	8	0	82	9
Shale, light	3	11	86	8
Sandstone	0	6	87	2
Shale, dark	0	8	87	10
Sandstone	2	7	90	5
Shale, dark	0	9	91	2
Sandstone	2	6	93	8
Shale, dark	9	0	102	8
Sandstone	2	5	105	1
Shale, dark	1	8	106	9
Shale, dark	0	2	106	11
Coal	2'	4"	Sewell (1689' L.)	4
Slate, streaked with coal	2	2		
Fire clay	2	10	111	5
Shale, dark	29	2	114	3
Slate, streaked with coal.....	0	8	143	5
Shale, dark	3	8	144	1
Shale, light	9	0	147	9
Sandstone, hard	12	2	156	9
Shale, dark	13	6	168	11
Coal, dirty	1	2	182	5
Fire clay	2	2	183	7
Sandstone	9	7	185	9
Shale, dark	140	7	195	4
Sandstone	25	8	335	11
Sandstone, conglomerate	3	7	361	7
Fire clay	4	0	365	2
Shale, dark	6	6	369	2
Fire clay	8	0	375	8
Sandstone	54	0	383	8
Shale, dark	1	6	437	8
Sandstone	2	6	439	2
Shale, dark	2	0	441	8
Sandstone	14	10	443	8
Fire clay	2	10	458	6
Sandstone	20	6	461	4
Shale, light, sandy.....	1	3	481	10
Sandstone	1	0	483	1
			484	4?

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, light, sandy.....	16	10	501	2
Sandstone	3	8	504	10
Shale, light, sandy.....	7	8	512	6
Sandstone	60	3	572	9
Shale, light, sandy	26	6	599	27
Sandstone	30	4	629	6
Shale, dark	10	8	640	2

**Gauley Coal Land Co. Coal Test Boring No. 13—No. 39
on Map II.**

Nicholas County, Kentucky District; 0.75 mile north of Ophella and 1.75 miles west of Nettle; elevation, 2162' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—New River Group (400'+)				
Surface	23	0	23	0
Shale, gray	21	0	44	0
Shale, light	0	6	44	6
Shale and coal, Castle	0	4	44	10
Shale and clay.....	1	0	45	10
Sandstone, Guyandot	30	2	76	0
Shale, dark	0	8	76	8
Coal, Sewell "B".....	0	3	76	11
Shaly clay	2	6	79	5
Shale, dark	37	1	116	6
Sandstone	2	0	118	6
Shale, light	28	1	146	7
Slate, black	2	5	149	0
Coal, Sewell "A"?	1	1	150	1
Shale, dark	6	8	156	9
Coal	1'	4½"	Sewell (2003' L.)....	
Coal, bony	0	2½		
Slate	0	0½		
Cannelized slate.....	0	5½		
Sandy clay	6	2	165	0
Sandstone	3	8	168	8
Shale, dark	20	4	189	0
Coal, Welch	0	4	189	4
Fire clay, dark.....	1	8	191	0
Sandstone, hard, Upper Raleigh.....	21	0	212	0
Shale, dark	31	0	243	0
Fire clay, soft, Little Raleigh Coal horizon?	4	0	247	0
Shale, light	32	0	279	0
Sandstone, light, Lower Raleigh.....	20	0	299	0
Shale, dark	12	0	311	0
Sandstone, hard, dark.....	4	0	315	0
Sandstone, hard, light.....	9	6	324	6
Slate, gray	6	8	331	2
Coal, bony, Beckley?.....	0	5½	331	7½
Shale, gray	68	4½	400	0

Gauley Coal Land Co. Coal Test Boring No. 12—No. 40 on Map II.

Nicholas County, Kentucky District; 0.55 mile southeast of Ophelia and 1.2 miles northwest of Tolbert; elevation, 2715' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (615'+)				
Surface	11	0	11	0
Shale, dark	7	0	18	0
Shale, sandy, gray.....	28	0	46	0
Shale, sandy, dark.....	27	6	73	6
Sandstone, Lower Dotson	19	6	93	0
Shale, sandy, dark	18	6	111	6
Slate, black	0	3	111	9
Coal, Lower Douglas	1	0½	112	9½
Fire clay	0	2½	113	0
Shale, dark, sandy.....	11	6	124	6
Sandstone, hard, gray, Upper Nuttall	50	0	174	6
Shale, dark	0	6	175	0
Sandstone	0	8	175	8
Slate, black	4	0	179	8
Sandstone, hard, gray.....	5	4	183	0
Coal	0	10	183	10
Clay, sandy	2	2	186	0
Shale, gray, with hard sand streaks.....	6	0	192	0
Shale, dark, with hard sand streaks.....	21	0	213	0
Shale, dark	26	6	239	6
Slate	0	7	240	1
Slate and bony coal.....	0	6	240	7
Shale, gray, with sand streaks	11	0	251	7
Shale, dark	6	3	257	10
Sandstone	0	7	258	5
Slate, black	5	5	263	10
Sandstone	0	8	264	6
Sandstone and shale	1	6	266	0
Sandstone, gray	6	0	272	0
Shale, dark-gray, mixed.....	8	4	280	4
Shale, dark	22	0	302	4
Coal, Hughes Ferry	0	7	302	11
Shale, gray, sandy.....	1	11	304	10
Fire clay	2	2	307	0
Shale, gray, sandy.....	10	0	317	0
Sandstone, hard, gray, with coal spars, Middle laeger	30	4	347	4
Slate	0	3	347	7
Clay, sandy	1	6	349	1
Shale, dark, sandy.....	11	5	360	6
Sandstone, gray	31	0	391	6
Shale, gray, sandy.....	11	6	403	0
Shale, dark	4	6	407	6
Coal, Lower laeger	0	10½	408	4½
Fire clay	0	7½	409	0
Sandstone, hard, Lower laeger	15	6	424	6
Shale, hard, dark.....	28	0	452	6
Shale, gray, sandy.....	11	0	463	6
Shale and sandstone, dark mixed.....	37	4	500	10

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Shale, dark			7	6	508	4
Slate			0	11	509	3
Coal, clean, Castle			1	11½	511	2½
Slate, soft			0	1	511	3½
Clay, sandy			0	2½	511	6
Slate, soft			0	1	511	7
Shale, sandy, with clay streaks.....			1	10	513	5
Sandstone, hard, Guyandot			32	3	545	8
Sandstone, with coal spars.....			3	0	548	8
Shale, dark			0	5	549	1
Sandstone, hard			4	5	553	6
Shale, hard, sandy.....			12	6	566	0
Shale, dark, with sand streaks.....			25	0	591	0
Sandstone, hard, Lower Guyandot			12	10	603	10
Coal, clean	0'	1½"	} Sewell (2103' B.).....			
Coal, dirty	0	1½				
Slate	0	0¾				
Coal, clean	0	3¼				
Sulphur ball,						
tapered	0	0½				
Coal, good, clean....	1	9				
Coal, dirty	0	1½				
Slate, black	1	5				
Coal, good, clean....	1	2½				
Coal, knife-edge						
slate'	0	2½				
Coal, good	0	7				
"Mother coal"	0	1				
Coal, good	0	7				
Coal, bony, hard.....	0	4½				
Coal, good	0	9¾				
Fire clay, sandy.....			4	0	615	7

The following quotation is part of the core record:

"The above record is correct except item 75, [third line from bottom of section (Coal, bony, hard....0' 4½")] which I am satisfied fell into the hole while the drill was withdrawn on account of core barrel being full. I watched the drill cutting the core and when necessary to pull out, it was still in good coal. The drill rod was marked and when put back into the hole would not go down to the bottom by four inches, which was caused, in my opinion, by the piece of bony coal falling in from the side of the hole nearer the top. After the drill got down as deep as it was before being pulled out, it continued to cut soft coal for 9¾ inches before getting into hard stuff, and when drill was pulled out, the coal measured 9¾". Core was in good shape. Coal looks good."

Gauley Coal Land Co. Coal Test Boring No. 19—No. 41 on Map II.

Nicholas County, Kentucky District; 1.05 miles northwest of Mayflower School; elevation, 2210' B.

		Thickness.		Total.	
Pottsville Series—New River Group (311'+)		Ft.	In.	Ft.	In.
Surface		16	0	16	0
Shale, soft		7	0	23	0
Shale, gray, sandy		16	6	39	6
Shale, dark		51	8	91	2
Coal	1' 1"				
Coal, bony	0 3				
Clay, shaly	1 11½				
Coal, bony	0 1				
Slate	0 1½				
Sandstone	0 3				
Coal, bony	0 1½				
Shale, dark, sandy.....		0	4½	95	5
Fire clay		6	1	101	6
Clay, gray, shaly.....		1	5	102	11
Sandstone		1	3	104	2
Clay		0	1	104	3
Clay and coal mixed		0	2	104	5
Sandstone		0	6	104	11
Shale, dark, lime.....		2	6	107	5
Shale, dark		8	4	115	9
Shale, gray		13	0	128	9
Shale, gray, sandy.....		20	9	149	6
Shale, dark, sand streaks.....		7	4	156	10
Sandstone, mottled		15	2	172	0
Shale, dark, hard sand streaks		33	4	205	4
Coal, bony	0' 1"				
Coal	0 3				
Coal, bony	0 0½				
Coal	1 1½				
Clay, soft		0	4½	207	2½
Clay, sandy		0	3½	207	6
Shale, hard, dark, sandy.....		6	10	214	4
Sandstone, hard, Lower laeger		4	8	219	0
Shale, hard, dark.....		23	0	242	0
Coal		0	3	242	3
Clay, sandy		1	1	243	4
Shale, gray, sandy.....		3	4	246	8
Sandstone		1	3	247	11
Shale, dark, sandy.....		1	6	249	5
Sandstone, hard, Harvey.....		23	1	272	6
Shale, hard, dark, sandy streaks.....		25	2	297	8

			Thickness.		Total.	
			Ft.	In.	Ft.	In.
Coal	0'	3"				
"Mother coal"	0	1				
Coal	0	3 $\frac{3}{4}$				
"Mother coal"	0	0 $\frac{1}{4}$				
Coal	0	8				
Variegated sandstone, with white and dark streaks	0	5 $\frac{1}{2}$				
Slate, black	0	6 $\frac{1}{4}$				
Slate, black, with coal streaks	1	1	Castle (1908' B.)..	4	0 $\frac{1}{4}$	301
Coal	0	7 $\frac{1}{2}$				
Clay, light, sandy.....						
Shale, light, sandy.....						
Shale, dark, sandy.....						
						8 $\frac{1}{2}$?
			1	8	303	4 $\frac{1}{2}$
			3	4	306	8 $\frac{1}{2}$
			4	3 $\frac{1}{2}$	311	0

**Gauley Coal Land Co. Coal Test Boring No. 3—No. 43
on Map II.**

Nicholas County, Kentucky District; on Deer Creek, 0.7 mile southeast from Trimble School; elevation, 2055' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—New River Group (445'+)				
Surface	20	0	20	0
Sandstone, Upper Nuttall	16	0	36	0
Fire clay	6	0	42	0
Shale, dark	95	0	137	0
Slate, gray	6	0	143	0
Coal, bony, Hughes Ferry.....	0	2	143	2
Slate, soft	3	8	146	10
Fire clay	2	6	149	4
Sandstone	2	4	151	8
Fire clay	1	2	152	10
Sandstone	76	9	229	7
Slate, dark	3	3	232	10
Bone	0	4	233	2
Coal, Lower laeger	1	0	234	2
Slate and coal	1	4	235	6
Fire clay	0	5	235	11
Shale, dark	2	9	238	8
Shale, light	9	7	248	3
Shale, dark	15	3	263	6
Sandstone	18	10	282	4
Sandstone and shale	17	6	299	10
Shale, dark	10	6	310	4
Sandstone	2	0	312	4
Shale, dark	3	0	315	4
Coal, Castle	1	10	317	2
Slate, soft	0	10	318	0
Fire clay	5	0	323	0
Shale, dark	12	3	335	3
Sandstone	4	5	339	8
Shale, light	17	3	356	11
Shale, dark	18	0	374	11

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Fire clay, sandy	1	10	201	6
Sandstone, hard	0	6	202	0
Continued with record of W. E. Deegans				
No. 1 (Above boring deepened, beginning at 202'; started, Jan. 13, 1930; completed, Jan. 15, 1930).				
Sandstone, hard	3	2	205	2
Shale, dark	8	11	214	1
Coal, Fire Creek?	2	5	216	6
Fire clay	3	4	219	10
Slate, black	3	2	223	0
Shale, dark, sandy	10	0	233	0
Shale, light, sandy	50	6	283	6
Sandstone	13	6	297	0
Sandstone, hard, with hard slate streaks.....	5	0	302	0

The following 12 records are of borings drilled northwest of Anjean for the Leekie Smokeless Coal Company, partly on their own property and partly on land leased from the Ganley Coal Land Company. As prospecting is still in progress, permission to publish the actual coal sections was withheld. The beds immediately above the coal seams are included in the measurement indicated by coal bed correlations:

Leekie Smokeless Coal Company Coal Test Boring No. 4— No. 5A on Map II.

Meadow Bluff District; on west side of Brown Creek 1.2 miles northwest of mouth, 2.1 miles west of Anjean; started, April 20, 1931; completed, May 29, 1931; elevation, 3357' L.

				Thickness.		Total.	
Pottsville Series (555'+)				Ft.	In.	Ft.	In.
Surface				18	6	18	6
Shale, dark, sandy				16	6	35	0
Sandstone, broken, hard	30'	0"	Upper Raleigh Sandstone	72	0	107	0
Sandstone and shale	2	0					
Sandstone, broken, hard	14	0					
Shale, dark	2	0					
Sandstone, broken, hard	17	0					
Shale, dark	1	0					
Sandstone, broken, hard..	6	0					
Shale, dark, sandy				20	0	127	0
Shale, dark, and coal, Little Raleigh				14	8	141	8
Fire clay, shaly				7	4	149	0
Shale, blue, sandy				16	0	165	0
Sandstone, broken, hard	20'	0"	Lower Raleigh Sandstone	50	0	215	0
Sandstone, shale streaks	5	0					
Sandstone, broken, hard	25	0					
Shale, blue, sandy				20	0	235	0
Shale, dark-blue, sandy				20	0	255	0
Shale, dark, and coal, Beckley (3085').....				16	8	271	8

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, dark	0	3	271	11
Fire clay	2	0	273	11
Fire clay, sandy	2	1	276	0
Shale, gray, sandy	4	0	280	0
Sandstone, hard	11'	0"	Quinnimont Sandstone 57	0
Shale, blue	2	0		
Sandstone, hard	14	0		
Shale, dark	1	0		
Sandstone, hard	29	0		
Shale, gray, sandy, shale, dark, and coal. (Little Fire Creek?) Fire Creek? (2998').....	22	0	359	0
Shale, soft	5	0	364	0
Shale, sandy, blue	11	0	375	0
Sandstone, hard, Pineville	32	0	407	0
Shale, blue, sandy	7	0	414	0
Shale, dark, and coal, No. 8 Pocahontas	8	6	422	6
Fire clay, soft	1	6	424	0
Shale, dark, sandy	20	0	444	0
Fire clay, splint	3	0	447	0
Shale, dark, sandy	12	0	459	0
Sandstone, soft, and coal, No. 7 Pocahontas....	21	10	480	10
Shale, blue, sandy	5	0	485	10
Shale, dark	10	0	495	10
Shale, blue, sandy	24	0	519	10
Shale, soft, blue	15	2	535	0
Shale, blue, sandy	4	0	539	0
Sandstone, hard, Eckman	16	0	555	0

Leckie Smokeless Coal Company Coal Test Boring No. 2— No. 5B on Map II.

Meadow Bluff District; on the southwest side of Pollock Knob,
1.6 miles northwest of Anjean; started, Aug. 29, 1930; completed, Sept.
16, 1930; elevation, 3303.9' L.

		Thickness.		Total.	
		Ft.	In.	Ft.	In.
Pottsville Series (325+)					
Surface		4	8	4	8
Shale, gray, sandy		6	4	11	0
Sandstone, reddish		18	0	29	0
Shale, light, sandy		60	0	89	0
Shale, dark, sandy, and slate, black, and coal. Beckley (3198')		16	6	105	6
Fire clay		1	6	107	0
Shale, gray, sandy		2	0	109	0
Sandstone, hard, Quinnimont		51	6	160	6
Shale, dark, sandy		3	6	164	0
Sandstone		3	4	167	4
Sandstone, with shale streaks, shale, sandy, and coal, (Little Fire Creek?) Fire Creek? (3113')		23	4	190	8
Slate, black		2	0	192	8
Fire clay, sandy		2	4	195	0
Sandstone, Pineville		31	6	226	6
Shale, sandy		2	0	228	6
Sandstone		6	0	234	6

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, sandy, and coal, No. 9 Pocahontas	33	9	268	3
Shale, dark, sandy	4	6	272	9
Shale, gray, sandy, and coal, No. 8 Pocahontas	10	8	283	5
Sandstone and shale streaks, Flat top	15	0	298	5
Shale, dark	1	3	299	8
Sandstone	3	0	302	8
Shale, dark, sandy	2	0	304	8
Shale, gray, sandy	3	4	308	0
Sandstone, with shale streaks, shale, dark, and coal, No. 7 Pocahontas (2983')	13	0	321	0
Shale, light, sandy	4	0	325	0

**Leckie Smokeless Coal Company Coal Test Boring No. 1—
No. 5C on Map II.**

Meadow Bluff District; on southeast side of Pollock Knob, 1½ miles north-northwest of Anjean; started, June 17, 1930; completed, July 17, 1930; elevation, 3446.7' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (587' +)				
Surface	3	6	3	6
Sandstone	23	0	26	6
Shale, dark	1	4	27	10
Sandstone	7	2	35	0
Shale, dark	14	6	49	6
Sandstone	2	6	52	0
Shale, dark, sandstone streaks, and coal, Little Raleigh "A"	67	5	119	5
Fire clay, sandy	2	7	122	0
Shale, sandy, light	17	0	139	0
Sandstone	6	0	145	0
Shale, sandy, light, black slate, and coal, Little Raleigh	11	7	156	7
Shale, sandy, light	5	5	162	0
Sandstone	10	0	172	0
Sandstone, with shale streaks	14	0	186	0
Shale, sandy, blue, shale, sandy, dark, and coal, Beckley "Rider"	21	8	207	8
Slate	0	2	207	10
Shale, sandy, dark	8	2	216	0
Shale, sandy, light, black slate, and coal, Beckley (3191')	39	3	255	3
Fire clay, sandy	1	9	257	0
Shale, gray, sandy, and bony coal	25	4	282	4
Fire clay, sandy	3	8	286	0
Shale, dark, sandy, and slate, dark, Fire Creek Coal horizon (3125')	36	0	322	0
Fire clay, sandy	3	0	325	0
Shale, sandy	10	0	335	0
Sandstone	25	0	360	0
Shale, sandy, dark	27	0	387	0
Shale and sandstone streaks	13	0	400	0
Sandstone, shale, dark, sandy, and coal, No. 9 Pocahontas	6	3	406	3
Fire clay, sandy	2	0	408	3

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, gray, sandy and coal, No. 8 Pocahontas	9	5	417	8
Shale, gray, sandy	8	0	425	8
Sandstone	3	4	429	0
Shale, dark, sandy	4	0	433	0
Fire clay, shaly	4	0	437	0
Shale, light, sandy, and coal, No. 7 Pocahontas	21	8	458	8
Fire clay, sandy	0	2	458	10
Shale, dark, sandy	1	10	460	8
Slate, soft	0	2	460	10
Shale, dark	1	0	461	10
Shale, sandy, light, and slate, black, No. 6				
Pocahontas Coal horizon (2948')	36	8	498	6
Shale, light, sandy	6	0	504	6
Shale, dark, sandy	13	6	518	0
Sandstone, hard	30	0	548	0
Sandstone, hard, with coal streaks	5	6	553	6
Sandstone, hard	12	0	565	6
Shale, dark, sandy	4	0	569	6
Sandstone	1	6	571	0
Slate, dark	1	2	572	2
Sandstone, conglomerate, and coal streak, No.				
3 Pocahontas Coal horizon (2862')	12	2	584	4
Sandstone, conglomerate	2	8	587	0

Leckie Smokeless Coal Company Coal Test Boring No. 8— No. 5D on Map II.

Meadow Bluff District; on Pollock Mountain 2.75 miles north of Anjean and 1.25 miles northwest of Sam Creek; started, April 8, 1936; completed, April 24, 1936; elevation, 3367.6' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (175' +)				
Sand, boulders, and clay	38	0	38	0
Shale, sandy	15	0	53	0
Shale, blue, hard, and coal, Beckley "Rider"....	16	9	69	9
Shale, sandy	3	3	73	0
Sandstone	3	0	76	0
Shale, sandy	1	0	77	0
Sandstone	3	6	80	6
Shale, sandy	5	6	86	0
Sandstone	1	0	87	0
Shale, dark, and coal, Beckley (3251')	29	2	116	2
Fire clay, sandy	3	1	119	3
Shale, sandy	3	9	123	0
Sandstone	2	0	125	0
Rock, hard, blue	7	6	132	6
Sandstone, white, crystallized, Quinlmont	30	11	163	5
Shale, sandy, sandstone, and coal, Fire				
Creek (3199')	4	11	168	4
Slate and bone	0	1	168	5
Shale, sandy	0	10	169	3
Fire clay	1	8	170	11
Shale, sandy	4	1	175	0

The record of coal test boring **No. 5E** will be found in Chapter V.

**Leckie Smokeless Coal Company Coal Test Boring No. 6—
No. 5F on Map II.**

Meadow Bluff District; on south end of Pollock Mountain, 0.55 mile northwest of Anjean; started, Nov. 11, 1935; completed, Nov. 21, 1935; elevation, 3295.8' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (158'+)				
Clay, boulders, and sand	16	0	16	0
Sand and boulders	17	0	33	0
Shale, blue	27	6	60	6
Shale, gray, slate, black, and coal, Beckley (3225')	10	8	71	2
Shale, gray	0	10	72	0
Shale, sandy	6	0	78	0
Sandstone	2	0	80	0
Shale, dark-blue	2	6	82	6
Sandstone, hard, white, Quinnimont	23	6	106	0
Shale, sandy, blue, and coal, Fire Creek (3172')	17	8	123	8
Fire clay	3	6	127	2
Slate, gray	8	0	135	2
Shale, sandy	2	0	137	2
Slate, black	0	10	138	0
Slate, gray	5	6	143	6
Slate, black	1	10	145	4
Fire clay	1	4	146	8
Slate, gray	1	8	148	4
Shale, sandy, yellow, and blue	9	8	158	0

**Leckie Smokeless Coal Company Coal Test Boring No. 7—
No. 5G on Map II.**

Meadow Bluff District; on east side of Pollock Mountain, 1.2 miles north of Anjean; started, Dec. 16, 1935; completed, Mar. 31, 1936; elevation, 3296.8' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (198'+)				
Clay, sand, and boulders	25	0	25	0
Shale, blue	46	6	71	6
Slate, black, and coal, Beckley (3213')	12	5	83	11
Shale, light	4	1	88	0
Sandstone, hard, white 18' 0"	43	0	131	0
Shale, blue				
Shale, sandy				
Sandstone, hard, white 11' 0"				
Shale, sandy	24	0	155	0
Sandstone, hard, white, Pineville	43	0	198	0

**Leckie Smokeless Coal Company Coal Test Boring No. 5—
No. 5H on Map II.**

Meadow Bluff District; on east side of Brown Creek, 1.85 miles north of mouth, 1.75 miles northwest of Anjean; started, Sept. 25, 1935; completed, Nov. 1, 1935; elevation, 3385.3' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (290' +)				
Clay, yellow, and boulders	11	0	11	0
Clay layers and sandstone	4	0	15	0
Sandstone, Upper Raleigh	20	0	35	0
Shale, sandy	16	0	51	0
Shale, blue, and coal, Little Raleigh.....	22	7	73	7
Fire clay	4	5	78	0
Shale, gray	2	0	80	0
Shale, gray, sandy	15	7	95	7
Shale and hard blue rock	4	5	100	0
Shale, gray	10	0	110	0
Sandstone, hard, white, Lower Raleigh	33	0	143	0
Slate and coal, Beckley "Rider"	9	4	152	4
Shale, sandy, shale, blue, and coal, Beckley (3184')	48	10	201	2
Fire clay	0	10	202	0
Sandstone, white, fine, Quinnimont	63	0	265	0
Shale, blue	1	10	266	10
Shale, sandy, slate, dark, slate, black, and coal, (Little Fire Creek?) Fire Creek? (3100')	18	7½	285	5½
Slate, black	1	5	286	10½
Shale, sandy	3	1½	290	0

**Leckie Smokeless Coal Company Coal Test Boring No. 3—
No. 5I on Map II.**

Meadow Bluff District; on east side of Brown Creek, 2.75 miles north of mouth and 2.55 miles north of Anjean; started, Aug. 26, 1930; completed, Sept. 4, 1930; elevation, 3436.7' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (330' +)				
Surface	4	0	4	0
Sandstone	1	0	5	0
Shale, gray, sandy	15	0	20	0
Sandstone, hard	6	0	26	0
Shale, dark, sandy	11	0	37	0
Sandstone and coal, Little Raleigh	6	8	43	8
Shale and sandstone streaks	53	4	97	0
Shale, dark, sandy, and coal, Beckley "Rider" (3311')	29	8	126	8
Shale, dark	2	4	129	0
Fire clay, sandy	2	0	131	0
Shale, light, sandy	12	0	143	0
Shale, dark, sandy	40	0	183	0
Sandstone, Quinnimont	31	0	214	0
Shale, light, sandy	21	0	235	0
Shale, dark, sandy, slate, broken up, and coal, Fire Creek (3186').....	15	6	250	6

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Sandstone, hard	18	6	269	0
Shale, light, sandy, and coal, Little Fire Creek (2161')	6	6	275	6
Fire clay	1	6	277	0
Shale, gray, sandy	2	0	279	0
Slate, black	3	6	282	6
Fire clay	1	6	284	0
Shale, light, sandy	32	6	316	6
Shale, dark	1	6	318	0
Shale, gray, sandy, and coal, No. 9 Pocahontas	8	0	326	0
Shale, dark, sandy	0	7	326	7
Shale, light, sandy	3	5	330	0

**Leckie Smokeless Coal Company Coal Test Boring No. 10A—
No. 5J on Map II.**

Meadow Bluff District; on west side of Pollock Mountain, 3.6 miles north of Anjean and 2 miles west of Duo; started, May 23, 1936; completed, June 2, 1936; elevation, 3637.4' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (80'+)				
Clay, yellow	8	0	8	0
Shale, yellow	4	0	12	0
Clay, yellow, and sand	3	0	15	0
Shale, yellow	2	6	17	6
Shale, blue, and coal, Sewell "A"	27	4	44	10
Fire clay	2	2	47	0
Shale, blue	3	0	50	0
Shale, sandy, and coal, Sewell (3560' L.)	27	5	77	5
Fire clay	0	4	77	9
Shale, sandy	2	3	80	0

**Leckie Smokeless Coal Company Coal Test Boring No. 10—
No. 5K on Map II.**

Meadow Bluff District; on west side of Pollock Mountain, 3.6 miles north of Anjean and 2.05 miles west of Duo; started, May 5, 1936; completed, May 25, 1936; elevation, 3597.4' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (324'+)				
Sand and clay; sandstone and shale, broken; shale and coal, Sewell (3567')	30	6	30	6
Fire clay	3	4	33	10
Sandstone	8	2	42	0
Shale, blue	17	0	59	0
Slate, black	3	0	62	0
Shale, blue	5	0	67	0
Sandstone, crystallized	17	0	84	0
Slate, black, and coal, Welch	12	0	96	0
Fire clay	2	6	98	6
Shale, dark	7	6	106	0
Slate, black	3	0	109	0
Shale, gray	7	0	116	0
Shale, sandy	59	0	175	0

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Shale, dark-blue, slate, black, and coal, Little Raleigh	15	5	190	5
Fire clay	0	7	191	0
Shale, hard, blue	18	0	209	0
Shale, sandy	28	0	237	0
Sandstone, hard, Lower Raleigh	30	0	267	0
Slate, black	36	0	303	0
Slate, blue, and coal, Beckley (3287')	7	7	310	7
Fire clay, sandy	4	5	315	0
Shale, sandy	7	0	322	0
Sandstone, hard, Quinlmont	2	6	324	6

**Leckie Smokeless Coal Company Coal Test Boring No. 11—
No. 5L on Map II.**

Meadow Bluff District; on east side of Huggins Ridge, 3.8 miles north of Anjean and 2.3 miles west of Duo; started, June 8, 1936; completed, June 19, 1936; elevation, 3695.3' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (215'+)				
Clay and boulders	8	0	8	0
Sandstone, brown 20' 0"	34	0	42	0
Sandstone, gray 14 0				
Shale, sandy and coal, Castle (3587')	65	11	107	11
Fire clay	0	1	108	0
Shale, sandy	4	0	112	0
Sandstone and shale streaks	28'	0"	54	0
Sandstone	5	6		
Shale, blue	5	6		
Sandstone	15	0		
Shale, sandy, slate, blue, and coal, Sewell "A"	15	2	181	2
Shale, sandy	6	2	187	4
Sandstone and shale streaks	1'	3"	20	2
Sandstone	1	4		
Sandstone and shale streaks	1	1		
Sandstone	4	0		
Shale, sandy	1	10		
Sandstone	10	8		
Shale, sandy, slate, dark, and coal, Sewell (3483')	4	6	212	0
Fire clay	1	7	213	7
Shale, sandy	1	5	215	0

**Leckie Smokeless Coal Company Coal Test Boring No. 12—
No. 5M on Map II.**

Meadow Bluff District; at head of Brown Creek, 4.7 miles north of Anjean and 2.5 miles northwest of Duo; started, June 27, 1936; completed, July 10, 1936; elevation, 3497.6' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (470'+)				
Clay and silt	4	0	4	0

				Thickness.		Total.					
				Ft.	In.	Ft.	In.				
Shale, sandy, and coal, Castle				26	8	30	8				
Slate, black				0	4	31	0				
Sandstone and shale				Guyandot	44	75	0				
streaks	10'	0"									
Sandstone	11	0									
Sandstone and shale				Guyandot	44	75	0				
streaks	23	0									
Slate, dark-blue				8	4	83	4				
Shale, sandy				1	8	85	0				
Sandstone				7	6	92	6				
Shale, sandy, shale, hard, blue, slate, black, and coal, Sewell "A"				14	9	107	3				
Slate, black				0	5	107	8				
Shale, sandy, slate, black, and coal, Sewell, (3367')				23	0	130	8				
Fire clay				3	6	134	2				
Sandstone	24'	4"	Welch?	31	10	166	0				
Sandstone, with shale streaks	7	6									
Slate, dark				5	4	171	4				
Sandstone				13	8	185	0				
Shale, sandy				6	0	191	0				
Slate, blue				7	6	198	6				
Shale, sandy				1	8	200	2				
Slate, black, and coal, Welch?				7	2	207	4				
Slate, black				0	8	208	0				
Shale, sandy				6	0	214	0				
Sandstone and shale				Upper Raleigh?	66	280	0				
streaks	6'	6"									
Sandstone, hard	8	6									
Shale, blue, and sand- stone				Upper Raleigh?	66	280	0				
stone	51	0									
Slate, black				3	6	283	6				
Sandstone, hard, and shale				9	6	293	0				
Slate, black				6	0	299	0				
Shale, dark-blue, slate, black, and coal, Little Raleigh? (Beckley?)				23	3	322	3				
Fire clay				2	3	324	6				
Shale, gray				19	0	343	6				
Sandstone, hard	13'	0"	Lower Raleigh?	66	6	410	0				
Shale, gray, sandy	13	6									
Sandstone, hard	18	0									
Shale, gray	1	5									
Sandstone, hard, white	17	7									
Sandstone and shale				Lower Raleigh?	66	410	0				
streaks	3	0									
Slate, black								1	8	411	8
Shale, sandy								25	2	436	10
Slate, dark-blue								33	2	470	0

The records of coal test borings Nos. 6 and 7 will be found in Chapter V.

The following three records were furnished the Survey by Mr. J. W. Raine, of Duo:

**Raine Lumber and Coal Company Coal Test Boring No. 6—
No. 8 on Map II.**

Meadow Bluff District; near the northern end of Smokehouse
Ridge, two miles east of Duo; elevation, 4085' L.

				Thickness.		Total.	
Pottsville Series—New River Group (432' +)				Ft.	In.	Ft.	In.
Surface				3	0	3	0
Sandstone, hard	39'	6"	Lower Nuttall	93	0	96	0
Shale, dark, with sand streaks	53	6		0	3	96	3
Clay, with coal streaks, laeger "A"				4	11	101	2
Fire clay							
Shale, dark, sandy	9'	6"	Upper laeger	44	10	146	0
Shale, dark	6	0					
Sandstone	0	6					
Shale, dark, with sand streaks	5	6					
Shale, dark	23	4					
Coal, Hughes Ferry (3938')				1	6	147	6
Fire clay, dark				2	4	149	10
Sandstone and shale				3	7	153	5
Shale, dark, with sandy streaks				18	8	172	1
Shale, dark, with lime (?) streaks				24	5	196	6
Coal	1'	1"	Lower laeger				
Fire clay	2	4		3	10	200	4
Coal	0	5					
Clay, sandy				11	0	211	4
Sandstone				4	5	215	9
Shale, sandy				0	7	216	4
Sandstone				1	8	218	0
Shale, sandy				0	8	218	8
Sandstone				2	9	221	5
Shale, sandy				2	8	224	1
Sandstone				0	10	224	11
Shale, sandy				2	6	227	5
Sandstone	16'	2"	Harvey Conglomerate				
Shale, dark	2	9					
Sandstone	5	9		30	11	258	4
Shale, dark, sandy	1	5					
Sandstone	4	6					
Coal	0	1					
Sandstone	0	3					
Shale, dark, sandy, Sandy Huff				21	8	280	0
Coal, Castle (3804')				1	4	281	4
Fire clay				4	0	285	4
Clay, sandy				2	9	288	1
Shale, dark, sandy				2	5	290	6
Sandstone	9'	2"	Guyandot Sandstone				
Sandstone, with shale streaks	17	8					
Shale, dark	4	7					
Shale, light	2	0		82	9	373	3
Sandstone	17	6					
Shale, dark, sandy	2	8					
Sandstone	14	0					
Limestone (?)	1	2					
Sandstone	14	0					

				Thickness.		Total.	
				Ft.	In.	Ft.	In.
Slate, black				5	9	379	0
Coal, cannel.....	0'	6"	} Sewell "A"	1	4	380	4
Coal	0	10					
Fire clay				1	10	382	2
Shale, dark, sandy				6	0	388	2
Sandstone, Lower Guyandot				7	4	395	6
Shale, dark.....				14	10	410	4
Slate, black	3'	9"	} Hartridge				
Shale, dark	6	11		10	8	421	0
Coal, cannel	0'	3"	} Sewell (3660')	3	11	424	11
Coal	3	8					
Fire clay, dark				7	1	432	0

**Raine Lumber and Coal Company Coal Test Boring No. 3—
No. 9 on Map II.**

Meadow Bluff District; 2.5 miles east of Duo; elevation, 3990' L.

				Thickness.		Total.	
				Ft.	In.	Ft.	In.
Pottsville Series—New River Group (450'+)							
Surface				9	6	9	6
Shale, gray, sandy, Lower Iaeger.....				20	0	29	6
Sandstone, Harvey Conglomerate.....				87	6	117	0
Shale, dark, and sandstone streaks, Sandy Huff				8	6	125	6
Slate				0	6	126	0
Coal, Castle (3863')				1	1	127	1
Fire clay				2	9	129	10
Shale, gray, sandy	14'	2"	} Guyandot Sandstone				
Sandstone, with shale streaks	12	0		34	8	164	6
Shale, dark, with sandstone streaks	8	6					
Shale, dark				11	0	175	6
Shale, dark, sandy				10	6	186	0
Sandstone				12	0	198	0
Shale, dark				8	0	206	0
Slate, black				4	2	210	2
Coal, Sewell "A"				0	7	210	9
Fire clay, sandy				2	3	213	0
Shale, dark, sandy				15	6	228	6
Shale, dark				40	6	269	0
Slate, black, Hartridge Black Shale.....				13	10	282	10
Coal, cannel	0'	3"	} Sewell (3696')				
Slate, black	1	4		11	7	294	5
Coal, dirty	1	6					
Slate	6	3					
Coal	1	3					
Fire clay, dark, with coal spars.....	1	0					
Fire clay, sandy				0	5	294	10
Shale, gray, sandy				1	6	296	4
Sandstone, hard, Welch				34	0	330	4
Shale, dark				1	8	332	0
Coal, Welch				0	2	332	2

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Sandstone, hard, Upper Raleigh	25	6	357	8
Shale, dark, sandy	2	0	359	8
Fire clay, dark	3	5	363	1
Fire clay, shaly	5	6	368	7
Sandstone and coal spars	20	5	389	0
Shale, dark, and sandstone, mixed	4	0	393	0
Shale, dark, sandy	8	6	401	6
Coal, dirty, Little Raleigh "A"	0	11	402	5
Fire clay	1	7	404	0
Shale, dark, sandy	18	6	422	6
Shale, gray, sandy	19	6	442	0
Sandstone	8	0	450	0

**Raine Lumber and Coal Company Coal Test Boring No. 7—
No. 10 on Map II.**

Meadow Bluff District; near Job Knob, 3 miles east of Duo; elevation, 4240' L.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—New River Group (460'+)	3	0	3	0
Surface				
Sandstone, brown.... 38' 6"	90	9	93	9
Shale, dark 1 6				
Sandstone, with shale streaks 50 9				
Coal, bony, laeger "A"	0	4	94	1
Fire clay, sandy	3	8	97	9
Sandstone, hard	8	3	106	0
Shale, dark, with sandstone streaks 30' 0"	36	2	142	2
Shale, dark 6 2				
Slate, black	2	10	145	0
Coal, Hughes Ferry (4094')	1	1	146	1
Sandstone, Middle laeger	10	6	156	7
Shale, dark, sandy	16	4	172	11
Shale, dark, Lower laeger?	19	9	192	8
Sandstone, Harvey Conglomerate?	55	4	248	0
Shale, dark	43	0	291	0
Shale, dark, sandy, Sandy Huff	10	0	301	0
Shale, dark	0	6	301	6
Coal, Castle	0	10	302	4
Sandy clay	6	8	309	0
Sandstone, Guyandot	10	0	319	0
Shale, dark, sandy	30	0	349	0
Shale, dark	35	6	384	6
Coal, Sewell "A"	0	6	385	0
Sandy clay	2	0	387	0
Sandstone	2	0	389	0
Shale, dark, sandy	51	0	440	0
Slate, black, Hartridge Black Shale	10	0	450	0
Coal, cannel 0' 2"	2	6	452	6
Coal 2 4				
Fire clay, light	1	6	454	0
Fire clay, dark	6	0	460	0

Attention is called to the fact that in borings Nos. 11 to 15 inclusive the measurements were not always made at right angles to the bedding-plane of the formations penetrated. Only parts of the cores of Nos. 11, 14, and 15 were found but they showed a variation of 3° to 20° off vertical. The harder sandstone beds caused the greater migration.

The record of boring No. 11 will be found on pages 172-4, Chapter V.

Gauley Coal Land Company Coal Test Boring No. 3— No. 12 on Map II.

Meadow Bluff District; on Rockcamp Ridge, 6.7 miles northeast of Anjean and 4.1 miles east of Duo; elevation, 3951' L.

Pottsville Series—New River Group (350' 6"+)				Thickness.		Total.		
				Ft.	In.	Ft.	In.	
Surface				4	0	4	0	
Sandstone, hard 21' 0"				Lower Raleigh	37	6	41	6
Sandstone, hard,								
coal spars 16 6								
Shale, dark, soft.....				3	0	44	6	
Sandstone, hard, dark, shale mixed				10	0	54	6	
Shale, dark, sandy				30	0	84	6	
Shale, dark				46	6	131	0	
Coal 1' 6"				Beekley (3718')	2	4	133	4
Bone coal 0 10								
Slate, black								
Shale, gray				0	4	133	8	
Sandstone				2	6	136	2	
Shale, gray				0	8	136	10	
Shale, gray, sandy				7	2	144	0	
Sandstone, hard, coal				Quinnimont ...	21	9	177	0
spars 10' 0"								
Coal 0 5								
Fire clay 0 4				Quinnimont	42	0	219	0
Sandstone, hard, white 11 0								
Shale, dark, sandy 17' 0"								
Shale, gray, sandy 25 0				Shale				
Fire clay, hard				3	0	222	0	
Shale, gray, sandy				7	0	229	0	
Slate, black, Fire Creek Coal horizon				1	6	230	6	
Shale, gray				21	0	251	6	
Slate, black, Little Fire Creek Coal horizon....				4	0	255	6	
Shale, gray, sandy				26	6	282	0	
Coal and slate.....				0	2	282	2	
Shale, dark, sandy				11	6	293	8	
Bone coal				0	2	293	10	
Fire clay				0	6	294	4	
Shale, gray				3	6	297	10	
Sandstone				4	10	302	8	
Shale, dark, sandy				6	0	308	8	
Slate				1	6	310	2	

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Slate, black, and fire clay mixed.....	1	4	311	6
Fire clay, soft	1	0	312	6
Shale, gray	17	6	330	0
Shale, gray, sandy	20	6	350	6
Pottsville Series—Pocahontas Group (115' +)				
Sandstone, hard, Flattop	40	0	390	6
Slate, black	1	4	391	10
Fire clay, dark	1	2	393	0
Shale, gray	5	0	398	0
Shale, gray, sandy	8	6	406	6
Sandstone, Pierpont	31	2	437	8
Coal, No. 6 Pocahontas?	0	1	437	9
Fire clay, soft	0	4	438	1
Fire clay, sandy	1	0	439	1
Shale, gray, sandy	11	11	451	0
Sandstone	14	6	465	6

The record of boring No. 13 will be found on pages 174-6 in Chapter V.

Gauley Coal Land Company Coal Test Boring No. 2— No. 14 on Map II.

Meadow Bluff-Williamsburg District line; seven miles N. 80° E. of Anjean on Little Clear Creek Mountain; elevation, 4168' L.

	Thickness.		Total.		
	Ft.	In.	Ft.	In.	
Pottsville Series—New River Group (280' 4" +)					
Surface	3	6	3	6	
Shale, brown, broken.....	5	0	8	6	
Shale, brown, sandy	36	6	45	0	
Shale, dark	9	8	54	8	
Slate, black	0	3	54	11	
Coal, Beckley (4110')	3	2	58	1	
Fire clay, light	3	7	61	8	
Shale, gray, sandy	3	4	65	0	
Sandstone, hard 43' 0"	55	7	120	7	
Sandstone, hard,					} Quinnimont...
with coal spars 12 7					
Slate, black, soft.....	0	9	121	4	
Coal, Fire Creek	1	2	122	6	
Clay shale, soft, dark	5	0	127	6	
Shale, dark	8	2	135	8	
Shale, dark, sandy	26	4	162	0	
Sandstone, hard, broken 57' 0"	69	0	231	0	
Sandstone, fine-grained 12 0					} Pineville....
Shale, dark, sandy					
Slate, black	0	10	234	1	
Bone coal, No. 9 Pocahontas?	0	7	234	8	
Shale, gray, sandy	7	4	242	0	
Sandstone	4	0	246	0	
Shale, dark, sandy	12	0	258	0	
Shale, dark	14	8	272	8	

			Thickness.		Total.		
			Ft.	In.	Ft.	In.	
Bone coal	0'	2"	No. 8 Pocahontas?..	3	8	270	4
Slate	0	11					
Bone coal	0	1					
Clay shale, dark	2	3					
Bone coal	0	3					
Shale, gray			4	0	280	4	
Pottsville Series—Pocahontas Group (286' 8"±)							
Sandstone, Flattop			25	0	305	4	
Coal, No. 7 Pocahontas			0	4	305	8	
Fire clay			1	8	307	4	
Shale, dark			17	2	324	6	
Shale, dark, sandy			21	0	345	6	
Sandstone, hard	5'	0"	Pierpont Sandstone	15	4	360	10
Shale, hard, gray, sandy	4	0					
Sandstone, hard	6	4					
Coal, No. 6 Pocahontas			1	0	361	10	
Clay shale, gray, sandy			1	2	363	0	
Sandstone			2	7	365	7	
Shale, gray			1	2	366	9	
Fire clay			4	6	371	3	
Clay shale			10	9	382	0	
Slate, gray			4	0	386	0	
Slate, coal, and sulphur			0	3	386	3	
Fire clay			0	11	387	2	
Slate and coal spars			0	3	387	5	
Shale, dark			5	5	392	10	
Slate, soft, broken			0	10	393	8	
Coal			0	10	394	6	
Fire clay			3	6	398	0	
Shale, gray			6	0	404	0	
Slate, black			1	0	405	0	
Fire clay			2	0	407	0	
Shale, gray			14	6	421	6	
Sandstone, Upper Pocahontas			34	0	455	6	
Slate, gray			0	2	455	8	
Coal, No. 3 Pocahontas?			0	1	455	9	
Fire clay			0	3	456	0	
Shale, gray, sandy			5	0	461	0	
Shale, gray			4	6	465	6	
Fire clay			1	0	466	6	
Shale, gray			3	6	470	0	
Clay shale, gray			3	6	473	6	
Shale, gray			16	0	489	6	
Slate and coal, No. 3 Pocahontas Coal?			0	3	489	9	
Fire clay			3	3	493	0	
Shale, gray			8	6	501	6	
Sandstone, Lower Pocahontas?			28	0	529	6	
Coal, No. 2 "A" Pocahontas?			0	4	529	10	
Sandstone			4	8	534	6	
Sandstone, with shalo streaks			6	0	540	6	
Shale, dark, sandy			15	8	556	2	
Slate, black	3'	0"	No. 2 Pocahontas Coal horizon?	3	10	560	0
Slate, black, soft	0	10					
Fire clay, sandy			2	0	562	0	
Shale, gray, sandy			6	0	568	0	
Shale, gray			7	0	575	0	

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Black slate, coal spars, No. 1 Pocahontas	0	3	575	3
Fire clay	1	9	577	0

DETAILED COAL TEST RECORDS, WILLIAMSBURG DISTRICT.

In Williamsburg District one test hole was drilled for coal. The record of this hole (No. 15) is supporting evidence of the comparative rapid dip of the rocks in the region of Grassy Knob. The top of the red shale that was found at an elevation of 3182' in boring No. 11, (1.8 miles northwest) was found at an elevation of 3868' in boring No. 15.

As stated on a foregoing page the measurements given in the following record are probably not true vertical measurements:

Raine Lumber and Coal Company Coal Test Boring No. 1— No. 15 on Map II.

Williamsburg District; 0.3 mile N. 70° E. from the Grassy Knob triangulation point; elevation, 4125' B.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series—Pocahontas Group (256'+)				
Surface	6	0	6	0
Sandstone, Pierpont	33	0	39	0
Shale, dark	13	0	52	0
Coal	1'	1"	No. 6 Pocahontas (4070')	
Fire clay, with coal spars.....	1	3		
Coal	1	1½		
Fire clay, soft.....	0	1½	55	7
Fire clay, hard	1	5	57	0
Shale, gray, sandy	27	0	84	0
Slate, black	1	0	85	0
Fire clay, soft	11	6	96	6
Sandstone	6	4	102	10
Shale, gray, sandy	28	2	131	0
Sandstone	11	6	142	6
Shale, gray, sandy	3	4	145	10
Shale, dark clay	9	6	155	4
Coal and clay mixed 1' 1"	0	7	No. 3 Pocahontas?	
Slate				
Coal, dirty				
Shale, gray	2	4	157	8
Shale, gray, sandy	2	4	160	0
Shale, gray clay	11	0	171	0
Shale, gray clay	2	0	173	0
Fire clay, with coal streaks	0'	8"	No. 3 Pocahontas? (3949')	
Fire clay, dark	1	5		
Black slate, coal spars	0	9		

For the Calendar Year 1927.

Name of Company	Name of Mine	Production of Coal. (Tons of 2000 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Elk Lick Coal Co.....	Spruce Knob.....	29,421	1,807	291	27,323
Frances Coal Co.....	Frances	77,456	893	76,563
Greenbrier Smokeless Coal Co.....	Bellburn	249,798	4,200	245,598
Imperial Smokeless Coal Co.....	Quhuwood	358,933	672	358,261
Johnstown Coal & Coke Co.....	Criehton	168,176	3,510	164,666
Margarette Coal Co.....	Margarette No. 2.....	356,360	7,200	1,685	347,475
Meadow River Fuel Co.....	Lincoln	1,064	830	234
Nelson Fuel Co.....	Nelson No. 1.....	358,954	3,230	355,724
Totals.....	1,600,162	9,007	15,311	1,575,844

For the Calendar Year 1928.

Name of Company	Name of Mine	Production of Coal. (Tons of 2000 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Elk Lick Coal Co.....	Spruce Knob and Turkey Run	16,657	874	193	15,590
Frances Coal Co.....	Frances	58,144	465	57,679
Greendrier Smokeless Coal Co.....	Bellburn	263,629	396	263,233
Imperial Smokeless Coal Co....	Quinwood	374,452	724	373,728
Johnstown Coal & Coke Co....	Crichton	191,927	765	191,162
Leckle Smokeless Coal Co....	Leckle Nos. 1, 2, 3, 4, & 5....	54,914	200	54,714
Margarette Coal Co.....	Margarette	339,089	6,500	1,340	331,249
Meadow River Fuel Co.....	Lincoln	2,000	2,000
Nelson Fuel Co.....	Nelson No. 1.....	357,397	3,500	353,897
Totals.....		1,658,209	7,374	9,583	1,641,252

For the Calendar Year 1929.

Name of Company	Name of Mine	Production of Coal. (Tons of 2000 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Clear Creek Coal Co.....	Clearco No. 1.....	907	175	750
Frances Coal Co.....	Frances	72,547	72,547
Greenbrier Smokeless Coal Co.....	Bellburn	269,822	230	269,592
Imperial Smokeless Coal Co.....	Quinwood	382,785	1,322	381,463
Johnstown Coal & Coke Co.....	Crichton No. 1.....	206,628	819	205,809
Leekie Smokeless Coal Co.....	Leekle Nos. 1, 2, 3, 4, & 5.....	252,775	775	252,000
Margarette Coal Co.....	Margarette No. 1.....	337,268	6,000	2,000	329,268
Meadow River Fuel Co.....	Lineoln	4,430	3,546	884
New River & Poca, Cons. Coal Co.....	Leslle	261,747	4,241	257,506
Puck Brothers.....	Dwyer	1,120	1,120
Totals.....	1,790,029	6,000	13,090	1,770,939

For the Calendar Year 1930.

Name of Company	Name of Mine	Production of Coal. (Tons of 2000 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Clear Creek Coal Co.....	Clearco Nos. 1 & 2.....	45,034	194	44,840
Frances Coal Co.....	Frances	97,675	96,675
Greenbrier Smokeless Coal Co.....	Orlehton No. 2.....	197,112	896	196,216
Imperial Smokeless Coal Co..	Quinwood	319,846	1,508	318,338
Johnstown Coal & Coke Co....	Orlehton	208,274	550	207,724
Leekio Smokeless Coal Co....	Leekie	327,992	800	327,192
Margaretto Coal Co.....	Margarette	386,177	6,000	4,000	376,177
Meadow River Fuel Co.....	Lincoln	9,353	3,726	5,627
New River & Poca. Cons. Coal Co.....	Leslie	433,169	6,439	426,730
Tuek Brothers.....	Dwyer	7,850	100	7,750
Totals.....		2,032,482	6,000	18,213	2,008,269

For the Calendar Year 1931.

Name of Company	Name of Mine	Production of Coal. (Tons of 2000 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Clear Creek Coal Co.....	Brooke Nos. 1 & 2.....	112,211	125	224	111,862
Frances Coal Co.....	Frances	58,219	500	500	57,219
Greenbrier Smokeless Coal Co.....	Cricketon No. 2.....	152,701	1,584	151,117
Imperial Smokeless Coal Co.....	Quinwood	248,117	1,116	247,031
Johnstown Coal & Coke Co.....	Cricketon No. 1.....	174,280	1,021	173,259
Leekle Smokeless Coal Co.....	Leekle	278,477	977	277,500
Margarette Coal Co.....	Margarette	372,113	6,000	2,000	364,113
Meadow River Fuel Co.....	Lincoln	1,102	1,102
Midland Smokeless Coal Co.....	Midland No. 1.....	1,851	955	896
New River & Poca, Cons. Coal Co.....	Leslie	374,100	5,650	368,450
Quek Brothers.....	Dwyer	12,771	800	41,971
Totals.....	1,816,272	6,625	16,229	1,793,418

For the Calendar Year 1932.

Name of Company	Name of Mine	Production of Coal. (Tons of 2000 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Clear Creek Coal Co.....	Brooke Nos. 1 & 2.....	68,051	156	155	67,740
Frances Coal Co.....	Frances	58,865	1,200	57,665
Greenbrier Smokeless Coal Co.....	Crichton No. 2.....	128,886	1,014	127,872
Imperial Smokeless Coal Co....	Quinwood	190,997	562	190,435
Johnstown Coal & Coke Co....	Crichton No. 1	123,248	769	122,479
Leckle Smokeless Coal Co....	Leckle	275,087	1,500	3,500	270,087
Margarette Coal Co.....	Margarette	254,125	8,000	4,200	241,925
Meadow River Fuel Co.....	Lineoln	2,528	2,076	452
Midland Smokeless Coal Co....	Midland	5,270	1,136	4,140
Now River & Poca. Cons. Coal Co.....	Leslie	236,212	2,537	233,675
Totals.....	1,343,269	9,656	17,143	1,316,470

For the Calendar Year 1933.

Name of Company	Name of Mine	Production of Coal. (Tons of 2000 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Clear Creek Coal Co.....	Brooke	126,278	159	212	125,907
Frances Coal Co.....	Frances	64,857	795	64,062
Greenbrier Smokeless Coal Co.....	Crichton No. 2	136,862	1,297	135,565
Imperial Smokeless Coal Co.....	Quinwood	240,074	536	239,538
Johnstown Coal & Coke Co....	Crichton No. 1	152,092	626	151,466
Leekie Smokeless Coal Co....	Leckle	317,296	2,099	315,197
Margarette Coal Co.....	Margarette	294,039	7,141	5,709	281,189
Meadow River Fuel Co.....	Lincoln	848	698	150
Midland Smokeless Coal Co....	Midland	4,256	658	3,598
New River & Poca, Cons. Coal Co.....	Leslie	275,359	130	4,801	270,428
Ralne Lumber & Coal Co.....	Duo	28,119	12	28,107
Totals.....		1,640,080	7,430	17,443	1,615,207

For the Calendar Year 1934.

Name of Company	Name of Mine	Production of Coal. (Tons of 2000 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Burley Coal Co.....	Burley	19,382	123	23	19,236
Clear Creek Coal Co.....	Brooke	153,642	132	318	153,192
Frances Coal Co.....	Frances	58,001	699	57,302
Greenbrier Smokeless Coal Co.....	Orlehton No. 2.....	120,629	1,319	119,310
Imperial Smokeless Coal Co..	Quinwood	243,728	564	243,164
Johnstown Coal & Coke Co....	Orlehton No. 1.....	148,610	683	147,927
Leekle Smokeless Coal Co....	Leekle	368,419	2,450	1,464	364,505
Margarette Coal Co.....	Margarette	279,814	12,000	4,000	263,814
Meadow River Fuel Co.....	Lineoln	1,044	1,044
Midland Smokeless Coal Co..	Midland	3,735	810	2,925
New River & Poca. Cons. Coal Co.....	Leslie	313,635	10,218	303,417
Ralme Lumber & Coal Co.....	Ralme	36,390	282	36,108
Totals.....	1,747,029	14,705	21,424	1,710,900

For the Calendar Year 1935.

Name of Company.	Name of Mine.	Production of Coal. (Tons of 2,000 lbs.)	Distribution of Coal.	
			Used Locally and Stocked.	Railroad Shipments.
Burley Coal Co.....	Burley	4,526	4,526
Clear Creek Coal Co.....	Brooko	161,006	555	160,451
Frances Coal Co. (I. F. Vass and Bonner H. Hill, Trustees)	Frances	31,834	763	31,071
Greenbrier Smokeless Coal Co.....	Crichton No. 2.....	124,080	1,514	122,566
Imperial Smokeless Coal Co.....	Quinwood	202,252	1,114	201,138
Johnstown Coal & Coke Co.....	Crichton No. 1.....	139,984	537	139,447
Leeklo Smokeless Coal Co.....	Leekle	321,964	321,964
Margarette Coal Co. (I. F. Vass and Bonner H. Hill, Trustees).....	Margarette	227,430	12,552	214,878
Meadow River Fuel Co.....	Lincoln	1,330	1,000	330
Midland Smokeless Coal Co.....	Midland	17,439	385	17,054
New River & Poca, Cons. Coal Co.....	Lestle	300,323	14,810	285,513
Ralme Lumber & Coal Co.....	Duo	55,868	257	56,391
Totals	1,588,036	33,487	1,555,329
Truck Mines	3,600	3,600
Totals	1,591,636	37,087	1,555,329

For the Calendar Year 1936.

Company.	Mine.	Production of Coal (Tons of 2,000 lbs.
Clear Creek Coal Co.....	Brooke Nos. 1 and 2	161,366
a—Franees Coal Co.....	Franees	15,160
*A—Gauley Coal Land Co.....
Greenbrier Smokeless Coal Co..	Criehton No. 2.....	167,780
C—Greenbrier Firecreek Coal Co....	Midland	10,979
Imperial Smokeless Coal Co.....	Quinwood	240,737
Johnstown Coal & Coke Co.....	Criehton No. 1.....	183,716
Leekie Smokeless Coal Co.....	Leekie Nos. 1 and 2	327,207
b—Margarette Coal Co.....	Margaretto
B—Margaretto Coal Corp.....	Margarette	234,737
*—Meadow River Fuel Co.....	Lincoln	225
c—Midland Smokeless Coal Co.....	Midland	10,791
New River & Poa. Cons. C. Co.	Leslie	360,594
Raine Lumber & Coal Co.....	Duo	76,719
Total	1,790,011

*Ceased (Mine or Company Ceased Operation).

a—Predecessor.

A—Successor.

b—Predecessor.

B—Successor.

c—Predecessor.

C—Successor.

RECORDS OF COAL TEST BORINGS.

SUMMARIZED RECORDS.

Within Greenbrier County 31 holes have been drilled for testing coal, 27 of which are located in Meadow Bluff District, one in Williamsburg District, and the remaining three in Irish Corner District. In the near-by parts of Nicholas, Fayette, and Monroe Counties there have been 78 cores drilled, most of which have bearing on the coal resources of Greenbrier County. Since the records of most of these holes were not available at the time of publication of the previous reports, it is deemed advisable to include them in this volume. The records of these holes have been correlated and will be found on succeeding pages.

It will be noted that the numbers given to the holes in adjoining counties are not always in sequence; this was done to avoid, so far as possible, the renumbering of cores listed in previous publications of the Survey.

The following table, while lacking some of the details it should contain, gives the surface elevation, ownership, and, when available, the key number on Map II, by which the locations of the borings may be found. In the elevation column the letter "L" signifies a spirit-level determination, the sign " \pm " means that the elevation was taken from the topographic map, and the letter "B" indicates that an aneroid barometer was used, checked on the nearest Government elevation. A question mark beside the depth to the various coals indicates uncertainty of the correlation. The following abbreviations of company names have been used:

W. E. Deegans C. C.....	W. E. Deegans Coal Corporation.
Margarette Coal, et al.....	Margarette Coal Company and W. E. Deegans.
W. Va. Coal & Coke.....	West Virginia Coal & Coke Com- pany.
Brackens Cr.....	Brackens Creek Coal & Land Com- pany.
Babcock C. & C.....	Babcock Coal & Coke Company.
N. R. & P. C. C.....	New River & Pocahontas Consol- dated Coal Company.

Summarized Record of Coal Test Borings

No. on Map II.	Name of Property.	Magisterial District.	Company.	Surface Elevation.
1	Mrs. E. T. Martin No. 1.....	Meadow Bluff	Nutter Hrs.	1930'B
2	Meadow River Coal & Land Co....	Meadow Bluff	W. E. Deegans C. C.1
3	Meadow River Coal & Land Co....	Meadow Bluff	W. E. Deegans C. C.1
4	Meadow River Coal & Land Co....	Meadow Bluff	W. E. Deegans C. C.1	3125'B
5	Gauley Coal Land Co.....	Meadow Bluff	Margarette Coal, et al.
5A	Leckie Smokeless Coal Co. No. 4....	Meadow Bluff	Leckie Smokeless Coal..	3357'L
5B	Leckie Smokeless Coal Co. No. 2....	Meadow Bluff	Leckie Smokeless Coal..	3304'L
5C	Leckie Smokeless Coal Co. No. 1....	Meadow Bluff	Leckie Smokeless Coal..	3447'L
5D	Leckie Smokeless Coal Co. No. 8....	Meadow Bluff	Leckie Smokeless Coal..	3368'L
5E	Gauley Coal Land Co. No. 30.....	Meadow Bluff	Gauley Coal Land.....	3832'
5F	Leckie Smokeless Coal Co. No. 6....	Meadow Bluff	Leckie Smokeless Coal..	3296'L
5G	Leckie Smokeless Coal Co. No. 7....	Meadow Bluff	Leckie Smokeless Coal..	3297'L
5H	Leckie Smokeless Coal Co. No. 5....	Meadow Bluff	Leckie Smokeless Coal..	3385'L
5I	Leckie Smokeless Coal Co. No. 3....	Meadow Bluff	Leckie Smokeless Coal..	3437'L
5J	Gauley Coal Land Co. No. 10A.....	Meadow Bluff	Leckie Smokeless Coal..	3637'L
5K	Gauley Coal Land Co. No. 10.....	Meadow Bluff	Leckie Smokeless Coal..	3597'L
5L	Gauley Coal Land Co. No. 11.....	Meadow Bluff	Leckie Smokeless Coal..	3695'L
5M	Gauley Coal Land Co. No. 12.....	Meadow Bluff	Leckie Smokeless Coal..	3498'L
6	Raine Lumber & Coal Co. No. 5....	Meadow Bluff	Raine Lumber & Coal..	3630'L
7	Raine Lumber & Coal Co. No. 4....	Meadow Bluff	Raine Lumber & Coal..	4015'L
8	Raine Lumber & Coal Co. No. 6....	Meadow Bluff	Raine Lumber & Coal..	4086'L
9	Raine Lumber & Coal Co. No. 3....	Meadow Bluff	Raine Lumber & Coal..	3990'L
10	Raine Lumber & Coal Co. No. 7....	Meadow Bluff	Raine Lumber & Coal..	4240'L
11	Raine Lumber & Coal Co. No. 2....	Meadow Bluff	Raine Lumber & Coal..	4010'L
12	Gauley Coal Land Co. No. 3.....	Meadow Bluff	Gauley Coal Land.....	3951'L
13	Gauley Coal Land Co. No. 1.....	Meadow Bluff	Gauley Coal Land.....	3808'L
14	Gauley Coal Land Co. No. 2.....	Meadow Bluff	Gauley Coal Land.....	4168'L
15	Raine Lumber & Coal Co. No. 15.	Williamsburg	Raine Lumber & Coal..	4125'B
16	Hunter Moore No. 1.....	Irish Corner	Homer Hoke, et al.....	1780'B
17	Mary E. Morris Hrs. No. 2.....	Irish Corner	Homer Hoke, et al.....	1895'B
18	A. W. Smith No. 3.....	Irish Corner	Homer Hoke, et al.....	1705'B
19	Harry Ellis No. 4.....	(Monroe Co.)	Homer Hoke, et al.....	1865'B
25	Gauley Coal Land Co. No. 28.....	(Nicholas)	W. Va. Coal & Coke..	2530'B
26	Gauley Coal Land Co. No. 26.....	(Nicholas)	W. Va. Coal & Coke..	2125'B
27	Gauley Coal Land Co. No. 24.....	(Nicholas)	W. Va. Coal & Coke..	2600'B
27A	Gauley Coal Land Co. No. 21.....	(Nicholas)	W. Va. Coal & Coke..	2190'+
29	Gauley Coal Land Co. No. 23.....	(Nicholas)	W. Va. Coal & Coke..	2590'B
30	Gauley Coal Land Co. No. 23A.....	(Nicholas)	W. Va. Coal & Coke..	2580'B
31	Gauley Coal Land Co. No. 11.....	(Nicholas)	W. Va. Coal & Coke..	2315'B
32	Gauley Coal Land Co. No. 2.....	(Nicholas)	Gauley Coal Land.....	2325'B
33	Gauley Coal Land Co. No. 25.....	(Nicholas)	W. Va. Coal & Coke..	2045'B
37	Gauley Coal Land Co. No. 4.....	(Nicholas)	Gauley Coal Land.....	1660'B
38	Gauley Coal Land Co. No. 1.....	(Nicholas)	E. F. Saxman.....	1800'L
39	Gauley Coal Land Co. No. 13.....	(Nicholas)	W. Va. Coal & Coke..	2162'L
40	Gauley Coal Land Co. No. 12.....	(Nicholas)	W. Va. Coal & Coke..	2715'B
41	Gauley Coal Land Co. No. 19.....	(Nicholas)	W. Va. Coal & Coke..	2210'B
43	Gauley Coal Land Co. No. 3.....	(Nicholas)	Gauley Coal Land.....	2055'B
44	Gauley Coal Land Co. No. 27.....	(Nicholas)	W. Va. Coal & Coke..	2225'B
45	Gauley Coal Land Co. No. 16.....	(Nicholas)	W. Va. Coal & Coke..	2400'B
46	Gauley Coal Land Co. No. 14.....	(Nicholas)	Gauley Coal Land.....	2300'+
46A	Gauley Coal Land Co. No. 15.....	(Nicholas)	Gauley Coal Land.....	2600'+
46B	Gauley Coal Land Co. No. 8.....	(Nicholas)	Gauley Coal Land.....	2375'+
46C	Gauley Coal Land Co. No. 22A.....	(Nicholas)	Gauley Coal Land.....	2480'+
47	Gauley Coal Land Co. No. 29.....	(Nicholas)	W. Va. Coal & Coke..	2740'B
48	Gauley Coal Land Co. No. 17.....	(Nicholas)	W. Va. Coal & Coke..	2420'B
49	Gauley Coal Land Co. No. 18.....	(Nicholas)	W. Va. Coal & Coke..	2510'B
50	Gauley Coal Land Co. No. 6.....	(Nicholas)	Gauley Coal Land.....	2375'B

for Greenbrier County and Adjoining Area.

Sewell Coal.			Little Raleigh Coal.		Beckley Coal.		Fire Creek Coal.		No. 6 Pocahontas Coal.		No. 3 Pocahontas Coal.		Total Depth Feet.	No. on Map II.
Depth Base.	Thickness Inches.	Elevation Base.	Depth Base.	Thickness Inches.	Depth Base.	Thickness Inches.	Depth Base.	Thickness Inches.	Depth Base.	Thickness Inches.	Depth Base.	Thickness Inches.		
.....	364.1	29	368.0	1
.....	2
.....	3
.....	4
.....	41.9	28	169.5	18	216.5	29	302.0	5
.....	141.7	251.7	359?	553.0	6A
.....	105.5	191?	325.0	6B
.....	156.5	255.2	322?	498.5	584.3	687.0	6C
.....	116.1	168.3	175.0	6D
381.9	41	3450'	387.0	6E
.....	71.1	123.7	158.0	6F
.....	84.0	198.0	6G
.....	73.5	201.1	285?	200.0	6H
.....	43.7	250.5	330.0	6I
77.4	3560'L	80.0	6J
30.5	3567'L	100.4	310.5	324.5	6K
12.0	3483'L	215.0	6L
30.7	3367'L	322?	322?	470.0	6M
70.2	51	3551'L	235.3	40	332.0	12	603.3	28	607.0	6
83.0	41	3532'L	489.0	7
25.0	47	3600'L	432.0	8
94.4	33	3696'L	450.0	9
52.5	30	3788'L	460.0	10
53.3	38	3957'L	105.0	2	338.0	9	447?	6	642.9	10	758.5	839.0	11
.....	133.3	28	230.5	438?	1	465.5	12
.....	56?	2	263.9	42	384.9	1	541.0	13
.....	58.1	38	122.5	26	361.9	12	490?	3	577.0	14
.....	55.5	41	158?	28	271.0	15
.....	329.0	16
.....	223.0	17
.....	216.0	18
.....	156.0	19
61.2	15	2269'B	467.9	16	550.0	25
97.1	60	1928'B	200.0	26
07?	1	2293'B	476.0	27
33.0	41	2067'+	137.0	27A
.....	29
73.1	22	2407'B	177.0	30
34.3	43	2181'B	138.0	31
31.5	44	2194'B	285.0	32
46.7	40	1898'B	150.0	33
61?	1499'B	236.0	37
11.4	54	1689'L	640.0	38
58.8	25	2003'L	243?	332?	6	400.0	39
11.5	93	2103'B	615.5	40
.....	311.0	41
42.0	26	1613'B	445.5	43
.....	450.0	44
64.1	34	2016'B	392.0	45
99.1	36	2201'+	196.0	46
16.8	26	2183'+	419.0	46A
82.3	52	2293'+	108?	3	312?	6	362?	7	520?	5	665.0	46B
85.0	43	2295'+	289.0	46C
21.8	38	2518'B	227.0	47
37.4	38	2283'B	141.5	48
48.2	48	2362'B	151.0	49
.....	118.9	8	268?	23	384.0	50

Summarized Record of Coal Test Borings for

No. on Map H.	Name of Property.	Magisterial District.	Company.	Surface Elevation.
51	Gauley Coal Land Co. No. 7.....	(Nicholas)	Gauley Coal Land.....	2435'B
52	Gauley Coal Land Co. No. 5.....	(Nicholas)	Gauley Coal Land.....	2090'B
93	Brackens Cr. C. & L. Co. No. 2...	(Fayette)	Nuttall Hrs.	2357'L
93B	Brackens Cr. C. & L. Co. No. 4...	(Fayette)	Brackens Cr.	2124'L
93C	Brackens Cr. C. & L. Co. No. 3...	(Fayette)	Brackens Cr.	2345'L
93D	Brackens Cr. C. & L. Co. No. 1...	(Fayette)	Brackens Cr.	2431'L
94	Charles White	(Fayette)	Nuttall Hrs.	2400'B
95	Nuttall Hrs.	(Fayette)	Nuttall Hrs.	2655'B
96	Nuttall Hrs.	(Fayette)	Nuttall Hrs.	2545'B
97	Jno. Jordan-Amick	(Fayette)	Stancecock C. & C.....	2430'B
111	Beury No. 3.....	(Fayette)	N. R. & P. C. C.....	2545'L
112	N. R. & Poca. C. C. Co. No. 19...	(Fayette)	N. R. & P. C. C.....	2641'L
113	N. R. & Poca. C. C. Co. No. 20...	(Fayette)	N. R. & P. C. C.....	2595'L
114	N. R. & Poca. C. C. Co. No. 8NS.	(Fayette)	N. R. & P. C. C.....	2976'L
115	N. R. & Poca. C. C. Co. No. 22...	(Fayette)	N. R. & P. C. C.....	2801'L
116	N. R. & Poca. C. C. Co. No. 18...	(Fayette)	N. R. & P. C. C.....	2571'L
117	N. R. & Poca. C. C. Co. No. 2....	(Fayette)	N. R. & P. C. C.....	2863'L
118	N. R. & Poca. C. C. Co. No. 1....	(Fayette)	N. R. & P. C. C.....	2621'L
119	N. R. & Poca. C. C. Co. No. 5....	(Fayette)	N. R. & P. C. C.....	2554'L
120	N. R. & Poca. C. C. Co. No. 21...	(Fayette)	N. R. & P. C. C.....	2560'L
121	N. R. & Poca. C. C. Co. No. 17...	(Fayette)	N. R. & P. C. C.....	2739'L
122	N. R. & Poca. C. C. Co. No. 15...	(Fayette)	N. R. & P. C. C.....	2781'L
123	N. R. & Poca. C. C. Co. No. 14...	(Fayette)	N. R. & P. C. C.....	2584'L
124	N. R. & Poca. C. C. Co. No. 10...	(Fayette)	N. R. & P. C. C.....	2564'L
124A	N. R. & Poca. C. C. Co. No. 11...	(Fayette)	N. R. & P. C. C.....	2722'L
125	N. R. & Poca. C. C. Co. No. 4....	(Fayette)	N. R. & P. C. C.....	2865'L
126	N. R. & Poca. C. C. Co. No. 3....	(Fayette)	N. R. & P. C. C.....	2904'L
127	N. R. & Poca. C. C. Co. No. 23...	(Fayette)	N. R. & P. C. C.....	2978'L
128	N. R. & Poca. C. C. Co. No. 7NS.	(Fayette)	N. R. & P. C. C.....	3213'L
129	N. R. & Poca. C. C. Co. No. 12...	(Fayette)	N. R. & P. C. C.....	2872'L
130	N. R. & Poca. C. C. Co. No. 6NS.	(Fayette)	N. R. & P. C. C.....	3129'L
131	N. R. & Poca. C. C. Co. No. 5NS.	(Fayette)	N. R. & P. C. C.....	2804'L
132	N. R. & Poca. C. C. Co. No. 8....	(Fayette)	N. R. & P. C. C.....	2741'L
133	N. R. & Poca. C. C. Co. No. 9....	(Fayette)	N. R. & P. C. C.....	2719'L
134	N. R. & Poca. C. C. Co. No. 13...	(Fayette)	N. R. & P. C. C.....	2844'L
135	N. R. & Poca. C. C. Co. No. 7....	(Fayette)	N. R. & P. C. C.....	2774'L
136	N. R. & Poca. C. C. Co. No. 14...	(Fayette)	N. R. & P. C. C.....	2845'L
137	N. R. & Poca. C. C. Co. No. 4NS.	(Fayette)	N. R. & P. C. C.....	2914'L
138	N. R. & Poca. C. C. Co. No. 6....	(Fayette)	N. R. & P. C. C.....	2741'L
139	N. R. & Poca. C. C. Co. No. 3NS.	(Fayette)	N. R. & P. C. C.....	2895'L
140	N. R. & Poca. C. C. Co. No. 24...	(Fayette)	N. R. & P. C. C.....	2746'L
141	Bellwood Coal Co. No. 5.....	(Fayette)	Bellwood Coal	3285'L
142	Bellwood Coal Co. No. 4.....	(Fayette)	Bellwood Coal	3006'L
143	Bellwood Coal Co. No. 6.....	(Fayette)	Bellwood Coal	3155'L
144	Bellwood Coal Co. No. 1.....	(Fayette)	Bellwood Coal	3105'L
145	Bellwood Coal Co. No. 7.....	(Fayette)	Bellwood Coal	3155'L
146	Bellwood Coal Co. No. 9.....	(Fayette)	Bellwood Coal	3192'L
147	Bellwood Coal Co. No. 10.....	(Fayette)	Bellwood Coal	3159'L
148	Bellwood Coal Co. No. 8.....	(Fayette)	Bellwood Coal	3153'L
149	Bellwood Coal Co. No. 12.....	(Fayette)	Bellwood Coal	3197'L
150	Bellwood Coal Co. No. 3.....	(Fayette)	Bellwood Coal	3144'L
151	Bellwood Coal Co. No. 11.....	(Fayette)	Bellwood Coal	3196'L
152	Bellwood Coal Co. No. 2.....	(Fayette)	Bellwood Coal	3196'L

Greenbrier County and Adjoining Area—(Continued).

Sewell Coal.			Little Raleigh Coal.		Beckley Coal.		Fire Creek Coal.		No. 6 Pocahontas Coal.		No. 3 Pocahontas Coal.		Total Depth Feet.	No. on Map II.
Depth Base.	Thickness Inches.	Elevation Base.	Depth Base.	Thickness Inches.	Depth Base.	Thickness Inches.	Depth Base.	Thickness Inches.	Depth Base.	Thickness Inches.	Depth Base.	Thickness Inches.		
.....	201?	5	385.3	51
.....	56?	9	175.0	52
59.4	18	2198'L	310.8	7	365.0	93
44.0	29	1980'L	247.5	93B
27.5	42	2117'L	229.0	93C
82.7	13	2248'L	300.0	93D
41.0	46	2218'B	94
.....	95
.....	(600?)	96
.....	225.2	38	97
.....	111.3	5	300.0	111
.....	128.9	18	279.8	43	282.0	112
.....	141.3	12	305?	14	410.8	17	495.0	113
.....	348?	4	436.5	18	114
.....	27.4	12	123.5	3	273?	11	277.4	115
.....	45?	3	176.9	50	327.7	34	336.2	116
.....	109.0	4	184?	419.7	117
.....	154.5	6	221.5	5	402.9	22	513.1	118
.....	149.9	13	223.4	292.1	119
.....	54.4	7	215.0	120
.....	108.0	6	315.9	16	386.0	121
.....	78?	13	343.3	122
.....	24?	4	159.2	8	219.0	123
.....	146.9	26	150.0	124
.....	186?	8	298?	19	328.0	124A
.....	436.9	10	459.3	125
.....	38.4	4	291?	4	425.3	21	521.3	13	532.9	126
.....	219?	7	383?	8	473?	6	511.0	127
.....	412?	20	498?	6	128
.....	130?	1	398?	69	439.0	129
.....	106?	12	532?	2	130
.....	131
.....	45?	2	215.7	57	232.8	132
.....	56?	8	201.2	16	305.3	2	335.0	133
.....	313.9	9	410.4	4	421.2	134
.....	432.5	135
.....	136.9	56	141.6	136
.....	184?	2	137
.....	110.2	14	227.9	34	232.0	138
.....	128?	1	209.2	50	139
.....	111.9	20	202.5	17	205.0	140
.....	141
56.6	3228'L	327?	6	394?	564.7	47	583.6	142
.....	46?	45	148.3	18	200.0	143
.....	180.1	38	282.7	26	442.8	144
.....	142.3	98	148.0	145
.....	177.1	128	180.0	146
.....	208.0	115	208.5	147
.....	192.9	118	194.6	148
.....	202.0	88	206.8	149
.....	196.8	124	201.1	150
.....	189.3	57	145.5	151
.....	17.1	18	176.7	107	183.7	152

**DETAILED COAL TEST RECORDS,
MEADOW BLUFF DISTRICT.**

Of the 27 coal test borings that have been drilled in Meadow Bluff District, the complete records of 12 have been secured for publication. The records of 12 borings, drilled for the Leckie Smokeless Coal Company were available to the Survey and permission was granted for publication of the records exclusive of the coal sections. The remaining three holes were drilled on Little Sewell Mountain but their records could not be found.

Borings Nos. 1, 5E, 6, 7, 11, and 13 were included in Chapter V because of their stratigraphic importance.

The following record is of a hole drilled by the Margarette Coal Company and W. E. Deegans. No elevation for the hole is available, but what are believed to be the correct correlations for the various beds are indicated in the record:

**Gauley Coal Land Company Coal Test Boring No. 1—No. 5 on
Map II.**

Meadow Bluff District; on south side of Meadow Creek, 0.4 mile northeast of Marfrance; or 0.2 mile northwest of location shown on map; started, Dec. 6, 1929; completed, Jan. 7, 1930.

	Thickness.		Total.	
	Ft.	In.	Ft.	In.
Pottsville Series (302'±)				
Surface	3	0	3	0
Sandstone	0	6	3	6
Shale, sandy	5	6	9	0
Shale, gray	17	0	26	0
Shale, dark, sandy.....	13	6	39	6
Coal, bony, Little Raleigh?.....	2	4	41	10
Fire clay, sandy.....	5	2	47	0
Shale, dark	6	0	53	0
Sandstone	9	0	Lower Raleigh?	0
Sandstone, hard	33	0		
Sandstone, mixed with coal streaks, hard.....	4	0		
Sandstone, hard	1	0		
Shale, light, sandy.....	4	6	104	6
Sandstone, hard.....	13	0	117	6
Shale, dark, sandy.....	50	6	168	0
Coal, Beckley?.....	1	6	169	6
Fire clay	3	6	173	0
Slate, blue.....	5	6	178	6
Sandstone	7	4	185	10
Slate, blue	6	2	192	0
Sandstone, hard.....	7	8	199	8

		Thickness. Total.	
		Fcet.	Fcet.
Shale, red.....		25	465
Shale, greenish-gray.....		20	485
Shale, red.....		4	489
Shale, gray, sandy.....		79	568
Shale, dark, carbonaceous, limy.....		10	578
Shale, gray, sandy.....		5	583
Shale, dark, slaty, pyritic.....		2	585
Shale, gray, sandy.....		5	590
Sandstone, white, coarse-grained.....		1	591
Shale, dark-gray.....		6	597
Sandstone, gray (gns, 649'=82,000 cu. ft.).....		73	670
Shale, black.....		1	671
Sandstone, white, quartzitic.....19'	} Droop Sandstone	50	721
Sandstone, white, intermingled with black shale.....31			
Shale, dark-gray.....		24	745
Shale, grayish-black, sandy.....		73	818
Shale, dark-gray, limy.....		24	842
Shale, dark-blue, limy, with calcite crystals.....		28	870
Shale, gray, soft, sandy.....		17	887
Limestone, Reynolds, bluish-gray, soft, sandy.....		37	924
Sandstone, Webster Springs, gray, compact, carrying some muscovite.....		16	940
Shale, dark-blue, limy, with calcite crystals.....		87	1027
Shale, soft.....		4	1031
Limestone, gray, sandy.....9'	} Glenray Limestone.....	20	1051
Sandstone, gray, impure.....5			
Limestone, bluish, soft, impure.....6			
Shale, gray, limy.....		56	1107
Shale, blue, platy, slightly limy.....123'	} Lillydale Shale....	173	1280
Shale, dark-blue, soft, platy.....50 { (Penell Cave)			

Greenbrier Series (1480')

Shale, bluish-gray, limy.....25	} Alderson Limestone	136½	1440
Shale, dark, bluish-gray, slightly limy			
Limestone, dark-gray, soft.....86½			

NOTE: 1339 on sand line=1362½ on steel measuring line. All measurements corrected from here on.
Error on sand line probably cumulative.

Shale, Greenville, dark-blue, gray, slightly limy.....	60	1500
Limestone, blue-black, composed of rounded dark granules in a light- colored ground-mass, (oolitic texture)	80'	} Union Limestone.....
Limestone, bluish-gray, platy, shaly	110	
Limestone, gray.....	95	
	285	1785

Thickness. Totnl.
Feet. Feet.

Limestone, dark, bluish-gray, im- pure	90'		
Limestone, bluish-black, oolitic.....	112		
Limestone, gray, compact, probably somewhat shaly.....	43	Pickaway	
Shale, gray, limy, and sandy.....	30	Limestone	477 2262
Limestone, gray, shaly.....	40		
Limestone, bluish-gray, dark, im- pure.....	45		
Limestone, bluish-gray.....	55		
Limestone, bluish-gray.....	62		
Limestone, gray, siliceous; traces of brachiopods	28'	Taggard	
Limestone, gray, shaly.....	30	Limestone	58 2320
Limestone, dark-gray, hard.....	47'		
Sandstone, gray, fine-grained, some- what limy.....	23		
Limestone, gray, shaly.....	28	Patton	
Sandstone, bluish-gray, fine-grained, shaly, somewhat limy.....	67	Limestone	190 2510
Limestone, gray, hard, oolitic; ex- terior composed of concentric cal- cite layers.....	25		
Limestone, dark-blue, almost black, impure	80'		
Limestone, gray, hard, sandy.....	16		
Shale, bluish-gray, limy.....	10		
Shale, dark, bluish-gray, limy and sandy	9	Sinks Grove	
Limestone, dark-gray and light-gray, compact	29	Limestone	177 2687
Shale, slate-gray, sandy and some- what limy.....	8		
Limestone, dark-gray, impure.....	25		
Limestone, gray, very sandy (should probably be called a shaly, limy sandstone)	7'		
Sandstone, dark-gray, fine-grained, shaly and limy.....	8	Hillsdale	
Limestone, gray, shaly.....	36	Limestone	73 2760
Shale, gray, and black, limy shale; fragments composed of quartz in minute grains with conchoidal sur- faces	5		
Limestone, dark-gray, sandy.....	7		
Limestone, grayish-white, sandy.....	10		
Maccrady Series (180')			
Shale, gray; with quartz as above in ¼" lenses.....	28		2788
Quartz, chiefly, with some gray shale (quartz in co- mmun grains larger than above with conchoidal surfaces)	22		2810
Note: 2796 on sand line = 2806½ on steel measuring line, probably cumulative error. Corrected measure- ment from here on.			

Thickness. Total.
Feet. Feet.

Shale, purple-red with $\frac{1}{4}$ " to $\frac{1}{2}$ " quartz lenses (little gray shale)	30	2840
Sandstone, gray, fine-grained, shaly; or sandy shale....	15	2855
Shale structures, gray, hard, which resemble slickensides; pyrite disseminated throughout.....	45	2900
Shale, purple-red (little gray shale).....	40	2940

Pocono Series (410')

Sandstone, well cemented, medium-grained (quartz, with a little muscovite)	8'	Squaw Sand?..	74	3014
Sandstone, dark-gray, fine-grained (containing a little calcite).....	38			
Shale, hard, gray (with pyrite disseminated throughout); probably contains some quartz grains; slickensided surfaces.....	15			
Shale, gray, sandy; and sandstone, gray, coarse-grained, with conspicuous muscovite.....	13			
Shale, dark-gray, limy (with some gray, fine-grained sandstone, some pyrite, and little slickensided coal)	3			3017
Shale, carbonaceous; and coal, hard, slickensided; horizon of Merrimac Coal?.....	5			3022
Sandstone, grayish-white, coarse-grained	38'	Broad Ford Sandstone	226	3248
Sandstone, gray, coarse-grained.....	24			
Sandstone, gray, compact, with muscovite	19			
Shale, gray and reddish, with scattered grains of quartz; and sandstone, gray, with muscovite, a pebbly conglomerate (gas at 3105')....	12			
Sandstone, gray, with muscovite and little veined dolomite.....	42	Berea Sandstone	90	3350
Sandstone, chiefly, gray, compact, with muscovite; some slickensided, dark shale with a few dolomite veins.....	23			
Sandstone, bluish-gray; and sandy shale	68			
Shale, Sunbury? (Coffee Shale), bluish-gray, sandy.....	12			
Sandstone, gray, fine-grained; and dark bluish-gray shale.....	17'	Berea Sandstone	90	3350
Sandstone, light-gray, shaly.....	3			
Sandstone, chiefly, dark-gray, fine-grained; with some shaly sandstone	35			
Some sandstone, gray, fine-grained but chiefly bluish shale.....	35			

Chemung Series (172'+)

Sandstone, bluish-gray, fine-grained; with braebio-pod fragments.....	5	3355
Sandstone, chiefly, gray and fine-grained, or bluish-gray	44	3399

	Thickness.	Total.
	Feet.	Feet.
Sandstone, gray, fine-grained; and dark-colored sandy shale.....	26	3425
Sandstone, chiefly, reddish, fine-grained and platy, but some dark-colored and platy.....	10	3435
Sandstone, chiefly, gray and fine-grained; some gray shale with scattered quartz grains.....	15	3450
Shale, blue-black, platy.....	14	3464
Sandstone, grayish-white.....	27	3491
Shale, blue-black.....	2	3493
Note: 3493' on sand line=3496½ on steel measuring line		3496½
Sandstone, light-gray, well cemented.....	5½	3502
Shale, slate-blue, chiefly, with 5% of light-gray well cemented sandstone which appears to occur as small lenses in the shale; little pyrite.....	20	3522
Total depth.....		3522

SUMMARY OF OIL AND GAS POSSIBILITIES IN GREENBRIER COUNTY.

Is it worth while to prospect for oil or gas in Greenbrier County? The answer to this question, for that portion of the County east of the Greenbrier River, is no. To answer this question for the western part of the County is not so easy. The answer largely depends upon two things; namely, source beds and the degree of metamorphism necessary to destroy or dissipate oil and gas. These factors have been discussed on an earlier page of this Chapter, where it was pointed out that source beds are probably present and while the chances of finding oil are very small, there is some chance of finding commercial quantities of natural gas.

From the standpoint of the petroleum geology of West Virginia as a whole, Greenbrier County is considerably farther east than any commercial oil or gas pool thus far discovered. While this fact does not necessarily condemn the territory, it does suggest that the search for gas in this county should be left to those that can afford to lose.

CHAPTER XI.

COMMERCIAL COAL.

INTRODUCTION.

In Chapter VI a systematic description of all the coal seams found in Greenbrier County has been given, together with their correlations. Many of the beds are too thin, lenticular, or impure to be of commercial rank and all such have been fully described in the Chapter named. In the present Chapter numerous measured sections for those coals that are of minable thickness and purity, with estimates of their probable tonnage, and etchings showing their areal extent are given.

In this county there appear to be six coals that have minable thickness and 24 others that are too thin, impure, or irregular to be of more than local value, some of these latter being thin beds that are of scientific interest only. The minable seams in descending order are the Sewell, Little Raleigh, Beekley, and Fire Creek of the New River Group; and the No. 6 Pocahontas and No. 3 Pocahontas of the Pocahontas Group; all in the Pottsville Series.

Figure 16 shows the different coal seams of the county, giving not only their thickness but also the average interval (base to base) between them. Figures 17, 19, 20, 21, and 23 show approximately where the commercial seams occur in possible minable thickness in the county.

In general, these coals are semibituminous, those northeast of Beech Ridge being on the dividing line in chemical composition between semibituminous and bituminous and those in the southwest part of the county approach the semianthracite classification.




FIGURE 16
 DIAGRAM SHOWING THE POSITION AND THICKNESS
 OF COAL SEAMS IN GREENBRIER COUNTY

The coals are variously used for steam and domestic fuel, for metallurgical purposes and for mixing with higher volatile coals to produce gas and by-product coke. Owing to their low ash and sulphur content, their low volatile content and the ease with which they may be crushed, the coals of Greenbrier County would be especially well adapted for use in mechanical stokers or for powdered fuel.

STATISTICS OF COAL PRODUCTION.

Commercial coal mining has been practiced in Greenbrier County for many years, the first record of production being in 1907; the record of mining being continuous from that year to date.

So far as the records of the Department of Mines show, the Lost Flat Mine (Mine No. 308) of the Elk Lick Coal Company, was the first commercial mine in the county. This mine is in what appears to be the Beekley seam and was opened in 1906 or 1907. In 1910 the same company opened the Spruce Knob Mine (Mine No. 225) on North Fork of Cherry River in the Sewell seam, abandoning the Lost Flat Mine the same year. In 1916 J. W. Dwyer opened the Dwyer Mine (Mine No. 424, now Tuck Brothers) near the Fayette County line in the No. 6 Pocahontas seam. It was not until 1922 that Greenbrier County came to the front as one of the major coal-producing counties in the State. In this year several mines were opened in the Sewell seam along Meadow Creek.

At the present time about 95 per cent. of the coal production in Greenbrier County comes from the Sewell seam, but there is a large reserve of excellent coal in the lower seams. If prospecting with good results is any guide, there should be several mines opened in these lower seams in the near future.

The following tables, mainly assembled from statistics given in the Annual Reports of the West Virginia Department of Mines, supplemented by certain unpublished data from N. P. Rhinehart, present Chief, gives the coal production of the county since 1907, the relative rank in production as compared to other counties and the production of coal by mines:

Greenbrier County Coal Production.

(Production by fiscal years ending June 30 of each year up to June 30, 1924; production by calendar years starting January 1, 1925).

Year.	Long Tons (2240 lbs.)	=Short Tons. (2000 lbs.)	Order.
1907	31,978	35,815	28
1908	36,066	40,394	28
1909	32,296	36,171	29
1910	21,688	24,290	32
1911	48,819	54,677	30
1912	52,358	58,641	27
1913	42,853	47,995	32
1914	22,633	25,349	33
1915	24,128	27,023	32
1916	35,692	39,975	32
1917	45,207	50,632	32
1918	37,311	41,788	32
1919	33,695	37,738	32
1920	52,398	58,686	33
1921	52,153	58,411	33
1922	400,938	449,050	21
1923	431,643	483,440	24
1924	826,185	925,327	18
1924(a)	503,383	563,789	18
1925(b)	1,181,016	1,322,738	18
1926(b)	1,278,688	1,432,131	19
1927(b)	1,428,716	1,600,162	18
1928(b)	1,480,544	1,658,209	18
1929(b)	1,598,240	1,790,029	18
1930(b)	1,814,716	2,032,482	14
1931(b)	1,621,671	1,816,272	13
1932(b)	1,199,347	1,343,269	14
1933(b)	1,464,357	1,640,080	14
1934(b)	1,559,847	1,747,029	13
1935(b)	1,421,104	1,591,636	14
1936(b)	1,598,224	1,790,011	14
Totals	20,377,894	22,823,239	

(a) Last six months of 1924.

Coal Tonnage Production in Greenbrier County by the Various Mines for the Year Ending June 30, 1907.

Name of Company	Name of Mine	Production of Coal. (Tons of 2240 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Liek Coal Co.....	Lost Flat.....	31,978	373	31,605
For the Year Ending June 30, 1908.					
Liek Coal Co.....	Lost Flat.....	36,066	711	35,355
For the Year Ending June 30, 1909.					
Liek Coal Co.....	Lost Flat.....	32,296	774	31,522
For the Year Ending June 30, 1910.					
Liek Coal Co.....	Spruce Knob.....	13,372	134	93	13,145
Liek Coal Co.....	Lost Flat.....	8,316	180	112	7,724
Totals.....		21,688	614	205	20,869

For the Year Ending June 30, 1911.

Name of Company	Name of Mine	Production of Coal. (Tons of 2240 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Elk Llek Coal Co.....	Spruce Knob.....	48,819	281	134	48,401

For the Year Ending June 30, 1912.

Elk Llek Coal Co.....	Spruce Knob.....	52,358	285	158	51,915
-----------------------	------------------	--------	-----	-----	--------

For the Year Ending June 30, 1913.

Elk Llek Coal Co.....	Spruce Knob.....	42,853	268	123	42,462
-----------------------	------------------	--------	-----	-----	--------

For the Year Ending June 30, 1914.

Elk Llek Coal Co.....	Spruce Knob.....	22,633	295	114	22,224
-----------------------	------------------	--------	-----	-----	--------

For the Year Ending June 30, 1915.

Elk Llek Coal Co.....	Spruce Knob.....	24,128	352	104	23,672
-----------------------	------------------	--------	-----	-----	--------

For the Year Ending June 30, 1916.

Name of Company	Name of Mine	Production of Coal. (Tons of 2240 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Dwyer, J. W.	Dwyer	10,058	10,058
Elk Lick Coal Co.	Spruce Knob	25,634	359	121	25,154
Totals.....		35,692	359	121	35,212

For the Year Ending June 30, 1917.

Elk Lick Coal Co.	Spruce Knob	27,038	468	88	26,482
Meadow River Smokeless Coal Co.	Dwyer	18,169	320	17,849
Totals.....		45,207	468	408	44,331

For the Year Ending June 30, 1918.

Elk Lick Coal Co.	Spruce Knob	19,401	611	71	18,719
Meadow River Smokeless Coal Co.	Dwyer	17,910	84	17,826
Totals.....		37,311	611	155	36,545

For the Year Ending June 30, 1919.

Name of Company	Name of Mine	Production of Coal. (Tons of 2240 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Elk Lick Coal Co.....	Spruce Knob.....	22,629	560	103	21,966
Meadow River Smokeless Coal Co.....	Dwyer	11,066	2,771	8,295
Totals.....		33,695	560	2,874	30,261

For the Year Ending June 30, 1920.

Elk Lick Coal Co.....	Spruce Knob.....	29,207	776	169	28,262
Lincoln Smokeless Coal Co....	Lincoln No. 1.....	3,191	300	2,891
Meadow River Smokeless Coal Co.....	Dwyer	20,000	240	19,760
Totals.....		52,398	776	709	50,913

For the Year Ending June 30, 1921.

Name of Company	Name of Mine	Production of Coal. (Tons of 2240 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Elk Lick Coal Co.....	Spruce Knob.....	32,893	1,135	252	31,506
Lincoln Smokeless Coal Co....	Lincoln No. 1.....	10,710	10,710
Meadow River Smokeless Coal Co.....	Dwyer	8,550	85	8,465
Totals.....	52,153	1,135	337	50,681

For the Year Ending June 30, 1922.

Elk Lick Coal Co.....	Spruce Knob.....	32,018	1,304	247	30,467
Frances Coal Co.....	Frances	7,680	7,680
Greenbrier Smokeless Coal Co.....	Greenbrier	37,535	191	37,344
Imperial Smokeless Coal Co....	Quinwood	79,178	79,178
McRoss Smokeless Coal Co....	Lincoln No. 1.....	4,500	1,100	3,400
Margarette Coal Co.....	Margarette Nos. 1 & 2.....	65,402	390	1,005	64,007
Meadow Creek Coal Co.....	Orlehton	52,411	400	52,011
Meadow River Smokeless Coal Co.....	Dwyer	5,890	1,500	4,390
Nelson Fuel Co.....	No. 1.....	44,817	400	44,417
Nelson Fuel Co.....	No. 2.....	71,507	270	71,237
Totals.....	400,938	1,694	5,113	394,131

For the Year Ending June 30, 1923.

Name of Company	Name of Mine	Production of Coal. (Tons of 2240 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Elk Llek Coal Co.....	Spruce Knob.....	37,896	1,526	248	36,122
Frances Coal Co.....	Frances	29,680	240	29,440
Greenbrier Smokeless Coal Co.....	Greenbrier	12,573	347	42,226
Imperial Smokeless Coal Co..	Quinwood	71,954	287	71,667
McRoss Smokeless Coal Co..	Lincoln No. 1.....	11,691	550	11,141
Margarette Coal Co.....	Margarette No. 1.....	35,398	964	34,434
Margarette Coal Co.....	Margarette No. 2.....	25,742	1,537	2,726	21,479
Meadow Creek Coal Co.....	Crichton	59,809	384	59,425
Meadow River Smokeless Coal Co.....	Dwyer	14,000	1,500	12,500
Nelson Fuel Co.....	Nelson No. 1.....	70,113	1,101	69,012
Nelson Fuel Co.....	Nelson No. 2.....	32,787	32,787
Totals.....	431,643	3,063	8,347	420,233

Name of Company	Name of Mine	Production of Coal. (Tons of 2000 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Elk Lick Coal Co.....	Spruce Knob.....	60,360	2,022	305	58,033
Frances Coal Co.....	Frances	65,607	300	65,307
Greenbrler Smokeless Coal Co.....	Greenbrler	94,968	277	94,691
Imperial Smokeless Coal Co.....	Quinwood	174,155	318	173,837
Margarette Coal Co.....	Margarette No. 1.....	91,504	1,200	600	89,704
Margarette Coal Co.....	Margarette No. 2.....	87,991	1,200	600	86,191
Meadow Creek Coal Co.....	Crichton	87,940	520	87,420
Midland Smokeless Coal Co.....	Midland No. 2.....	3,800	3,800
Nelson Fuel Co.....	Nelson Nos. 1 & 2.....	259,002	813	258,189
Totals.....	925,327	4,122	3,733	917,172

Coal Tonnage from July 1, 1924, to December 31, 1925.

Elk Lick Coal Co.....	Spruce Knob.....	64,448	2,782	459	61,207
Frances Coal Co.....	Frances	74,297	403	73,894
Greenbrler Smokeless Coal Co.....	Greenbrler	202,251	900	201,351
Imperial Smokeless Coal Co.....	Quinwood	352,149	25	576	351,548
Margarette Coal Co.....	Margarette	464,826	6,547	2,138	456,141
Meadow Creek Coal Co.....	Crichton	214,376	708	213,668
Midland Smokeless Coal Co.....	Midland	2,213	1,077	1,136
Nelson Fuel Co.....	Nelson Nos. 1 & 2.....	511,967	2,088	509,879
Totals.....	1,886,527	9,354	8,349	1,868,824

For the Calendar Year 1926.

Name of Company	Name of Mine	Production of Coal. (Tons of 2000 lbs.)	Distribution of Coal.		
			Used in Operation of Mine.	Furnished local trade and tenants.	Shipped from Mine.
Elk Llek Coal Co.....	Spruce Knob.....	43,259	1,953	342	40,964
Francees Coal Co.....	Francees	62,172	480	61,692
Greenbrier Smokeless Coal Co.....	Greenbrier	202,480	725	201,755
Imperial Smokeless Coal Co..	Quinwood	287,328	530	286,798
Margarette Coal Co.....	Margarette No. 2.....	290,044	7,373	1,845	280,826
Meadow Creek Coal Co.....	Crichton	168,783	576	168,207
Meadow River Fuel Co.....	Lincoln	1,645	300	1,345
Midland Smokeless Coal Co..	Midland	3,000	1,800	1,200
Nelson Fuel Co.....	Nelson No. 1.....	362,741	2,230	360,511
Nelson Fuel Co.....	Nelson No. 2.....	10,679	10,679
Totals.....		1,432,131	9,326	8,828	1,413,977

The West Virginia Geological Survey has long classified the Red Medina as Silurian and there is considerable evidence to substantiate such a classification*. The Pennsylvania and New York Geological Surveys also classify the Juniata and Queenston as Silurian but the Richmond beds are classified (by some) as Ordovician. On the basis of the supposed equivalence of the Red Medina with the Richmond, the U. S. Geological Survey places this series in the Ordovician.

ECONOMIC ASPECTS, RED MEDINA SERIES.

From an economic standpoint, the Red Medina Series in this area is of minor importance, the shales being generally too sandy for brick, while the sandstones are not suitable for building stone.

*See Mercer, Monroe, and Summers, Pocahontas, Pendleton, Mineral and Grant, and the Hampshire and Hardy County Reports of the West Virginia Geological Survey.

PART III.

Mineral Resources.

CHAPTER X.

PETROLEUM AND NATURAL GAS.

GENERAL STATEMENT.

In evaluating the chances of finding commercial deposits of petroleum or hydrocarbon natural gas in any area, certain fundamental factors must be taken into account. (1) There must be source beds from which the hydrocarbons may be derived. (2) There must be reservoir beds, in which the hydrocarbons can collect, that will yield these substances in commercial amounts. (3) The source beds and reservoir beds must be near enough to one another that the oil and gas can migrate from the former to the latter. (4) There must be a suitable structure or trap to permit segregation of gas, oil, and water. (5) The degree of metamorphism of the beds must not be too great. All of the above factors must be taken into consideration by the petroleum geologist in the search for new oil and gas pools and each will be considered in turn as to the manner in which it affects Greenbrier County.

(1) Practically all petroleum geologists are now agreed that petroleum and the associated natural gas have been derived from organic matter of vegetable or/and animal origin. Therefore, oil and gas deposits can only be found in regions where source beds contain a sufficient amount of organic matter that is suitable for the formation of these hydrocarbons. There

is a great deal that is not known about what constitutes source beds of petroleum but for the present it may be assumed that adequate source beds are present in Greenbrier County.

(2) The exposed rocks of Greenbrier County show a number of beds that appear to be suitable as reservoir rocks for petroleum and natural gas.

(3) The distance that oil and gas may migrate has been the subject of much debate. The writers are inclined to the view that in most cases the source beds and reservoir beds must be in actual contact at some point but it is conceded that the oil and gas may migrate considerable distances laterally along the beds. It is probable that there are several areas in Greenbrier County that meet this requirement.

(4) There are several structures in Greenbrier County that appear to be suitable for the accumulation of oil and gas. In addition to anticlinal structures, oil and gas are often trapped in sand lenses that are sealed updip by pinching out of the permeable rock. Judging from the outcropping rocks such traps may be expected in Greenbrier County.

(5) Commercial oil and gas deposits are never found in highly metamorphosed rocks. In nature all gradations between unmetamorphosed sedimentary rocks and their metamorphosed equivalents are found. Since commercial deposits of oil are found in the former and never in the latter it is apparent that somewhere in between there must be a zone where the degree of metamorphism has been sufficient to destroy or dissipate any oil or gas that may have been present. White has pointed out that the zone between 62 and 65 per cent. fixed carbon ratio in coal may be considered the extinction zone for the occurrence of commercial deposits of oil but natural gas may be found in areas that have suffered more advanced metamorphism¹. As shown in the Table of Coal Analyses published at the end of Chapter XI, the coals of Greenbrier County all show more than 70 per cent. fixed carbon when calculated on a moisture and ash free basis. This would seem to eliminate Greenbrier County as prospective

¹White, David, *Metamorphism of Organic Sediments and Derived Oils*; Bull. Amer. Assoc. of Petroleum Geologists, Vol. 19, p. 592; 1935.

oil territory. A fact that is difficult to explain under David White's theory is that the percentage of fixed carbon in the coal of individual coal beds increases from east to west in Greenbrier County.

There is one other factor that should be given careful consideration. That factor is the distribution of oil and gas in the State as a whole. The oil pools nearest Greenbrier County are about 30 miles to the northwest, in Clay and Kanawha Counties. Gas has been produced somewhat nearer the boundaries of Greenbrier County, the commercial production in Fayette and Nicholas Counties being only 16 to 20 miles west or northwest of the county line. What is much more encouraging is the fact that gas has been found near Bozoo (Chestnut Hill) in Monroe and Summers Counties where the rocks are more severely metamorphosed than are the rocks in western Greenbrier County. Wells drilled much nearer Greenbrier County in Nicholas, Fayette, Summers, and Monroe Counties have found shows of natural gas but no oil.

From the foregoing discussion it is seen that the chances of finding oil in Greenbrier County are very slight but there does seem to be some chance of finding natural gas.

PROSPECTIVE OIL AND GAS AREAS.

There are four areas in western Greenbrier County in which closures probably exist on subsurface beds. (1) Judging from surface exposures there is a small closure on the south end of the Webster Springs Anticline. As shown by the green contours on Map II, this "dome" is located about 3 miles north of Anjean and 2 miles west of Dno. It is possible that this "dome" is tilted enough with depth, due to the convergence of the Mississippian beds, to eliminate the northeast closure on all beds below the base of the Greenbrier Limestone.

(2) About one mile south of Mann Knob in Williamsburg District there is another closure on the same anticline. The closure is not readily apparent from the green contours but here the convergence of the underlying beds will greatly increase the closure on each successively lower horizon.

(3) In the vicinity of Cold Knob it is reasonably certain that there is a structural closure on the subsurface horizons.

Structure contours on the Sewell Coal horizon fail to show a closure but a small closure is present on the Princeton Sandstone. Here, too, the convergence of the underlying beds will increase the closure on each successively lower horizon.

(4) From a structural standpoint, the most favorable area for gas production in Greenbrier County is on Brushy Ridge. This topographic feature coincides with the Williamsburg Anticline and the surface rocks show a closure, in all directions, of at least 1,000 feet. The cross-section of this structure is clearly shown on Cross-Section D—D' printed on the margin of Map II. The narrow crest and steep sides of this structure may make the drilling of a straight hole difficult.

The area east of the Greenbrier River can be eliminated as prospective oil and gas territory because all of the horizons known to be productive in West Virginia either outcrop at the surface or have been removed from the area by erosion.

PROSPECTIVE OIL AND GAS HORIZONS.

In the area under discussion the known productive sands of the Monongahela, Conemaugh, Allegheny, and upper Pottsville Series do not now exist, as they belong above the youngest remaining formations. Small areas of rocks of the middle and lower Pottsville Series remain but in their present position they may be definitely eliminated from prospective oil and gas horizons. In the Mauch Chunk Series the upper and middle portions outcrop widely and offer little hope of oil or gas. In the lower portion occur the Droop and Webster Springs Sandstones that offer slight possibilities of production in areas 1 and 2 described above.

The Greenbrier Series, or Big Lime of the drillers, probably has a thickness of 400 feet in western Greenbrier County. At the outcrop, this series contains several oolitic layers, as described in Chapter VII. The oolitic beds might serve as reservoir beds and production is possible from these horizons in areas 1, 2, and 3 as outlined above. In a well drilled near Lookout, Fayette County, a good show of gas and several separate salt water horizons were found in this series. (See record of well No. 6 published on a succeeding page of this Chapter.)

The Pocono Series, which occurs just below a protective mantle of red shales (Maecrady), contains several coarse sandstone beds interbedded with shales, that would appear to be good reservoir beds. It is from sands of this horizon that a considerable quantity of gas was found in the Wills and Johnson wells of Monroe County, and the Shumate well of Summers County. The records of these wells are published on subsequent pages of this Chapter. It is possible that production will be found at this horizon. These horizons outcrop at the surface or are very near the surface in area number 4.

There are several sandstones in the Upper Devonian that appear to be suitable for reservoir beds. It is possible that production will be secured from these horizons.

The Middle Devonian is mainly composed of black shale in Greenbrier County. Devonian beds of a similar character are producing commercial quantities of gas in southwestern West Virginia.

The Oriskany Series of the Lower Devonian has recently become a valuable gas producing horizon in the West Virginia area. The Ridgeley Sandstone is producing great volumes of gas in Kanawha County and the Huntersville Chert is producing gas in Fayette County, Pennsylvania, near this State's northern boundary. It is the Ridgeley Sandstone that appears to be the most promising horizon for production in Greenbrier County.

The Helderberg Series coming next below the Oriskany contains two sandstones (Healing Springs and Clifton Forge) that appear to be suitable as reservoir beds. Any well drilled to the Oriskany in the Greenbrier area should not be abandoned without testing these horizons.

The Silurian rocks in Greenbrier County outcrop along the Browns Mountain Anticline in a broken and greatly mashed condition so that the evaluation of the various beds as prospective horizons for gas (or oil) is very difficult. In western Greenbrier County the Keefer and White Medina Sandstones may be porous enough to serve as reservoirs for fluids.

Table Showing the Estimated Depths to Geologic Horizons at Various Points in Greenbrier County.

Geologic Horizon	3 Mi. N. of Anjean.	1. Mi. S. of Mann Knob.	Cold Knob.	Alta.	Russellville.	County Line. N. F. Cherry River.
Greenbrier Series (Top).....	2,300	1,450	1,450	1,550	1,300
Greenbrier Series (Base).....	2,725	1,875	1,925	1,930	1,700
Pocono (Base).....	3,050	2,175	2,200	300	2,400	2,000
Oriskany Series (Top).....	8,250	7,650	7,800	5,800	6,750	7,600
Helderberg (Base).....	8,650	8,050	8,200	6,200	7,150	8,000
White Medinn (Top).....	9,950	9,350	9,450	7,450	8,500	9,400

In the above table the wells at the localities noted are assumed to start at the following horizons:

Well 3 miles north of Anjean, Sewell Coal.

Well 1 mile south of Mann Knob, top of Manah Chnuk.

Well near Cold Knob, top of Princeton Sandstone.

Well near Alta, 100 feet below the top of Pocono.

Well near Russellville, 150 feet below Sewell Coal.

Well near the county line on North Fork of Cherry River, top of Princeton Sandstone.

Several samples of the Lower Devonian and Upper Silurian sandstones were collected by the writers in connection with the preparation of this report. These samples were collected for comparison with drill cuttings from the same horizons taken from wells in various parts of the State. The samples were examined by Professor Martens and those familiar with the work of Dr. Martens will be interested in the following table of mineral identifications:

Mineralogy of Lower Devonian and Upper Silurian Sandstones
in
Greenbrier and Pocahontas Counties, W. Va.
(Mineral Identifications by J. H. C. Martens).

Alvon Section.

Sample No.	HEAVY								LIGHT				Formation	
	Pyrite	Zircon	Rutile	Leucosene	Brown Tourmaline	Green Tourmaline	Blue Tourmaline	Anthogenic Tourmaline	Monazite	Quartz	Anthogenic Quartz	Feldspar		Glauconite
1M.....	C	VS	VA	C	S	VA	A	Huntersville
2M.....	C	S	VA	C	C	VS	VA	C	S	Ridgeley
3M.....	A	S	A	C	S	S	VA	VS	Ridgeley
4M.....	VA	S	C	C	C	VS	VS	VA	Ridgeley
5M.....	A	VS	C	A	C	VS	C	VS	VA	Healing Springs
6M.....	A	S	C	A	S	S	A	VA	Healing Springs
7M.....	A	S	A	C	S	S	S	VS	VA	Healing Springs
8M.....	A	A	A	S	S	C	VA	Clifton Forge
9M.....	C	C	VA	S	S	C	VA	Clifton Forge
10M.....	S	C	A	S	C	VA	C	Clifton Forge
11M.....	S	C	VA	S	C	VA	Keeler
12M.....	VA	VS	C	A	S	S	A	VA	Keeler
13M.....	C	C	A	C	S	C	VA	C	Keeler
14M.....	C	VS	A	A	S	S	VA	A	Keeler
15M.....	C	C	S	A	A	C	S	C	VA	A	Keeler
16M.....	C	C	VA	S	S	A	VA	C	Keeler
17M.....	C	C	VA	S	S	A	VA	A	Keeler
18M.....	C	C	C	VA	S	S	A	VA	C	Keeler
19M.....	C	A	C	A	C	S	C	VA	C	Keeler
20M.....	VA	A	VA	A	Keeler
21M.....	S	C	C	VA	C	S	A	VA	A	Keeler

Burr Valley Section.

1M.....	S	A	S	A	C	S	A	C	A	Huntersville
2M.....	VA	C	C	VS	VA	S	Ridgeley
3M.....	A	S	A	C	S	S	C	VA	A	C	Ridgeley
4M.....	A	S	A	C	S	S	C	VA	C	Ridgeley (?)
5M.....	A	S	A	C	S	S	C	VA	C	S	Ridgeley (?)

Bobs Ridge Section.

1M.....	A	A	C	S	C	VA	A	Healing Springs
2M.....	A	S	A	C	S	C	VA	A	Healing Springs
3M.....	A	A	C	S	C	VA	A	Healing Springs
4M.....	A	S	A	C	S	C	VA	A	Healing Springs
5M.....	C	S	A	A	C	S	A	VA	A	Healing Springs
6M.....	A	S	A	A	C	A	VA	A	Healing Springs
7M.....	A	S	A	A	C	S	A	VA	A	Healing Springs
8M.....	A	S	A	A	C	A	VA	A	Healing Springs

Key.
VA—Very abundant. C—Common VS—Very scarce.
A—Abundant. S—Scarce.

The amounts of heavy minerals in the table are relative to the total heavy fraction and not to the rock as a whole, since the amount of heavy minerals in each of the samples is very small.

WELL RECORDS.

In Greenbrier County five wells have been drilled in search of oil or gas, none of which obtained production. Three shallow wells were drilled north of Sam Black Church, one of which may have reached the Greenbrier Limestone (Big Lime). The records of these wells (Nos. 1, 2, and 3 on Map II) could not be found, but it is reported that the deepest well (No. 2) reached a depth of 1,600 feet. No. 1 reached a reported depth of 800 feet but the total depth of No. 3 is unknown.

The record of the **S. W. Hinkle well (No. 4 on Map II)**, drilled about one mile north of Trout P. O., reached a total depth of 1,600 feet. This well penetrated 400 feet of the Chemung beds. The well is unfavorably located from a structural standpoint as it is in the syncline between the "dome" at Cold Knob and the Williamsburg Anticline. In 1936 the drilling machine was still setting at the location of this well and an artesian flow of fresh water was emerging between the 8- and 10-inch casings. The source of the water is probably the Droop or Webster Springs Sandstone. The record of this well is published in Chapter V, pages 180-1, in connection with the Cold Knob—Hinkle Well Section.

So far as known to the writers, only one other test well for oil or gas has been drilled in Greenbrier County. This well (**No. 5 on Map II**) was located on the east side of the Greenbrier River, 1.5 miles northeast of Anthony. Located just east of the axis of the Caldwell Syncline, the well was drilled at a structurally unfavorable position. The formations shown in the log of the well outcrop along Anthony Creek a short distance to the east of the well. The well log is as follows:

S. M. Jones Well—No. 5 on Map II.

Falling Springs District; on east side of Greenbrier River, 1.5 miles northeast of Anthony and 0.38 mile southeast of mouth of Laurel Run; drilled in 1913 by South Penn Oil Co.; elevation, 1810-15' (by topo. contour).

	Thickness. Total.	
	Feet.	Feet.
Wooden conductor.....	9	9
Slate	11	20
Lime, (Berea).....	30	50

	Thickness. Feet.	Total. Feet.
Red rock (Top of Catskill).....	2	52
Red sand.....	48	100
White slate.....	5	105
Red sand.....	45	150
Sandy lime.....	70	220
White slate.....	5	225
Sandy lime.....	15	240
Slate and shells.....	15	255
Sandy lime.....	20	275
Brook	2	277
Sandy lime.....	17	294
Slate	6	300
Lime	10	310
Red rock (Base of Catskill).....	15	325
Lime	15	340
Sandy lime (Hendricks).....	30	370
Black slate.....	8	378
Lime	6	384
Slate	3	387
Lime	13	400
Sandy lime.....	35	435
Slate	15	450
Lime	25	475
Slate	4	479
Lime	21	500
Sandy lime.....	60	560
White slate.....	15	575
Lime	25	600
Slate	25	625
Sandy lime.....	20	645
Slate and shells.....	35	680
Sandy lime.....	40	720
Slate	13	733
Slate and shells.....	17	750
Lime	40	790
Slate	10	800
Sandy lime.....	45	845
Slate	40	885
Lime	40	925
Slate and shells.....	75	1000
Lime	40	1040
Slate	20	1060
Lime	30	1090
Slate	20	1110
Lime	160	1270
Slate and shells.....	40	1310
Lime	15	1325
Slate	25	1350
Lime	50	1400
Slate and shells.....	40	1440
Slate	30	1470
Lime	80	1550
Sandy lime.....	50	1600
Lime	40	1640

	Thiekness.	Total.
	Feet.	Feet.
Sandy lime.....	50	1690
Lime	35	1725
Slate	145	1870
Black lime.....	50	1920
Slate	30	1950
Slate and shells.....	75	2025
Slate	35	2060
Slate and shells.....	40	2100
Sandy lime.....	20	2120
Slate	70	2190
Sandy lime.....	15	2205
Slate and shells.....	45	2250
Limo	50	2300
Slate	50	2350
Slate and shells.....	25	2375
Limo	25	2400
Sand	20	2420
Slate	15	2435
Limo	15	2450
Slate	15	2465
Lime	25	2490
Slate and shells.....	30	2520
Slate	20	2540
Total depth		2540
Total depth (steel-line measurement).....		2575

The following is the record of a well drilled near Lookout, Fayette County. This well is of particular interest in that it illustrates the possibility of several producing horizons within the Greenbrier Series. The well is located rather far down on the plunging end of the Mann Mountain Anticline, which perhaps accounts for the salt water. The fact that salt water was found proves the presence of reservoir beds in this series. The presence of an estimated flow of 300,000 cu. ft. of gas is also of interest. The carbon ratio of the Sewell Coal at Lookout is slightly higher than the carbon ratio of the Sewell Coal in Greenbrier County:

John Nuttall Estate No. 1 Well—No. 6 on Map II.

Fayette County, Nuttall District; authority, Joseph H. Holmes; on Keeney Creek, at Lookout, near mouth of Lookout Mine of Lookout Coal & Coke Company; completed, May 13, 1926, after one movement of rig; elevation, 2257' B. 13" casing. 36'; 10", 260' (pulled 240'); 8", 1370' (pulled); 6 $\frac{5}{8}$ ", 1830' (pulled 79 joints; 6 joints left in); 5 $\frac{3}{8}$ ", 2095' (pulled). Show of gas at 1900-1910', estimated 300,000 cu. ft. Gas drowned out by water but when casing was pulled the gas showed considerable pressure. Well plugged.

	Thickness. Feet.	Total. Feet.
Pottsville Series (825' +)		
Soil and clay.....	47	47
Lime shell.....	3	50
Coal, Sewell (2210' B., in mine).....	3	53
Slate	67	120
Lime, sandy, very hard.....	4	204
Slate	43	247
Sand and lime.....	63	310
Slate	35	345
Lime and slate.....	113	458
Coal, Fire Creek.....	7	465
Slate	10	475
Fire clay	6	481
Lime and slate.....	344	825
Mauch Chunk Series (990')		
Red rock.....	20	945
Lime, black.....	30	975
Slate, white.....	10	985
Red rock.....	185	1170
Lime	3	1173
Red rock.....	102	1275
Lime shell.....	15	1290
Slate	12	1302
Lime, sandy.....	8	1310
Slate	2	1312
Lime, grlty.....	2	1314
Slate	2	1316
Lime	8	1324
Slate	16	1340
Lime shell.....	5	1345
Slate	13	1358
Lime shell.....	2	1360
Sand (water, 1393').....	35	1395
Red rock.....	10	1405
Lime	30	1435
Red rock.....	10	1445
Lime	30	1475
Sand, Maxton (?).....	50	1525
Slate	10	1535
Red rock.....	15	1550
Slate	18	1568
Little Lime.....	12	1580
Slate	8	1588
Sand, white.....	8	1596
Slate	4	1600
Lime, sandy, white.....	56	1656
Slate	44	1700
Slate and shells.....	90	1790
Lime	9	1799
Penell Cnve.....	16	1815
Greenbrier Series (371')		
Big Lime, black (gas, 1868'; quite a puff but lasted only short time; gas, 1900-1910').....	100	1915
Big Lime, white.....	100	2015

	Thickness. Feet.	Total. Feet.
Big Lime, light-gray, sandy.....	2	2017
Big Lime, light-gray, sandy, (salt water).....	13	2030
Big Lime, white.....	57	2087
Big Lime, red (salt water).....	4	2091
Big Lime, white.....	19	2110
Big Lime, red.....	12	2122
Big Lime, white.....	64	2186
Pocono Series (507')		
Sand, Big Injun.....	81	2267
Sand, Squaw.....	43	2310
Slate, gray.....	7	2317
Shale and slate.....	23	2340
Slate, sandy.....	85	2425
Slate, Weir.....	18	2443
Slate, sandy.....	82	2525
Sand, soft, Weir.....	5	2530
Slate, sandy.....	97	2627
Sand, Weir.....	15	2642
Slate and shells, blue.....	22	2664
Slate and lime shells.....	18	2682
Slate, black, Coffee.....	6	2688
Sand, Berea.....	5	2693
Chemung Series (115'+)		
Slate, sandy, blue.....	12	2705
Slate and shells.....	103	2808

The record of the J. H. Gwinn No. 1 Well (No. 7 on Map II) is published in connection with the Green Sulphur Springs Section in Chapter V. page 195. The well was drilled near the town of Green Sulphur Springs, Summers County. Several shows of gas were reported from horizons in the Pocono and Chemung Series. The well was not located in a favorable structural position.

The record of the Gauley Coal Land Company (Granville O'Dell) No. 1 Well (No. 8 on Map II) is published in connection with the Hominy Falls Section in Chapter V, pages 177-8. This well, drilled 1.4 miles south of Hominy Falls, Nicholas County, was abandoned as a dry hole. The well was not located in a favorable structural position.

The following is the record of a well drilled near Johnson Crossroads, Monroe County. The well is located west of the axis of the Abbs Valley Anticline. It is reported that one of the reasons the well was abandoned was because of salt water coming along with the gas. The well may have been located too low on the structure:

L. E. Johnson et al. No. 4209 Well.

Monroe County, Wolf Creek District; 0.3 mile northeast of Johnson Crossroads; 3.1 miles W. of 80° 35' and 3.3 miles S. of 37° 40', Alderson Quadrangle. By United Fuel Gas Company, Charleston, W. Va. Rig commenced March 25, 1930; completed March 29, 1930. Drilling commenced April 15, 1930; completed October 17, 1930. Drilled by Drilling Dept. of Company; drillers, B. J. Dotson and W. H. McClane. 13½" casing, 287' (left in); 10", 671'; 8¼", 1578'. Shot, October 7, 1930, with 40 qts. Shot exploded by tools, 1562-1572'. Test before shot 12/10 W. in 1"=42,000 cu. ft. Test after shot 12/10 W. in 1"=42,000 cu. ft. 1—10" Star rimmer, 1—10" Wing Sub, and 1—8¼x10x13 B. H. packer left in hole.

Well plugged and abandoned October 17, 1930.

Starts about 300' below top of Greenbrier Series.

Elevation, 1904' B.

	Thickness. Total.	
	Feet.	Feet.
Greenbrier Series (1038'+)		
Clay, yellow, soft.....	12	12
Lime, black, hard.....	298	310
Sand, red, soft.....	5	315
Lime, black, hard.....	677	992
Lime, gray, hard.....	46	1038
Maccrady Series (173')		
Sand, red, soft.....	12	1050
Lime, gray, hard.....	25	1075
Shale, red, soft.....	136	1211
Pocono Series (426'+)		
Sandy lime, blue, soft (10/10 W. in 1"=37,680 cu. ft. gas, at 1230-35').....	49	1260
Slate, white, soft.....	5	1265
Sand, gray, hard.....	35	1300
Lime, white, hard.....	68	1368
Slate, black, soft.....	16	1384
Coal, black slate with some coal.....	6	1390
Lime shell.....	50	1440
Sand	106	1546
Slate	4	1550
Sand	38	1588
Lime	7	1595
Unrecorded to bottom (steel-line measure).....	42	1637

The following two records are of wells drilled near Bozoo (Chestnut Hill), Monroe County. Both wells found gas in the Pocono Series:

John T. Shumate No. 1 Well.

Summers County, Forest Hill District; by The Bozoo Company; on Crooked Run of New River, 0.6 mi. N. E. of Neponset; well was completed September 2, 1929; Contractors, Dunham & Titus; elevation,

2040' B. Gas test, 2,500,000 cu. ft. Rock pressure, 635 lbs. in 30 min. in 8¼" casing, after which well was opened and allowed to flow into Wills well at Bozoo.

Thickness. Total.
Feet. Feet.

Mauch Chunk Series—Hinton Group (50' +)

Surface soil and loose sandstone.....	12		12
Sandstone, gray (small amount of water at 20'.....13'	} Stony Gap Sandstone	38	50
Sandstone, grayish-white.....25'			

Mauch Chunk Series—Bluefield Group (1235')

Shale, red and brown.....	110		160
Limestone, bluish-gray, shaly.....	10		170
Shale, reddish-brown or gray.....	130		300
Shale, gray, calcareous.....	25		325
Shale, reddish-brown, sandy.....	55		380
Limestone, gray, shaly.....	20		400
Shale, gray, calcareous.....	62		462
Shale, red.....	28		490
Shale, gray.....	128		618
Shale, dark-gray.....	4		622
Sand, grayish-white.....	53		675
Shale, soft, gray.....	2		677
Sandstone, Droop, grayish-white.....	65		742
Shale, dark-gray, calcareous.....	108		850
Shale, brownish-gray.....	10		860
Shale, gray, sandy.....	40		900
Shale, gray, sandy and calcareous.....	50		950
Limestone, Glenray, dark-gray, shaly.....	53		1003
Shale, gray, soft.....52'	} Lillydale Shale (Penell Cave)	257	1260
Shale, gray, soft.....10'			
Shale, gray, soft, calcareous.....145'			
Shale, dark-brown, soft.....50'			
Sandstone, Edray, dark-gray, shaly, impure.....	25		1285

Greenbrier Series (1375')

Limestone, gray.....25'	} Alderson Limestone	250	1535
Limestone, dark-gray.....90'			
Limestone, bluish-gray.....135'			
Limestone, gray, hard, sandy.....30'	} Union Limestone	350	1915
Limestone, dark-gray.....110'			
Limestone, light-gray.....75'			
Limestone, gray.....100'			
Limestone, light-gray.....65'	} Pickaway Limestone	385	2300
Limestone, dark-gray, shaly.....100'			
Limestone, light-gray.....10'			
Limestone, light-gray.....65'			
Limestone, gray, shaly.....10'			
Limestone, dark-gray.....200'	} Patton Limestone	140	2540
Limestone, Taggard, gray, shaly.....			
Limestone, bluish-gray.....120'			
Limestone, gray, hard.....20'			
Limestone, Sinks Grove, dark, bluish-gray.....	120		2660

Maccrady Series (160')

Shale, gray.....	25		2685
Shale, gray, very soft.....	10		2695

	Thickness. Feet.	Total. Feet.
Shale, gray, sandy and calcareous.....	35	2730
Shale, reddish-brown and gray.....	90	2820
Pocono Series (103')		
Sandstone, gray, calcareous.....30'	} Squaw Sand?.. 103	2923
Shale, bluish-gray.....20		
Limestone, bluish-gray, shaly.....10		
Shale, dark, soft, sandy.....5		
Sandstone, grny, shaly.....31		
Sandstone, dark-gray, with quartz pebbles (gas, 2916-2923').....7		
Total depth.....		2923

G. K. Wills No. 1 Well.

Monroe County, Red Sulphur District; by The Bozoo Company; on New River plateau and axis of Abbs Valley Anticline, 0.5 mi. S. W. of Bozoo (Chestnut Hill); completed, February 21, 1929; drilled by L. H. Harrison et al.; Contractor, C. M. Means; elevation, probably somewhat less than 2100'. Shot, February 9, 1929, with 140 qts., at 3097-3129', with no increase in gas. Shot, February 14, 1929, with 40 qts., at 649'; Paeker set above lower pay sand but leaks and gas comes from this sand up into upper sand. A rock pressure reading under this condition showed 215 lbs. After allowing the well to blow off 17 hrs. and 45 min., gas test at 649' was 82,000 cu. ft.; at 3105', was 157,000 cu. ft. per day.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Hinton Group (30'+)		
Loose sandstone rocks.....	30	30
Mauch Chunk Series—Bluefield Group (1250')		
Shale, bluish-grny and reddish.....	26	56
Shale, greenish-gray (water at 70').....	16	72
Shale, reddish-brown.....	6	78
Sandstone, red, fine-grained.....	7	85
Shale, gray.....	9	94
Shale, reddish-brown.....	9	103
Sandstone, light greenish-gray, fine-grained, compact..	4	107
Shale, green and dark-green, containing some organic matter	33	140
Shale, reddish-brown.....	33	173
Sandstone, gray.....	4	177
Shale, red, platy, with conspicuous mica flakes.....	51	228
Shale, hard, slaty.....	12	240
Shale, reddish-brown.....	5	245
Shale, greenish-gray.....	35	280
Shale, dark-gray, sandy.....	22	302
Sandstone, greenish-gray, shaly.....	18	320
Shale, dark-gray.....	10	330
Shale, greenish-gray.....	25	355
Sandstone, gray, shaly.....	12	367
Shale, dark-gray, sandy.....	36	403
Limestone, grny, shaly.....	6	409
Shale, gray, sandy.....	31	440

sandstones with an occasional thin limestone. In most localities the presence of the **Fossil Ore Horizon** is found above the middle, which in turn was preceded by the deposition of beds predominantly shaly but containing platy sandstones with occasional thin limestones. Toward the base the sandstones increase in number and thickness and are more compact and greenish to gray in color, except the **Iron Sandstone**, which occurs in the lower portion, and which is generally more massive and red in color. The Clinton Series, although no complete and continuous exposures are found in this area, is approximately 600 feet in thickness.

The Clinton Series has received considerable attention from many geologists and hence there is a great deal of literature available with reference to it. The early work on these beds was done by Eaton, Hall, and others in New York State where its character is such that subdivisions as found there can not be applied with certainty in this area. In later work in Pennsylvania, the subdivisions of H. D. Rogers, as later revised by Dr. I. C. White¹, seem best adapted to the local area, except that the Keefer Sandstone that is now recognized as of Clinton age was not included. Their subdivision follows in descending stratigraphic order:

- Upper Shales.
- Ore Sandstone and Fossil Ore.
- Middle Shales.
- Iron Sandstone and Block Ore.
- Lower Shales.

In a still later work, Swartz² has given these beds the following classification:

- Clinton Group.
 - Rochester Formation.
 - Upper Shale and Limestone.
 - Roberts Iron Ore.
 - Keefer Sandstone Member.
 - Rose Hill Formation.
 - Upper Shale beds with some purplish bands.
 - Cresaptown Iron Sandstone.
 - Lower shale and sandstone beds.

¹See Second Geol. Surv. of Pa., Reports G7, pp. 111-112; 1883; and T3, p. 132; 1885.

²Charles K. Swartz, Silurian volume, Md. Geol. Surv., pp. 27-35; 1923.

It has been previously stated that in this county the Clinton is confined to those beds occurring between the top of the White Medina Sandstone and the top of the Keefer Sandstone, but at the same time recognizing the possibility of a small portion of those beds occurring immediately above the Keefer as being of Rochester or Clinton age. Since sufficient exposures are not available in this area to add much to a detailed discussion of the finer aspects of this series, the following subdivisions are used:

Upper Shales.

Keefer Sandstone.

Shales and thin limestones.

Fossil Ore Horizon

Middle Shales (including platy sandstones and thin limestones).

Iron Sandstone.

TOPOGRAPHIC EXPRESSION, CLINTON SERIES.

The Keefer Sandstone at or near the top of the Clinton as well as the Iron Sandstone in the lower portion are both quartzitic in character and resistant to weathering and frequently form sharp and prominent ridges. The lower portion is more sandy than the Upper and Middle Shales so that the lower portion of this series forms prominent shoulders along with the underlying White Medina. The upper and middle shaly members are less resistant and form a line of weakness in the Clinton outcrop represented by a depression in the topography along the east side of Beaver Lick Mountain.

AREAL EXTENT, CLINTON SERIES.

On Figure 15, the outcrops of the Clinton are shown, and can be seen in greater detail and on a larger scale on Map II accompanying this report. The exposures of the Clinton are confined to the Beaver Lick Mountain area in the northeastern portion of the county. Several isolated outcrops of this series occur in the folds of Beaver Lick Mountain but are all poorly exposed. The highway west of Alvon cuts across the Browns Mountain Anticline, but an accumulation of talus practically conceals these outcrops.



PLATE XLII.—Apparent unconformity at or near base of Helderberg Limestone. At spring-house of Alvon
springs Nos. 1 and 2 southwest of Alvon.




PLATE XLIII.—Niagara Limestone showing numerous veins of calcite, on the north side of Anthony Creek, 0.5 mile west of Alvon.





PLATE XLV.—La Barr Evergreen Nursery north of White Sulphur Springs.



CONTACTS, CLINTON SERIES.

The upper contact of the Clinton Series with the overlying Niagara has already been discussed under the latter series on page 333. Attention was called to the fact that in mapping the contact was placed at the top of the Keefer Sandstone and also to the possibility of a few feet of the overlying shales being of Rochester or Clinton age. The lower contact with the White Medina appears, in places, to be transitional, while at others evidence of erosional unconformity is noticeable. In the local area the contact is placed above the grayish-brown to white, quartzitic sandstones that are devoid of any organic remains other than *trilobites* and *Arthropycus alleghaniensis*.

FOSSIL LIFE, CLINTON SERIES.

The Clinton Series was found to be sparingly fossiliferous in this area, marine fossils being noted.

CORRELATION, CLINTON SERIES.

Under the heading "General Account," attention has already been called to certain relationships of the Clinton Series as found in Greenbrier County to those of its more northeastern counterparts in Maryland, Pennsylvania, and New York. The **Keefer Sandstone** of Stose and Swartz² was named from its occurrence in south-central Pennsylvania and has now been traced southwestward across Maryland and West Virginia. Farther down in the series a thin bed of iron ore occurs, along with shaly and siliceous, fossiliferous limestones, that correlates with the Fossil Ore Horizon of eastern Pennsylvania. In the basal portion of the Clinton in this area there is a prominent red sandstone which attains a thickness of as much as 50 feet. It weathers into rectangular blocks and makes a heavy talus. There is little doubt that this sandstone correlates with the **Iron Sandstone and Block Ore** of Rogers and White, which in turn is synonymous with **Cresaptown Iron Sandstone** of the Maryland Geological Survey. It has also been included with the **Cacapon** by Darton in the Monterey Folio.

²G. W. Stose and C. K. Swartz, Pawpaw-Hancock Folio, No. 179, U. S. Geol. Surv.: 1912.

DESCRIPTION OF MEMBERS, CLINTON SERIES.

UPPER SHALES.

The shales coming above the Keefer and referred to as the **Upper Shales** do not appear to be generally present in Greenbrier County. At most points where the Keefer is exposed the immediately overlying interval is concealed so that their presence can not be definitely proved.

KEEFER SANDSTONE.

The **Keefer Sandstone** was first named by Stose and Swartz⁴ from its occurrence in Keefer Mountain, Pennsylvania, a few miles northeast of Hancock, Md., where it forms a thick and massive bed. In Greenbrier County this same member is present, being composed of grayish sandstones that vary from four to eight feet in thickness, often quartzitic in character, and having a total thickness of 35 to 60 feet. This member was noted at several points in the county and can best be seen 0.5 mile west of Alvon.

SHALES AND THIN LIMESTONES.

Between the Keefer Sandstone and the Fossil Iron Ore Horizon there occurs a succession of yellowish-gray to olive, thin shales and platy sandstones with occasional limestones one to six inches in thickness.

FOSSIL ORE HORIZON.

The **Fossil Ore Horizon** was noted at only one locality, that being on Beaver Lick Mountain just south of the position of Cross-Section A—A' as shown on Map II.

MIDDLE SHALES.

The **Middle Shales** occupy the interval between the Fossil Ore Horizon and Iron Sandstone. These shales vary in color from yellow and olive to green, red, or dark, and attain a thickness of approximately 250 feet. Occasionally calcareous lenses and streaks occur along with thin sandstones.

⁴Ibid.

IRON SANDSTONE.

The **Iron Sandstone** in this area has a deep-red color and consists of quartz grains cemented with hematite. It is often oolitic in texture. The more ferruginous beds resemble a low-grade iron ore but the proportion of silica is entirely too high to permit their use as a source for commercial iron at the present time.

ECONOMIC ASPECTS, CLINTON SERIES.

At certain points in Greenbrier County the Clinton Series contains local deposits of iron ore that are of good enough quality to encourage more thorough prospecting. At no point in fresh exposures was there found ore of minable thickness in the Fossil Ore Horizon, and the Iron Sandstone is too low in ore to be used for this purpose. At some points, however, where the rocks are so folded as to form troughs or basins, there will probably be found better grade ores due to local enrichment from leaching of the higher beds. This will require further prospecting at such points. A further discussion of these ores will appear under their respective headings in Chapter XIII.

Many of the Clinton sandstones are of sufficient hardness to be used as a building stone. The Iron Sandstone breaks into rectangular blocks and is of a pleasing red color so that it is admirably adapted for that purpose.

WHITE MEDINA SERIES.

GENERAL ACCOUNT, WHITE MEDINA SERIES.

The **White Medina Series**, coming just below the Clinton Series and at the top of the three Medinas as recognized in West Virginia, is present in Greenbrier County, being a prominent white quartzite and varying in thickness from 50 to 100 feet. Its greater portion is thick-bedded and carries a siliceous cement so that it is very resistant to weathering and makes prominent ridges. It often contains rounded white quartz pebbles.

TOPOGRAPHIC EXPRESSION, WHITE MEDINA SERIES.

The White Medina, on account of its quartzitic character and massive bedding, is the most resistant to weathering of any rock exposed in the county. Its exposures are always

marked by rugged topography. In the area of its outcrop it is the chief ridge-forming rock and great blocks of the sandstone, which frequently forms the crests of the mountains, break away from the ledge and work by gravity down the steep slopes and frequently conceal the underlying formations.

AREAL EXTENT, WHITE MEDINA SERIES.

On Figure 15, the outcrop of the White Medina can be seen at a glance together with the other Silurian Rocks, while on Map II these same exposures are shown in greater detail and on a larger scale. These exposures are limited to the Browns Mountain Anticlinal area, being confined to Beaver Lick Mountain. Along the crest of Beaver Lick Mountain this sandstone stands at very high angles.

CONTACTS, WHITE MEDINA SERIES.

The upper contact of the White Medina has already been discussed in the description of the Clinton Series. Its base rests upon the red shales and red sandstones of the Red Medina Series. The contact, however, is not so pronounced as would generally be expected between beds so vastly different. The change from red to white is transitional.

FOSSIL LIFE, WHITE MEDINA SERIES.

The White Medina in Greenbrier County, as in other localities, is sparingly fossiliferous. The most abundant species is *Arthropycus alleghaniensis*, a trail resembling a seaweed, which is often found covering the under-side of these beds with its numerous interlacing "stems." Straight tubular borings occasionally refilled and standing at right angles to the bedding are found and are believed to be the same as similar borings found in the Medina of New York and named *Scolithus verticalis* by Hall. This is one of the characteristic fossils of the White Medina, being widely distributed at this horizon throughout the Appalachian area. In the upper portion of these beds there occurs an abundance of small stem-like, rounded and semirounded forms, that are both single and branching. The surface is smooth and without markings but does not retain a uniform width as in *Arthropycus alleghaniensis*. These forms, while probably of organic origin, are only classed in general as fucoids.

CORRELATION, WHITE MEDINA SERIES.

The White Medina as recognized in West Virginia and where it has been traced entirely across the State, following the Appalachian counties as it does, has been described under different names in other localities. That it corresponds to the White Medina of the New York and Pennsylvania Surveys appears to be without doubt. The name **Albion** was given it in New York by Kindle². In various Folios of the U. S. Geological Survey it is called **Tuscarora**, named from its outcrop in Tuscarora Mountains in Pennsylvania. In the adjoining State of Virginia and other southern Appalachian States it correlates with the **Clinch**.

ECONOMIC ASPECTS, WHITE MEDINA SERIES.

The White Medina, while very hard and resistant, has not been used as a building stone because it can not be satisfactorily split into blocks. It has been used, particularly in adjoining counties of Virginia, as a base for hard-surfaced roads. It contains a high percentage of silica but its use as a glass-sand has not proved satisfactory because of its conglomeratic character. The white quartzitic members are suited for gneiss and should be suitable for various trap-rock uses.

RED MEDINA SERIES.

GENERAL ACCOUNT, RED MEDINA SERIES.

In Greenbrier County the Red Medina Series is exposed only along the west side of Beaver Lick Mountain, outcropping in a small area about 3 miles long by 0.2 mile wide. In this area the rocks have been faulted by overthrusting and the Red Medina is now resting on the crumpled black shales of the Marcellus Series. As is to be expected, the rocks along the fault have been greatly mashed and in the case of the Red Medina, the bedding-planes and direction of dip can be ascertained only with the greatest difficulty. So far as can be told under such circumstances, the Red Medina rocks are composed of alternating red sandstones and red sandy shales.

²E. M. Kindle and F. B. Taylor, U. S. Geol. Survey, Niagara Folio, No. 190; 1913.

One of the surprising things about the Red Medina rocks along the fault is that they have not been fused into quartzites but rather they appear to have been thoroughly disintegrated. Just how much of this disintegration is due to weathering and how much is due to lack of metamorphism is unknown.

As exposed in Greenbrier County, the Red Medina appears to be about 800 feet thick but as pointed out above the bedding is indistinct so that the measurement is really the interval between the fault-plane and the White Medina and may not be the true vertical thickness of the series.

TOPOGRAPHIC EXPRESSION, RED MEDINA SERIES.

Outcropping as they do along the Burr Fault on the west side of Beaver Lick Mountain, the Red Medina rocks can not be spoken of as possessing a type of topographic expression.

AREAL EXTENT, RED MEDINA SERIES.

The Red Medina is exposed only along the west side of Beaver Lick Mountain in an area about 3 miles long by 0.2 mile wide, near the Pocahontas County line as shown on Map II.

CONTACTS, RED MEDINA SERIES.

The upper boundary of the Red Medina with the White Medina has already been discussed in the description of the latter formation, attention being called to the gradual change from one to the other. As the lower limit of the Red Medina, in this area, is a fault-plane, the nature of the contact can not be discussed in a stratigraphic sense.

FOSSIL LIFE, RED MEDINA SERIES.

The Red Medina has been quite generally considered non-fossiliferous and this same condition prevails in Greenbrier County.

CORRELATION, RED MEDINA SERIES.

The Red Medina of Greenbrier County appears to be the same as the **Juniata** of the various Folios of the U. S. Geological Survey. The Juniata is called by the same name in Pennsylvania which is in turn correlated with the **Queenston** of New York. The Juniata and Queenston are believed by some to be the same as the **Richmond** group of Ohio.

New Scotland Member.

The New Scotland is represented in Greenbrier County by a calcareous sandstone 25 to 40 feet thick. This sandstone has been named the **Healing Springs** by Swartz¹⁶, from its occurrence near Healing Springs, Virginia. In general the sandstone is medium- to fine-grained, light-gray to light-brown in color and is characteristically marked by numerous casts of medium- to large-sized crinoid stems. Being somewhat quartzitic, the sandstone weathers in bold relief and is quite conspicuous in the vicinity of Alvon. As mentioned above, the lower part of the limestone herein described as Beerat carries some fossils that are suggestive of the New Scotland. There is no prominent lithologic break within the limestone and the contact of the limestone with the underlying Healing Springs is blended as indicated in the following section:

White Sulphur Springs Section.

White Sulphur District, 1 mile north of White Sulphur Springs, on Howard Creek; measured on the west side of Bobs Ridge; arrangement in descending stratigraphic order.

Thickness, Total.
Feet. Feet.

Heiderberg Series (110'±)	
Limestone, blue-black, nodular, with blue-black nodular chert	80
Sandstone Healing Springs, gradual transition from overlying limestone, light-gray to light-brown on fresh exposure, calcareous, fossiliferous.....	30.
The lower part of the Healing Springs Sandstone is not well exposed in Greenbrier County and the exact nature of its contact with the underlying limestone is not known.	110

The fossils collected from the horizon of this sandstone were badly weathered and due to this fact the New Scotland age of this sandstone may be considered to be slightly in doubt. However, on the basis of its lithologic characteristics, its stratigraphic position, and its fauna, which strongly suggest New Scotland forms, the correlation of this sandstone with the Healing Springs of the type locality appears to be established. Chemical analyses of samples collected from the Healing Springs Sandstone are presented and discussed in Chapter XII.

¹⁶Ibid., p. 41.

New Scotland Member.

The New Scotland is represented in Greenbrier County by a calcareous sandstone 25 to 40 feet thick. This sandstone has been named the **Healing Springs** by Swartz¹⁶, from its occurrence near Healing Springs, Virginia. In general the sandstone is medium- to fine-grained, light-gray to light-brown in color and is characteristically marked by numerous casts of medium- to large-sized crinoid stems. Being somewhat quartzitic, the sandstone weathers in bold relief and is quite conspicuous in the vicinity of Alvon. As mentioned above, the lower part of the limestone herein described as Beecraft carries some fossils that are suggestive of the New Scotland. There is no prominent lithologic break within the limestone and the contact of the limestone with the underlying Healing Springs is blended as indicated in the following section:

White Sulphur Springs Section.

White Sulphur District; 1 mile north of White Sulphur Springs, on Howard Creek; measured on the west side of Bobs Ridge; arrangement in descending stratigraphic order.

	Thickness. Total.	
	Feet.	Feet.
Holderberg Series (110'+)		
Limestone, blue-black, nodular, with blue-black nodular chert	80	80
Sandstone Healing Springs , gradual transition from overlying limestone, light-gray to light-brown on fresh exposure, calcareous, fossiliferous.....	30.	110

The lower part of the Healing Springs Sandstone is not well exposed in Greenbrier County and the exact nature of its contact with the underlying limestone is not known.

The fossils collected from the horizon of this sandstone were badly weathered and due to this fact the New Scotland age of this sandstone may be considered to be slightly in doubt. However, on the basis of its lithologic characteristics, its stratigraphic position, and its fauna, which strongly suggest New Scotland forms, the correlation of this sandstone with the Healing Springs of the type locality appears to be established.

Chemical analyses of samples collected from the Healing Springs Sandstone are presented and discussed in Chapter XII.

¹⁶Ibid., p. 41.

Coeymans Member.

The horizon at which the Coeymans would be expected is usually not exposed in Greenbrier County. Under these circumstances it was not possible to prove either the presence or absence of rocks of this age in the territory of this report. If it is present in Greenbrier County the Coeymans Member is not over 40 feet thick and is probably less than 10 feet thick.

Keyser Member.

In Greenbrier County the Keyser Member is best exposed along the north side of Anthony Creek, on the west limb of the Browns Mountain Anticline, just west of Alvon. The following section illustrates its occurrence at this point and is part of the Alvon Section—West Side, published in Chapter V:

Part of Alvon Section—West Side.

Devonian.	Thickness. Total.	
Helderberg Series.	Feet.	Feet.
Keyser Member (215'±).		
Concealed and shaly limestone (Coeymans, if present)	35	35
Limestone, sandy to shaly.....	20	55
Limestone, gray, platy, calcite streaks.....	20	75
Limestone, blue-gray, massive, calcite streaks.....	40	115
Limestone, blue-gray.....	20	135
Concealed and gray limestone.....	65	200
Sandstone, Clifton Forge, fine-grained, hard, porous, and limonite stained from weathering, upper portion strongly cemented by silica.....	15	215
Silurian.		

At the above locality the rocks are vertical or slightly overturned and accurate total measurements are difficult to obtain. The sandstone noted at the base of the Keyser in the above section is considered to be the same as the Clifton Forge Sandstone of Swartz¹⁷. This sandstone is either concealed or absent at many localities in the county. The lower contact of the Clifton Forge at the above section is poorly exposed but it is assumed that it marks the base of the Helderberg.

¹⁷Ibid., p. 29.

Some of the limestone beds of the Helderberg are of sufficient thickness and purity for lime-burning and other purposes for which a fairly pure limestone is required. It is doubtful if the Helderberg limestones will be used for this purpose, however, since the pure limestones of the Silurian and Mississippian present better quarry sites in the area. The Healing Springs Sandstone might be used for glass-sand. The commercial possibilities of the limestones and sandstones will be discussed in Chapter XII. In some places the residual soil left from weathering of the Keyser beds contains nodules of manganese ore and this together with a discussion of the springs that emerge from Helderberg rocks will be discussed in Chapter XIII.

CHAPTER IX.

STRATIGRAPHY—SILURIAN ROCKS.

GENERAL STATEMENT.

The Silurian Rocks as found in Greenbrier County, West Virginia, and as indicated in the General Columnar Section, page 133, have been classified, in descending stratigraphic order, with certain titles being added in parentheses to indicate supposed contemporaneous nomenclature, as follows:

	Thickness.	Total.
	Feet.	Feet.
Salina Series		
Bossardville Group (Tentaculite, Manlius, Tonoloway)	250	250
Rondout Waterlime Group (Wills Creek).....	200	450
Niagara Series (McKenzie).....	100	550
Clinton Series (Rockwood and Caeapon of U. S. G. S. Folios; Rochester of Maryland and Clinton of New York)	600	1150
White Medina Series (Tuscarora, Albion, Clinch, of U. S. G. S. Folios).....	100	1250
Red Medina Series (Juniata of U. S. G. S. and Queenston Shale of other Reports).....	800	2050

As shown on Figure 15, the outcrop of the Silurian in Greenbrier County is confined to the immediate vicinity of Beaver Lick Mountain. The rocks have been severely folded, mashed and in some places metamorphosed so that accurate measurements are very difficult to obtain. At no point was it possible to measure a complete succession of the beds in detail, because of duplication by folding, or on account of concealed intervals, but exposures of all the individual groups are available at one point or another.

FIGURE 15

MAP

OF

GREENBRIER COUNTY

SHOWING

OUTCROPS OF

SILURIAN ROCKS



The upper boundary of the Silurian is generally agreed upon as coming at the base of the Helderberg. The lower boundary or the contact of the Silurian with the Ordovician has long been a subject of debate, nor is there yet a general agreement on this point. It has been the policy of the West Virginia Geological Survey to place the base of the Silurian at the bottom of the **Gray Medina Sandstone**. As the oldest rocks exposed in Greenbrier County are the Red Medina sandstones and shales, there is no information available in this area that might throw any additional light on this controversy. In this report the Medinas are considered to be of Silurian age.

The Silurian as thus delimited begins in Greenbrier County with Red Medina beds, suggesting rapid deposition with poor sorting of the materials and estuarine or land deposits. In the following epoch the beds are mainly of white sand with white quartz pebbles that represent a shore phase of a transgressing sea. As the sea deepened there followed a succession of shales and sandstones of lower and middle Clinton with marine fossils. Following this the beds became more calcareous, contain marine fossils, and show the effect of a retreating sea marked by the Bossardville laminated limestones.

The Silurian deposition is a good illustration of a cycle of sea inundation and retreat, marked by times of recession, slight reversals, and the separation of sea basins.

SALINA SERIES.

GENERAL ACCOUNT, SALINA SERIES.

The Salina Series as found in Greenbrier County is divided, following earlier subdivisions, into the upper portion, Bossardville, containing platy and laminated limestones and the lower portion, Rondout, which is made up of interbedded calcareous shales and limestones. In this area clean-cut exposures of the Salina are not available and while, in general, there is a marked contrast between the two groups, the change from one to the other is gradual rather than abrupt.

It is the Salina Series that contains the rock salt and anhydrite that is found in the drilled wells in the western part of the State. A search was made for gypsum or anhydrite at the outcrop of this series in Greenbrier County but none was found.

TOPOGRAPHIC EXPRESSION, SALINA SERIES.

The Salina Series in Greenbrier County can not be spoken of as having a characteristic topography. In this area the beds have been greatly folded and are now found standing at steep angles along the east side of Beaver Lick Mountain. The Salina limestones, along with the Niagara, due to their soluble character, are generally found in narrow valleys between the more resistant Lower Devonian rocks and the underlying Clinton and Medina Series.

AREAL EXTENT, SALINA SERIES.

The areal extent of the Salina Series is shown on Figure 15 along with the rest of the Silurian. These same exposures are shown on Map II in much greater detail and on a larger scale. The outcrop of this series is confined to Beaver Lick Mountain and the northern end of Coles Mountain.

CONTACTS, SALINA SERIES.

The upper contact of the Salina Series with the Helderberg of the Devonian has already been discussed under the latter series. The lower contact of this series is not well exposed in Greenbrier County and its exact nature is not known. It is assumed, however, that the beds above and below it are conformable and that the Bloomsburg Group is absent because of non-deposition or changing conditions of sedimentation.

FOSSIL LIFE, SALINA SERIES.

In Greenbrier County the Salina rocks do not yield their fossils readily but fragments of marine fossils were noted in these rocks at various localities. *Camarotoechia tonolowayensis* and *Camarotoechia hitchfieldensis* were the most abundant of those noted in the field.

CORRELATION, SALINA SERIES.

Certain relationships of the Salina Series as found in Greenbrier County with their northeastern counterparts in other States have already been suggested. More definite detailed correlation necessitates better exposures and more numerous systematic fossil collections than are available in this area. That the upper or Bossardville Group of this series

correlates with this same formation to the northeast is quite certain, and this in turn is essentially synonymous with the Tentaculite, Manlius, and Tonoloway, as pointed out by Reger in Chapter XIV of the Mineral and Grant County Report. The Rondout Waterlime Group although somewhat attenuated retains in general the same character as found at its type locality in New York, and can safely be correlated with it. This group correlates with the **Wills Creek Formation** of Maryland, and is included under the **Lewistown Limestone** in the Monterey Folio.

DESCRIPTION OF GROUPS, SALINA SERIES.

BOSSARDVILLE LIMESTONE GROUP.

The Bossardville Group is made up largely of limestone which is thin-bedded and laminated. These thin beds of laminated limestone are often separated by thin shale partings, the limestone slabs weathering out and frequently covering the surface slopes in the area of the outcrop, so that it is easy to distinguish this formation at some distance. These slabs or fragments often have a noticeable cleavage, and break in rough geometric figures. Certain beds carry an abundance of fossils of few species. In the general section this group is shown to be about 250 feet in thickness. The figure may be excessive as no complete exposures were available for accurate measurement.

RONDOUT WATERLIME GROUP.

The Rondout Waterlime consists of interbedded calcareous shale, calcareous mud rock, and argillaceous limestone with an occasional sandstone. When seen in fresh exposures many of the strata seem to consist of compact, dark, purplish-blue limestone of considerable durability, but on weathering, however, the color of these strata changes to a dirty greenish hue. This same characteristic was noted by the writer in other counties of this State to the northeast, especially Pocahontas, Pendleton, Hampshire, and Hardy, and is also reported in Maryland. This feature is due to the large amount of clay that is present in the rock. Alternating with these rocks are beds of thin-bedded, fissile, and calcareous shale that are occa-

sionally dark. With these highly argillaceous beds are occasional strata of purer limestone. The Rondout Group as found in this area has a thickness of about 200 feet.

ECONOMIC ASPECTS, SALINA SERIES.

In Greenbrier County the principal economic value of the Salina Series is its use for agricultural purposes, a great deal of the limestone being suitable for burning, both for agricultural lime and Portland cement. The upper portion, or Bos-sardville Group, generally carries a high calcium carbonate content, the main impurity being silica or alumina which breaks down readily, so that long burning is not necessary. In the Rondout Group certain portions have been used for the manufacture of natural cement in northeastern West Virginia and western Maryland, but in Greenbrier County chemical analyses have not been made. Because of its generally inaccessible location, its value for road material in this area is overshadowed by the more readily obtainable limestone from the Greenbrier Series and the Huntersville Chert of the Oriskany Series.

NIAGARA SERIES.

GENERAL ACCOUNT, NIAGARA SERIES.

The **Niagara Series**, coming just below the Salina Series and slightly above the **Keefer Sandstone** of the Clinton Series, is a succession of shales and beds of limestone. The shales are generally buff or drab while the limestones vary from bluish-gray to dove-colored. This series, because of its non-resistant nature and its occurrence at high angle dips is poorly exposed, so that accurate measurements were difficult to get. An interval of 50 to 100 feet will include both the minimum and maximum thicknesses of this series in Greenbrier County.

The Niagara beds of New York were early subdivided by James Hall into Niagara or **Lockport Limestone** at the top, followed by the Niagara or **Rochester Shale** at the base. In the Pawpaw-Hancock Folio, Stose and Swartz described those beds occurring between the Bloomsburg red sandstone member of the Wills Creek Shale and the Clinton Series as **McKenzie Formation**, including the Keefer Sandstone. In its Silurian

volume, the Maryland Geological Survey considers the Keefer Sandstone as of Clinton age. It is, therefore, the beds that occur between the Salina Series and the Keefer Sandstone that are classified as the Niagara Series in this report. In this area there is not sufficient variation in lithology from top to bottom to form the basis of any subdivision.

TOPOGRAPHIC EXPRESSION, NIAGARA SERIES.

The Niagara Series, being predominantly shaly, is much less resistant to weathering than the Keefer Sandstone below. It has no tendency to cliff forming and is seldom seen in good exposures save in localities where it has been uncovered in stream gullies or by artificial cuts.

AREAL EXTENT, NIAGARA SERIES.

The Niagara Series with its narrow outcrop, the beds areal extent in Greenbrier County. Its exposures are shown on Figure 15 along with the Silurian Rocks, on page 328, and in much greater detail and on a larger scale on Map II accompanying this report. These exposures are limited to the Browns Mountain Anticlinal area, which is located east of the Greenbrier River.

CONTACTS, NIAGARA SERIES.

The upper contact of the Niagara with the Salina above has already been discussed under the same heading on the Salina Series, page 330. The lower limit of the Niagara is difficult to determine both because of the scarcity of fossils in this horizon and because there are few localities where the rocks immediately above the Keefer Sandstone are well exposed. For this reason and to facilitate areal mapping, the contact is placed at the top of the Keefer Sandstone. It is probable, however, that a few feet at least of those beds occurring above the Keefer are of Rochester age.

FOSSIL LIFE, NIAGARA SERIES.

Few collections were made from the Niagara Series but marine fossils in this series are quite common, the following being particularly noted: *Favosites*, both *marylandica* and *niagarensis*, *Leptaena rhomboidalis*, gastropods, and several species of ostracods.

volume, the Maryland Geological Survey considers the Keefe Sandstone as of Clinton age. It is, therefore, the beds that occur between the Salina Series and the Keefe Sandstone that are classified as the Niagara Series in this report. In this area there is not sufficient variation in lithology from top to bottom to form the basis of any subdivision.

TOPOGRAPHIC EXPRESSION, NIAGARA SERIES.

The Niagara Series, being predominantly shaly, is much less resistant to weathering than the Keefe Sandstone below. It has no tendency to cliff forming and is seldom seen in good exposures save in localities where it has been uncovered in stream gullies or by artificial cuts.

AREAL EXTENT, NIAGARA SERIES.

The Niagara Series with its narrow outcrop, the beds of which are usually standing at steep dips, has a very limited areal extent in Greenbrier County. Its exposures are shown on Figure 15 along with the Silurian Rocks, on page 328, and in much greater detail and on a larger scale on Map II accompanying this report. These exposures are limited to the Browns Mountain Anticlinal area, which is located east of the Greenbrier River.

CONTACTS, NIAGARA SERIES.

The upper contact of the Niagara with the Salina above has already been discussed under the same heading on the Salina Series, page 330. The lower limit of the Niagara is difficult to determine both because of the scarcity of fossils in this horizon and because there are few localities where the rocks immediately above the Keefe Sandstone are well exposed. For this reason and to facilitate areal mapping, the contact is placed at the top of the Keefe Sandstone. It is probable, however, that a few feet at least of those beds occurring above the Keefe are of Rochester age.

FOSSIL LIFE, NIAGARA SERIES.

Few collections were made from the Niagara Series but marine fossils in this series are quite common, the following being particularly noted: *Favosites*, both *marylandica* and *niagarensis*, *Leptaena rhomboidalis*, gastropods, and several species of ostracods.

sionally dark. With these highly argillaceous beds are occasional strata of pure limestone. The Rondout Group as found in this area has a thickness of about 200 feet.

ECONOMIC ASPECTS, SALINA SERIES.

In Greenbrier County the principal economic value of the Salina Series is its use for agricultural purposes, a great deal of the limestone being suitable for burning, both for agricultural lime and Portland cement. The upper portion, or Bos-sardville Group, generally carries a high calcium carbonate content, the main impurity being silica or alumina which breaks down readily, so that long burning is not necessary. In the Rondout Group certain portions have been used for the manufacture of natural cement in northeastern West Virginia and western Maryland, but in Greenbrier County chemical analyses have not been made. Because of its generally inaccessible location, its value for road material in this area is overshadowed by the more readily obtainable limestone from the Greenbrier Series and the Huntersville Chert of the Oriskany Series.

NIAGARA SERIES.

GENERAL ACCOUNT, NIAGARA SERIES.

The Niagara Series, coming just below the Salina Series and slightly above the Kefer Sandstone of the Clinton Series, is a succession of shales and beds of limestone. The shales are generally buff or drab while the limestones vary from bluish-gray to dove-colored. This series, because of its non-resistant nature and its occurrence at high angle dips is poorly exposed, so that accurate measurements were difficult to get. An interval of 50 to 100 feet will include both the minimum and maximum thicknesses of this series in Greenbrier County.

The Niagara beds of New York were early subdivided by James Hall into Niagara or Lockport Limestone at the top, followed by the Niagara or Rochester Shale at the base. In the Pawpaw-Hancock Folio, Stose and Swartz described those beds occurring between the Bloombsburg red sandstone member of the Wills Creek Shale and the Clinton Series as McKenzie Formation, including the Kefer Sandstone. In its Silurian

sionally dark. With these highly argillaceous beds are occasional strata of purer limestone. The Rondout Group as found in this area has a thickness of about 200 feet.

ECONOMIC ASPECTS, SALINA SERIES.

In Greenbrier County the principal economic value of the Salina Series is its use for agricultural purposes, a great deal of the limestone being suitable for burning, both for agricultural lime and Portland cement. The upper portion, or Bos-sardville Group, generally carries a high calcium carbonate content, the main impurity being silica or alumina which breaks down readily, so that long burning is not necessary. In the Rondout Group certain portions have been used for the manufacture of natural cement in northeastern West Virginia and western Maryland, but in Greenbrier County chemical analyses have not been made. Because of its generally inaccessible location, its value for road material in this area is overshadowed by the more readily obtainable limestone from the Greenbrier Series and the Huntersville Chert of the Oriskany Series.

NIAGARA SERIES.

GENERAL ACCOUNT, NIAGARA SERIES.

The **Niagara Series**, coming just below the Salina Series and slightly above the **Keefer Sandstone** of the Clinton Series, is a succession of shales and beds of limestone. The shales are generally buff or drab while the limestones vary from bluish-gray to dove-colored. This series, because of its non-resistant nature and its occurrence at high angle dips is poorly exposed, so that accurate measurements were difficult to get. An interval of 50 to 100 feet will include both the minimum and maximum thicknesses of this series in Greenbrier County.

The Niagara beds of New York were early subdivided by James Hall into Niagara or **Lockport Limestone** at the top, followed by the Niagara or **Rochester Shale** at the base. In the Pawpaw-Hancock Folio, Stose and Swartz described those beds occurring between the Bloomsburg red sandstone member of the Wills Creek Shale and the Clinton Series as **McKenzie Formation**, including the Keefer Sandstone. In its Silurian

volume, the Maryland Geological Survey considers the Keefer Sandstone as of Clinton age. It is, therefore, the beds that occur between the Salina Series and the Keefer Sandstone that are classified as the Niagara Series in this report. In this area there is not sufficient variation in lithology from top to bottom to form the basis of any subdivision.

TOPOGRAPHIC EXPRESSION, NIAGARA SERIES.

The Niagara Series, being predominantly shaly, is much less resistant to weathering than the Keefer Sandstone below. It has no tendency to cliff forming and is seldom seen in good exposures save in localities where it has been uncovered in stream gullies or by artificial cuts.

AREAL EXTENT, NIAGARA SERIES.

The Niagara Series with its narrow outcrop, the beds of which are usually standing at steep dips, has a very limited areal extent in Greenbrier County. Its exposures are shown on Figure 15 along with the Silurian Rocks, on page 328, and in much greater detail and on a larger scale on Map II accompanying this report. These exposures are limited to the Browns Mountain Anticlinal area, which is located east of the Greenbrier River.

CONTACTS, NIAGARA SERIES.

The upper contact of the Niagara with the Salina above has already been discussed under the same heading on the Salina Series, page 330. The lower limit of the Niagara is difficult to determine both because of the scarcity of fossils in this horizon and because there are few localities where the rocks immediately above the Keefer Sandstone are well exposed. For this reason and to facilitate areal mapping, the contact is placed at the top of the Keefer Sandstone. It is probable, however, that a few feet at least of those beds occurring above the Keefer are of Rochester age.

FOSSIL LIFE, NIAGARA SERIES.

Few collections were made from the Niagara Series but marine fossils in this series are quite common, the following being particularly noted: *Favosites*, both *marylandica* and *niagarensis*, *Leptaena rhomboidalis*, gastropods, and several species of ostracods.

CORRELATION, NIAGARA SERIES.

The relationship of the Niagara Series as found in Greenbrier County to its counterparts, particularly to the northeast, in West Virginia, Maryland, and New York, has already been briefly touched upon under previous headings. It is not considered advisable to attempt any subdivision of this series other than to note the points of similarity with synonymous beds in other areas. In the upper two-thirds of the Niagara beds, there occurs an assemblage of fossils, all of which are found in the McKenzie Formation of Maryland, and would seem to be synonymous with it. In view of this similarity it would seem that the Niagara Series as found in this area is essentially of the same age as the McKenzie of Maryland.

DESCRIPTION OF MEMBERS, NIAGARA SERIES.

As already stated the local Niagara appears to be confined to a single lithological unit and hence the general description of the series, as already given, embraces the description of the members.

ECONOMIC ASPECTS, NIAGARA SERIES.

From an economic standpoint the Niagara Series is of minor importance, its chief value being, when found on comparatively level land, as an agricultural soil. The shales are excellent for surfacing light-traffic roads as they contain a natural mixture of sand and clay with some lime to act as a cementing agent. Just west of Alvon the C. C. C. workers have established a small temporary quarry in this series and are using the limestone for masonry work.

CLINTON SERIES.

GENERAL ACCOUNT, CLINTON SERIES.

The **Clinton Series**, occurring next below the Niagara, is largely of arenaceous and argillaceous character. The shales are usually a yellowish-buff or greenish to gray and have thin beds of buff-weathering sandstones. The upper limit, as defined in this report, is marked by the Keefer Sandstone, beneath which lie yellow and gray thin-bedded shales and platy

FIGURE 14

MAP


OF

GREENBRIER COUNTY

SHOWING

OUTCROPS OF

LOWER DEVONIAN ROCKS



CONTACTS, ORISKANY SERIES.

The upper contact of the Oriskany Series with that of the Marellus has already been referred to under the discussion of the latter series. In Greenbrier County these two series are apparently conformable although there is an abrupt change in the lithology of the beds at the contact. The lower limit of the Oriskany is easily recognized by the lithologic difference between the Ridgeley Sandstone and the underlying Beersville Limestone of the Helderberg. The contact appears to be conformable.

FOSSIL LIFE, ORISKANY SERIES.

The sandstones of the Oriskany in Greenbrier County contain an abundance of marine fossils. The chert proper appears to be non-fossiliferous but where sandy layers are present within the chert they generally contain marine fossils. The age relationship and therefore the fossil life of this series will be discussed at greater length under "Correlation, Oriskany Series" and "Description of Members, Oriskany Series."

CORRELATION, ORISKANY SERIES.

It has already been noted that the sandstone at the base of the Oriskany Series as found in Greenbrier County, correlates with the Ridgeley Sandstone of the Potomac region of West Virginia and Maryland. The writers are inclined to agree with Swartz¹⁰ that the Shriver Chert and the Beersville Limestone are equivalent in time of deposition. Swartz has suggested that the difference between the two in lithologic and faunal properties may be accounted for by different environments during deposition. The following quotation from Swartz¹⁰ explains why the Beersville is classified as Helderberg while its time equivalent is classified as Oriskany:

"Acceptance of the conclusions suggested above [time equivalence of Shriver and Beersville] would still leave some questions as to terminology. If Helderberg group is used primarily with a time significance, I think that the Shriver would necessarily be included in it, although admitting that the Shriver and the Beersville of the Virginia-Maryland area range above the top of the Beersville, and thus above the

¹⁰Ibid., p. 48.
Swartz, Frank McKim, The Helderberg Group of Parts of West Virginia and Virginia, U. S. G. S., Prof. Paper 158-C, pp. 47, 48; 1929.

Ridgeley Sandstone.

The Ridgeley Sandstone as found in Greenbrier County is a medium to coarse, yellowish to earthy-brown, massive sandstone varying from 12 to 20 feet in thickness. The yellow-brown color is due, no doubt, to weathering and in some places limonite is so concentrated that it approaches a low-grade iron ore. The limonite appears to be a secondary concentration and was probably derived from the weathering of pyrite. The sandstone is quite fossiliferous and since it has usually been leached of its lime content it is characteristically marked by numerous fossil pits.

In Greenbrier County as well as in Pocahontas and most of the counties to the northeast, there occurs near the top of the Ridgeley a conglomerate, composed of small quartz grains that in size and shape resemble rice or wheat. This is often called the "Wheat Grain" Conglomerate. There are numerous points at which the Ridgeley outcrops in this area but due to the ease with which it weathers only a few of these points offer clean exposures. The best of these exposures are in the vicinity of Bobs Ridge and Eckle School.

ECONOMIC ASPECTS, ORISKANY SERIES.

The Ridgeley Sandstone member weathers into a loose grained sandstone which is easily broken down into sand. This same member has been used extensively in other areas for a glass-sand. Although no sample of this member was taken for analysis, its suitability for glass-sand, as found in this area, is somewhat doubtful, as it contains a much greater amount of impurities than it does farther northeast in West Virginia.

The Huntersville Chert, standing as it does at steep angles, breaks down readily into large deposits of chert "gravel" which is excellent material for road surfacing. These deposits generally contain sufficient lime, iron, and alumina to cement readily when subjected to the crushing effect of traffic.

Both members of the Oriskany Series are proving to be major reservoirs for natural gas in some parts of the Appalachian region. The Oriskany produces oil and gas in Ohio, gas and a little oil in West Virginia, and gas in Pennsylvania

Ridgeley Sandstone.

The Ridgeley Sandstone as found in Greenbrier County is a medium to coarse, yellowish to earthy-brown, massive sandstone varying from 12 to 20 feet in thickness. The yellow-brown color is due, no doubt, to weathering and in some places limonite is so concentrated that it approaches a low-grade iron ore. The limonite appears to be a secondary concentration and was probably derived from the weathering of pyrite. The sandstone is quite fossiliferous and since it has usually been leached of its lime content it is characteristically marked by numerous fossil pits.

In Greenbrier County as well as in Pocahontas and most of the counties to the northeast, there occurs near the top of the Ridgeley a conglomerate, composed of small quartz grains that in size and shape resemble rice or wheat. This is often called the "Wheat Grain" Conglomerate. There are numerous points at which the Ridgeley outcrops in this area but due to the ease with which it weathers only a few of these points offer clean exposures. The best of these exposures are in the vicinity of Bobs Ridge and Eckle School.

ECONOMIC ASPECTS, ORISKANY SERIES.

The Ridgeley Sandstone member weathers into a loose grained sandstone which is easily broken down into sand. This same member has been used extensively in other areas for a glass-sand. Although no sample of this member was taken for analysis, its suitability for glass-sand, as found in this area, is somewhat doubtful, as it contains a much greater amount of impurities than it does farther northeast in West Virginia.

The Huntersville Chert, standing as it does at steep angles, breaks down readily into large deposits of chert "gravel" which is excellent material for road surfacing. These deposits generally contain sufficient lime, iron, and alumina to cement readily when subjected to the crushing effect of traffic.

Both members of the Oriskany Series are proving to be major reservoirs for natural gas in some parts of the Appalachian region. The Oriskany produces oil and gas in Ohio, gas and a little oil in West Virginia, and gas in Pennsylvania



PLATE XXXVIII.—Marcellus Shale showing calcareous (Onondaga?) beds, near the mouth of Slash Lick Run.



PLATE XXXIX.—Contorted Marcellus Shale, one mile northwest of Alvon.



PLATE XL.—Gullying in Marcellus Shale on east side of Coles Mountain, one mile south of Alvon.



PLATE XLl.—Healing Springs Sandstone exposed in stream bed where Howard Creek cuts through Bobs Ridge.

and New York. Most of the production is found in the Ridgeley Sandstone member but it now appears that a major gas field has been found in the Huntersville Chert in Fayette County, Pennsylvania. The chances of gas in this series in Greenbrier County will be discussed in Chapter X.

HELDERBERG SERIES.

GENERAL ACCOUNT, HELDERBERG SERIES.

The Helderberg Series, coming just below the Oriskany and being the basal subdivision of the Devonian System in the Appalachian region, is present in Greenbrier County but is thinner than it is throughout the region to the northeast. The Helderberg is essentially a limestone formation. Its lithologic character varies not only in different beds but also in the exposures of different regions. It ranges in color from light-blue to dark-gray and in texture from a massive limestone to a calcareous shale or sandstone. Although the Helderberg, in Greenbrier County, is not exposed in a manner that permits exact measurements, its thickness has been determined as approximately 300 feet.

The Helderberg Series has been extensively studied in Maryland, Virginia, and West Virginia and has generally been divisible into four members, based on both lithologic and faunal grounds. These members in descending stratigraphic order are as follows:

- Beerft Member.
- New Scotland Member.
- Coeymans Member.
- Keyser Member.

All of these members are probably present in Greenbrier County. The presence of the Beerft, New Scotland, and Keyser Members is apparently proved and although it could not be definitely identified, the Coeymans is probably represented in this area.

TOPOGRAPHIC EXPRESSION HELDERBERG SERIES.

In Greenbrier County, the Healing Springs Sandstone (New Scotland) is somewhat more resistant to erosion than is the Oriskany. This effect is plainly shown along the eastern

side of Beaver Lick Mountain where the Healing Springs is often found on the more prominent knobs. Due largely to the resistant character of the sandstone, the Helderberg Series outcrops on much of the south end of Beaver Lick Mountain and the north end of Coles Mountain.

AREAL EXTENT, HELDERBERG SERIES.

The areal extent of the Helderberg Series is shown on Figure 14 along with the Oriskany Series under the title Lower Devonian Rocks. It is also shown on Map II by a separate color in much more detail and on a larger scale. As mentioned in the foregoing paragraph this series outcrops along the eastern side of Beaver Lick Mountain, on the south end of the same mountain, on the north end of Coles Mountain, and the upper part of the series is exposed along Howard Creek and Jericho Draft.

CONTACTS, HELDERBERG SERIES.

The upper contact of the Helderberg Series with the Oriskany has already been discussed under the same heading in the description of the latter formation. The lower limit has long been the subject of many lengthy papers and discussions. In the local area conditions are not favorable for a detailed study of this question, the exposures being few and poor. In conformity with former studies of this contact, with particular reference to the northeastern counties of West Virginia, the boundary that seems best fitted is the plane between the more massive limestones of the Helderberg and the more flaggy and purer beds of the Bossardville. This division seems best adapted on both lithologic and faunal grounds, although certain species of Silurian age are found to exist on into the Helderburg.

FOSSIL LIFE, HELDERBERG SERIES.

The Helderberg Series of Greenbrier County is abundantly

CORRELATION, HELDERBERG SERIES.

Attention has already been called to the fact that all of the members of the Helderberg are represented in Greenbrier County with the single possible exception of the Coeymans member. Under "Correlation, Oriskany Series" the apparent time equivalence of the Shriver Chert and the Beeraft member has been discussed. Swartz¹ has made a regional study of the Helderberg in near-by areas and while his work does not include the Greenbrier area, the Helderberg of this area does fit his discussion nicely. The reader is referred to Swartz's paper for the more technical aspects of the correlations. In many publications and particularly the U. S. Geological Survey Folios, the Helderberg Series is included under the description of Lewistown Limestone.

DESCRIPTION OF MEMBERS, HELDERBERG SERIES.

Beeraft Member.

In Greenbrier County the Beeraft Member is a light-gray to dark bluish-gray limestone, somewhat argillaceous or arenaceous at the top, purer near the middle, and arenaceous toward the base. The limestone carries numerous nodules of black chert and silicified fossils are common. The thickness of the member appears to vary between 60 and 100 feet but areas in which it outcrops have generally been so disturbed by folding that accurate measurements are difficult to obtain. The contact of the Beeraft with the overlying Oriskany and with the underlying New Scotland appears to be transitional, with the upper contact the more distinct of the two. It is possible that the extreme lower part of the Beeraft, as herein described, carries New Scotland fossils and this possibility needs further attention from the paleontologists.

In some places the middle portion of the Beeraft is fairly pure limestone and may furnish some agricultural lime or road material. In general, however, the Beeraft is too impure for most uses and the chert nodules would probably interfere with satisfactory crushing. See Chapter XII for a further discussion of the commercial possibilities of the Beeraft.

¹Swartz, Frank McKim, The Helderberg Group of Parts of West Virginia and Virginia, U. S. G. S., Prof. Paper 158-C; 1929.

side of Beaver Lick Mountain where the Healing Springs is often found on the more prominent knobs. Due largely to the resistant character of the sandstone, the Helderberg Series outcrops on much of the south end of Beaver Lick Mountain and the north end of Coles Mountain.

AREAL EXTENT, HELDERBERG SERIES.

The areal extent of the Helderberg Series is shown on Figure 14 along with the Oriskany Series under the title Lower Devonian Rocks. It is also shown on Map II by a separate color in much more detail and on a larger scale. As mentioned in the foregoing paragraph this series outcrops along the eastern side of Beaver Lick Mountain, on the south end of the same mountain, on the north end of Coles Mountain, and the upper part of the series is exposed along Howard Creek and Jericho Draft.

CONTACTS, HELDERBERG SERIES.

The upper contact of the Helderberg Series with the Oriskany has already been discussed under the same heading in the description of the latter formation. The lower limit has long been the subject of many lengthy papers and discussions. In the local area conditions are not favorable for a detailed study of this question, the exposures being few and poor. In conformity with former studies of this contact, with particular reference to the northeastern counties of West Virginia, the boundary that seems best fitted is the plane between the more massive limestones of the Helderberg and the more flaggy and purer beds of the Bossardville. This division seems best adapted on both lithologic and faunal grounds, although certain species of Silurian age are found to exist on into the Helderburg.

FOSSIL LIFE, HELDERBERG SERIES.

The Helderberg Series of Greenbrier County is abundantly fossiliferous. A number of collections were made from these rocks and lists of the fossils identified are published in Chapter XIV. Excellent exposures for the collection of the marine fossils of the Helderberg were noted along Howard Creek in the vicinity of Bobs Ridge and at Eckle School.

PLATE XXXVI.—Portage strata in bed of Anthony Creek, 2.3 miles southwest of Neola.





PLATE XXXVII.—Portage shale and flaggy sandstone along the C. & O. Railroad at Tuckahoe.

both marine and plant fossils in Greenbrier. No exposure of this series is so complete that its entire thickness could be measured, but it is approximately 2,000 feet.

In Greenbrier County there exists no basis for a subdivision of this series since it is devoid of any lithologic changes. Paleontologically, however, the fossils collected in this area show some similarity to those in more distant areas. In Maryland, Dr. Swartz¹ has divided those beds lying between the Chemung and Genesee, which correspond to the Portage, as follows:

Parkhead Sandstone Member. Recurrent *Tropidoleptus carlnatus* fauna.

Shale beds.

Conglomeratic sandstone beds.

Cyclonemina multistriata zone.

Camarotoecchia congregata var. *parkheadensis* zone.

Liorhynchus mesacostale zone.

Woodmont Shale Member.

Beds containing *Ithaea* fauna. (*Spirifer mucronatus* var. *posterus* fauna).

Liorhynchus globuliforme zone.

Cladochonus—*Reticularia laevis* zone.

Beds containing the Naples fauna. (*Buchiola speciosa* fauna).

As noted in preceding paragraphs lithologic characteristics that would warrant any subdivision in this area are absent, there being a monotonous succession of shales and flaggy sandstones, with no occurrence of conglomeratic beds. An examination of the fossils, however, reveals a similarity of the fauna of the upper half of this series to the Parkhead fauna while the lower half retains fossils characteristic of the Naples.

TOPOGRAPHIC EXPRESSION, PORTAGE SERIES.

The topography formed by the Portage Series is, in general, much like that of the Chemung, except less severe. Due to its less resistant character the ridges and slopes are more gentle and not so high. Where the strata are not greatly disturbed the more sandy ledges of the Chemung form steep ridges which are paralleled by the more gentle slopes of the Portage.

¹Charles K. Swartz, Middle and Upper Devonian, Md. Geol. Surv., p. 411; 1913.

AREAL EXTENT, PORTAGE SERIES.

The areal extent of the Portage Series is included in Figure 13 under the heading Upper and Middle Devonian Rocks, on page 298. There are only two long outcrops, both of which are on the eastern side of the Greenbrier River and enter the county on the north on either side of Browns Mountain Anticline and parallel this structural feature to a point about one mile southwest of White Sulphur Springs where they join on this plunging anticline to pass beneath the Chemung Series about 3 miles above the mouth of Harts Run.

CONTACTS, PORTAGE SERIES.

The contact of the Portage Series with the overlying Chemung has already been discussed under the same heading in the description of that series on page 302. At the base of the Portage its contact with the Genesee is generally marked by a change from olive and greenish-gray shales and flagstones of the former to that of brown, or black and sandy, usually fissile, and sometimes slaty shales of the latter, which contain no sandstone flags; and also by the presence of typical Genesee fossils.

FOSSIL LIFE, PORTAGE SERIES.

The Portage Series throughout West Virginia to the northeast has generally been found to carry few fossils except in the Eastern Panhandle where they are fairly abundant. In southern West Virginia this series contains only infrequent fossils. In Greenbrier County fossils were noted at frequent points and several collections were made, although a thorough search was not attempted. Several species of marine fossils and impressions of plants are listed in Chapter XIV, under collections from this series.

CORRELATION, PORTAGE SERIES.

The relationship of the Portage Series of Greenbrier County to its more northeastern counterparts in other States has already been touched upon under the subject of "General Account." Owing to a lack of any apparent lithologic subdivisions, and to the absence of definite faunal subdivisions,

it is inadvisable to make detailed comparisons in this report. Attention has, however, been called to the presence of Naples fauna in the lower portion and of Parkhead fossils in the upper.

The Portage Series occupies the interval between the Genesee and Chemung members of the Jennings Formation of the U. S. Geological Survey.

ECONOMIC ASPECTS, PORTAGE SERIES.

The Portage Series contains neither precious metals nor any other products of present economic interest. The shales are too sandy for brick purposes and the sandstones are too thin for building stone, and also weather into small blocks too small for flagstone walks. The soil is thin and poor except along the bottoms, and here the soil has been carried in and impregnated with that from other series. The soil does seem suitable for timber growth.

GENESEE SERIES.

GENERAL ACCOUNT, GENESEE SERIES.

The Genesee Series, coming just below the Portage and being the basal group of the Upper Devonian, is made up of black, fissile, argillaceous shales, with occasional streaks of bluish-black limestone, followed by dark but more arenaceous beds. These beds are followed by a greenish-gray arenaceous shale with occasional thin sandstone bands. In physical appearance the Genesee resembles the Marcellus but on close examination exhibits a difference, being harder, more arenaceous, and having a slaty cleavage. The thickness of the Genesee varies from 50 to 100 feet and may be even greater, but complete exposures are not available for accurate measurement.

The Genesee Series has not generally been subdivided, being considered as an individual lithologic unit. In Grant County, however, Prouty^{*} recognizes two divisions of this series, a lower and black, argillaceous and carbonaceous shale,

^{*}W. F. Prouty, Hampshire and Hardy Report, W. Va. Geol. Surv., pp. 323-324; 1927.

and an upper portion of more arenaceous and thin-bedded sandstone. On detailed examination these general divisions are noted in Greenbrier County.

TOPOGRAPHIC EXPRESSION, GENESEE SERIES.

The Genesee Series, in conjunction with the underlying shales of the Hamilton and Marcellus, is usually found in comparatively narrow valleys or lowlands. Its upper portion is more sandy, is slightly more resistant, and forms a gentle sloping topography between the Middle Devonian shales and the overlying Portage Series.

AREAL EXTENT, GENESEE SERIES.

On Figure 13, the areal extent of the Genesee is included under that of the Upper and Middle Devonian Rocks, but is delineated on Map II in much greater detail and on a larger scale. Its thickness is so small in comparison to that of the Upper Devonian that its area of outcrop is very limited. Its exposures are confined to the area east of the Greenbrier River, and limited to two narrow outcrops paralleling either side of the Browns Mountain Anticline from the Pocahontas County line to a point one mile southwest of White Sulphur Springs where they unite on the southern end of this structural fold to pass beneath the Portage Series.

CONTACTS, GENESEE SERIES.

The upper contact of the Genesee with the Portage Series has already been discussed in connection with the latter series on page 306. At its base it rests upon the Hamilton Series which is poorly exposed in Greenbrier County. For some time the writers were uncertain whether or not the Hamilton Series was present at all, but certain collections (Nos. 51 and 55), made from a brown, arenaceous and calcareous shale at points where this series should occur, contain characteristic Hamilton fossils. The lower contact is therefore placed at the base of the black, carbonaceous, fissile shale, with thin limestones, containing a Genesee fauna and at the top of a brown arenaceous shale with a sparse Hamilton fauna.

FOSSIL LIFE, GENESEE SERIES.

The most common fossils in the Genesee are pelecypods, cephalopods, and pteropods. The most abundant species are: *Paracardium doris*, *Pterochaenia fragilis*, *Buchiola livoniae*, *Styliolina fissurella*, and *Baetrites aciculus*. These species apparently range through the series. Several collections were made, the identifications of which were made by Dr. John L. Tilton and these appear in Chapter XIV.

CORRELATION, GENESEE SERIES.

The Genesee of Greenbrier County retains the same general character, both lithologic and faunal, as this same member of the Jennings Formation of New York, which is the type locality of the Genesee, and can definitely be correlated with it. It has been recognized and described in other Appalachian counties of West Virginia as well as in Maryland, Pennsylvania, and New York.

DESCRIPTION OF MEMBERS, GENESEE SERIES

As previously stated the Genesee is generally considered as a single unit with no distinct faunal break by which it might be subdivided. Even though there is a gradual change from predominantly shaly material at the base to sandy beds at the top the transition from one to the other is not sufficiently abrupt to warrant further subdivision. This is also further emphasized by the range of typical Genesee fossils throughout.

ECONOMIC ASPECTS, GENESEE SERIES.

From an economic standpoint the Genesee is of minor importance. It is possible that a portion of these shales would prove suitable for building brick or as a flux with limestone for the manufacture of Portland cement. These shales have frequently been prospected for coal but so far as known no coal has ever been found associated with them. The soils from these weathered shales are usually quite thin and barren and unsuitable for cultivation. The more sandy portions make excellent road-surfacing material where more durable stone is not available.

MIDDLE DEVONIAN ROCKS.

GENERAL STATEMENT.

The Middle Devonian Rocks, as indicated by the classification adopted for the Devonian in Greenbrier County, includes beds of Hamilton, Marcellus, and Onondaga age and rocks of the same age have been grouped under the name **Romney** by the U. S. Geological Survey and others. Rocks of this age have a combined thickness, in Greenbrier County, of approximately 500 feet. At all points observed these rocks are intricately folded and mashed so that accurate measurements, either in whole or in part, are not possible.

In many places it is very difficult to separate the Hamilton from the Marcellus and the outcrops of the two series are shown together on Map II under the name of Marcellus. In Greenbrier County, as well as in many other counties of West Virginia, the lower part of the Middle Devonian carries a mixed Marcellus and Onondaga fauna and this portion is considered to be the equivalent of the Onondaga of New York.

It has been found to be impractical to attempt a detailed subdivision of the Middle Devonian Rocks of Greenbrier County and it is to be remembered that the areas shown as Marcellus on Map II contain beds that are Hamilton, Marcellus, and Onondaga. In a similar manner the Middle Devonian Rocks are described under the Marcellus Series on the following pages.

MARCELLUS SERIES.

GENERAL ACCOUNT, MARCELLUS SERIES.

The Marcellus Series, coming below the Genesee and above the Oriskany, is composed for the most part of black, fissile shale, which becomes flaky and slickensided on compression. These shales are so black and contain so much carbon that they are frequently prospected for coal. Because of this carbon content they have a tendency to weather light colored on exposure. Toward the base of this series there occur thin impure limestones along with calcareous shales. At many localities large concretionary and septarian nodules of ferru-

ginous and calcareous character are common and these concretions often contain considerable barite. In Greenbrier County, the Marcellus Series is confined to the area comprising the Browns Mountain Anticline, and has therefore been subjected to considerable pressure by folding. For this reason it is impossible to get the exact thickness in any of the exposures visited, because of the repetition of beds by minor folding or thinning due to lateral compression, but the Marcellus retains, in this area, an approximate thickness of 500 feet.

TOPOGRAPHIC EXPRESSION, MARCELLUS SERIES.

The Marcellus shales are the most easily eroded series of rocks exposed in Greenbrier County. The low valleys on either side of the Beaver Lick-Coles Mountain area are largely formed in this series, as well as the flat land around White Sulphur Springs. These bottoms are frequently covered by alluvial material.

AREAL EXTENT, MARCELLUS SERIES.

On Figure 13 the Marcellus Series is included under the Upper and Middle Devonian Rocks, but it can be seen in much greater detail and on a larger scale on Map 11. This series is also confined to the east side of Greenbrier River, and to the Browns Mountain Anticline. It enters the county from the north on either side of this complex folded area and parallels this structural feature to the vicinity of White Sulphur Springs where its outcrop broadens by minor folding and passes beneath the younger rocks. The Marcellus Series can be seen to good advantage at many points along its outcrop. Along the highway on either side of Coles and Beaver Lick Mountains many opportunities are afforded to examine these rocks.

CONTACTS, MARCELLUS SERIES.

The upper contact of the Marcellus, as herein defined, with the overlying Genesee, has already been discussed under the description of the latter series. At the base the contact is more pronounced, with the black, fissile, typical Marcellus shale resting upon a yellowish-gray or greenish sandstone or

where this sandstone is absent, upon a yellowish to dark, sandy chert. The sandstone and chert are of Oriskany age, a fact that will be described in more detail under the description of the Oriskany Series. Although the contact at the base of the Marcellus is quite distinct there is no concrete evidence of an unconformity.

FOSSIL LIFE, MARCELLUS SERIES.

The Marcellus Series is, as a whole, sparingly fossiliferous. Aside from fossils occurring in the calcareous zones of the lower part and in the occasional brown shale at the top, the life forms are limited to a few species. *Styliolina fissurella* is the most common, with *Liorhynchus limitare* and a few other forms occasionally found. Since the fossil collections were made primarily for stratigraphic mapping, and as the Marcellus is generally followed with slight difficulty because of its lithologic character, few collections were made from this series. At the top of the series the brown shales interfinger with the black shales and two collections from this portion of the Middle Devonian show typical Hamilton forms. In the lower part, lenticular black limestones carry a mixed Marcellus and Onondaga fauna.

CORRELATION, MARCELLUS SERIES.

In view of the foregoing discussion it is clear that the Middle Devonian of Greenbrier County is the equivalent of the Hamilton and Marcellus Series as described in other counties of the State. The upper part has a lithology that is in part similar to the Hamilton of other areas and contains some black shale of the character typical of the Marcellus. The lower portion carries a mixed Marcellus and Onondaga fauna, a relationship that is well recognized in the Allegheny area⁶.

DESCRIPTION OF MEMBERS, MARCELLUS SERIES.

As described in the foregoing discussion, it is not feasible to subdivide the Middle Devonian in Greenbrier County. In the counties to the northeast it is possible to differentiate be-

⁶See, Kindle, E. M., Onondaga Fauna of the Allegheny Region, U. S. Geol. Sur., Bull. 508; 1912; see also, Prosser, C. S., Kindle, E. M., and Swartz, C. K., The Middle Devonian Deposits of Maryland, Maryland Geol. Sur., 1913.

tween the various lithologic and paleontologic units as described by Price⁷ in Pocahontas County. While similar subdivisions might be made in the northern part of Greenbrier County, they can not be carried the full length of the outcrop of the Middle Devonian.

The **Lower Selinsgrove (Onondaga) Limestone** and **Lower Selinsgrove Shale** of White⁸ are represented in Greenbrier County but as the limestone merges into typical Marcellus shale it can not always be recognized.

ECONOMIC ASPECTS, MARCELLUS SERIES.

The Marcellus Series weathers into a gray plastic clay soil which in itself is poor for cultivation, but is generally enriched by a wash from the adjoining hills, and locally by the presence of the Lower Selinsgrove (Onondaga) Limestone. The local limestones, while comparatively pure, are too thin for commercial purposes, their greatest value being in addition of lime to the soil in situ.

The Marcellus shales have a comparatively high carbon content from which various petroleum products may be distilled. No prospecting was done for oil shales in Greenbrier County, in the preparation of this report, but a sample was collected by the senior author from this series in Hardy County, and distilled in the Chemical Engineering laboratory at West Virginia University, which showed the presence of both oil and gas in these shales. Their value for this purpose will need to have further investigation at some future date. These shales have frequently been prospected for coal but so far as known no coal has ever been found associated with them and it is likely that none will ever be found at this horizon in Greenbrier County.

LOWER DEVONIAN ROCKS.

GENERAL STATEMENT.

The Lower Devonian Rocks, composed of the Oriskany and Helderberg Series, are represented in Greenbrier County by limestones, sandstones, and chert, having a total thick-

⁷Price, Paul H., Pocahontas Report, W. Va. Geol. Sur., pp. 221-230; 1929.

⁸White, I. C., Report G-7, Sec. Geol. Sur. of Pa., pp. 79-81; 1883.

ness of approximately 400 feet. A discussion of these beds will appear on succeeding pages. Figure 14 shows the distribution of the Lower Devonian Rocks in the county, while on Map II the same information is shown in much greater detail and on a larger scale.

ORISKANY SERIES.

GENERAL ACCOUNT AND SECTION, ORISKANY SERIES.

The Oriskany Series, which forms the upper subdivision of the Lower Devonian Rocks, is represented in Greenbrier County by a gray or brown, massive, coarse, fossiliferous sandstone at the base, by a gray and dark chert and a thin yellowish- or greenish-gray fossiliferous sandstone at the top. The sandstone at the base of the series generally contains in its upper part a bed of small quartz pebbles which resemble rice or wheat grains and this bed has often been referred to as the "Wheat Grain" Conglomerate. In some places this sandstone contains pockets of limonite (iron ore) with traces of manganese. The ore is apparently of secondary origin and is not everywhere present.

The series has been divided into two members on the basis of their lithologic characteristics. The Huntersville Chert, first named and described in Pocahontas County, is the upper member and its outcrop in West Virginia is apparently confined to Pocahontas, Greenbrier, and part of Pendleton Counties^{2a}, its occurrence in the latter county being only recently discovered by Price. The lower member, the Ridgeley Sandstone, makes an excellent lithologic unit and has been traced south across the State from its type locality in Maryland. Although varying in thickness, its general character, both lithologic and faunal, is retained throughout its outcrop in West Virginia. The Shriver Chert, which is described as the basal member of the Oriskany in the Potomac region of West Virginia, was not recognized in Greenbrier County and its apparent absence will be discussed in more detail under "Cor-

^{2a}Since the above was written, two feet of Huntersville Chert has been found in Grant County, along State Route 42, 1.6 miles south of Seherr.

relation, Oriskany Series" and "Correlation, Helderberg Series" on subsequent pages. The following generalized section of the Oriskany has been compiled for Greenbrier County:

General Section of Oriskany Series for Greenbrier County.

	Thickness. Feet.	Total. Feet.
Oriskany Series		
Sandstone, classlled with Huntersville Chert, yellowish or greenish-gray, fine- to medium-grained, calcareous, usually contains abundant glauconite, contains marine fossils.....	8 to 0	0
Chert, Huntersville, gray to black, hard, tough, bedded, contains occasional layers of fine-grained, glauconitic sandstone, weathers to light-gray "gravel".....	60 to 70	70
Sandstone, Ridgeley, gray, calcareous, stained brown on weathering, medium- to coarse-grained, usually contains small quartz pebbles resembling rice or wheat grains near top, marine fossils.....	12 to 20	90
Helderberg Series (Becraft).....		

TOPOGRAPHIC EXPRESSION, ORISKANY SERIES.

Due to its massive, cherty, and sandy character the Oriskany Series is generally found making a bold topography. Most of Bobs Ridge and Coles Mountain are covered with this series and the outcrop of the Oriskany makes bold shoulders or "knobs" paralleling Beaver Lick Mountain.

AREAL EXTENT, ORISKANY SERIES.

On Figure 14, the outcrops of the Oriskany Series are shown along with the underlying Helderberg Series, under the title Lower Devonian Rocks. On Map II the outcrops of this series are shown in greater detail and on a larger scale. The best exposures for study of the Oriskany Series are to be found where Howard Creek cuts through Bobs Ridge and on Jericho Draft near Eckle School.

FOSSIL LIFE, CATSKILL SERIES.

The typical Catskill of Greenbrier County appears to be devoid of fossil fauna and the fossil flora are rare and poorly preserved. No fossil collections were made from this series in this county.

CORRELATION, CATSKILL SERIES.

It is evident from the foregoing discussion that the Catskill Series as found in Greenbrier County correlates, at least in part, with this same series in the other counties of this State where it has often been designated as **Hampshire Formation** by members of the U. S. Geological Survey.

ECONOMIC ASPECTS, CATSKILL SERIES.

From an economic standpoint the Catskill Series is of minor importance in Greenbrier County. The soils are generally best suited to timber growth and its sandstones are, as a rule, not suitable for use as building stone. Its shales could possibly be used for making brick or tile but materials of this type are widely distributed and quite common in Greenbrier County.

CHEMUNG SERIES.

GENERAL ACCOUNT AND SECTION, CHEMUNG SERIES.

The Chemung Series of the Upper Devonian, coming just below the Catskill Series and just above the Portage Series, comprises the largest single assemblage of beds in Greenbrier County. It is composed of a mass of interbedded sandstones ranging from flags to massive ledges, alternating with green, olive, and brown shales, and it attains a thickness of 3,000 feet. The sandstones, which are greenish-gray to brown, fine-grained, and micaceous, very hard and compact, and often lenticular, occur throughout the series.

Owing to the lithologic similarity throughout the Chemung, attempts to subdivide it by physical appearance have been rather unsuccessful. The Hendricks Sandstone is apparently present in Greenbrier County and its presence just

beneath the red shales of the Catskill provides a valuable field marker. This sandstone is designated as marking the top of the *Chemung Series*. Near the middle of the series the sandstones become massive and sometimes contain conglomeratic beds. In the lower half of the series thin beds of limestone composed entirely of shells of marine animals are found and marine fauna and land flora are present at various horizons throughout the series. The following is a generalized section of this series in Greenbrier County:

General Section, Chemung Series, Greenbrier County.

	Thickness. Total.	
	Feet.	Feet.
1. Sandstone, Hendricks , grayish-brown, weathering white, flattened quartz pebbles and with occasional plant and marine fossils.....	10- 50	50
2. Shales, sandy, green to brown, with some sandstones; ripple-marked beds common.....	200- 400	450
3. Sandstone zone, with some shales, beds brown and iron-stained on weathering, frequently green on fresh exposure.....	100- 150	600
4. Shale zone, with some sandstones, gray and green, sandstones flaggy, shales sandy.....	800-1000	1600
5. Sandstone zone, with some shales, sandstones generally thick-bedded, greenish to reddish-brown, shales sandy, olive or gray.....	290- 400	2000
6. Shales, alternating with sandstones, shales olive or gray, sandstones greenish-brown, thinner bedded toward base.....	600-1000	3000

TOPOGRAPHIC EXPRESSION, CHEMUNG SERIES.

In Greenbrier as well as the remaining West Virginia Counties to the northeast, the Chemung Series, where unaffected by superjacent rocks or structural disturbances, exhibits a topographic relief that is characteristic of this series. The usual topography is that of sharp, narrow ridges with a general profile like that of an inverted V, separated by normal V-shaped valleys. When steeply dipping, this series forms a row of knobs or ridges parallel to the mountains formed by the overlying rocks, as well as to the valleys formed by the underlying and less resistant Middle Devonian shales. (See Plate III). The Chemung Series can be followed across the

State to the northeast, in Pocahontas, Randolph, Pendleton, Tucker, Mineral, Grant, Hardy, and Hampshire Counties where it forms these characteristic rows of sharp knobs and ridges.

AREAL EXTENT, CHEMUNG SERIES.

On Figure 13, page 298, the Chemung Series is outlined along with the remainder of the Upper and Middle Devonian Rocks, and comprises a larger areal extent than the Catskill, Portage, and Genesee Series combined. On Map II the outcrop of this series is shown in much greater detail and on a larger scale. The surface exposures of this series are limited to the eastern portion of the county and lie entirely east of the Greenbrier River. Along Allegheny Mountain the Chemung Series is extensively exposed and forms the greater part of this mountain, the younger Catskill and Pocono Series being retained along the crest at occasional high points. The remaining and longest continuous single exposure lies east of the Greenbrier River throughout the entire length of the county and west of those mountains included in the Browns Mountain Anticline.

CONTACTS, CHEMUNG SERIES.

The contact of the Chemung with the overlying Catskill Series has already been discussed under the description of the latter series, page 299. At the base of the Chemung or at its contact with the Portage Series, the sedimentary record is not clear. There is, however, a rather noticeable change, both lithologic and faunal, between those beds which are typical Portage and those which are Chemung. The former series is predominantly shaly and generally sparing in fossils, with flaggy or platy sandstone members which weather into rectangular blocks. The latter series contains sandstones which are much more massive, and also contains numerous marine horizons, with the guide fossil *Spirifer disjunctus* in profusion. As has been the policy of the West Virginia Geological Survey, the contact of these two series is therefore placed at the point where the flaggy and platy sandstone of the sparingly fossiliferous Portage is succeeded by the more massive sandstones, and abundantly fossiliferous Chemung. Because of the variation in the sandstones a decided break in the topography is often noted which is of great help in areal mapping.

FOSSIL LIFE, CHEMUNG SERIES.

Throughout the limits of Greenbrier County the Chemung Series carries marine fossils in profusion and at several places fossil land plants were noted. Although no attempt was made to obtain a complete fossil record, numerous collections were made from this series. *Lists of the fossils identified from these collections were made by the late Dr. John L. Tilton and Prof. Dana Wells and these lists are published in Chapter XIV.* The guide fossil **Spirifer disjunctus** is probably the most conspicuous and abundant form but **Spirifer mesacostalis** and **Atrypa hystrix** are quite common.

CORRELATION, CHEMUNG SERIES.

From the above discussion it is evident that the Chemung Series of Greenbrier County correlates with the same series in New York, Pennsylvania, and Maryland and it retains the same lithologic and faunal characteristics. This series has often been described along with the Portage and Genesee Series under the term **Jennings Formation**.

DESCRIPTION OF MEMBERS, CHEMUNG SERIES.

Hendricks Sandstone.

The **Hendricks Sandstone**, comprising the upper member of the Chemung Series and marking the lower limit of the Catskill Series in Greenbrier County, was observed at several points throughout the area. It is generally grayish-brown to reddish-brown, massive, and contains numerous flattened quartz pebbles. It is frequently white on weathered surfaces, occasionally contains marine fossils along with fragments of plants and varies in thickness from 10 to 50 feet.

As noted under "General Account and Section, Chemung Series," there is little upon which to base divisions of the Chemung Series. It appears probable that No. 3 of the General Section, page 301, may in general represent the **Valley Head Sandstone** and No. 5 of the same section may represent the **Elkins Sandstone**. Both of the sandstones mentioned were first named and described by Reger³.

³Reger, David B., The Tygart Valley Devonian Trees of West Virginia, Am. Jour. Sci., Vol. XV; pp. 52 and 53; Jan., 1928.

ECONOMIC ASPECTS, CHEMUNG SERIES.

From an economic standpoint the Chemung Series is of minor importance. The sandstone members are generally too cross-bedded, or shaly and sometimes quartzitic to be used for building stone, while the shales are too sandy for brick or tile purposes. There is a possibility that some of the sandstones from this series would be suitable for grindstones. Many of the flags of this series are suitable for flagstone walks, the demand for which is now on the ascendency, the chief objection being, of course, their distance to market. These sandstone flags have been used rather extensively in roads, walls, culverts, walks, etc., by C. C. C. workers.

The shales weather to a thin, yellow soil, quite poor in fertility, so that their use for agricultural purposes is not extensively followed. In the area of this report the outcrops of this series seem well adapted to timber growth.

This series so far as known contains no minerals of value except in regions farther west, although its frequent pockets of iron pyrites have often caused it to be prospected for gold in mountain counties, with invariably disappointing results. To the west, southwest, and northwest, where it is deeply buried under younger rocks, there are rich deposits of oil and gas in some of its coarser members. East of the Greenbrier River there is no possibility of their presence, as these horizons appear at the surface. West of the Greenbrier River the chances of obtaining oil or gas from this series are very slight, as will be discussed in Chapter X under Petroleum and Natural Gas.

PORTAGE SERIES.

GENERAL ACCOUNT, PORTAGE SERIES.

The Portage Series of the Upper Devonian, coming just below the Chemung and just above the Genesee, is composed of a succession of shales and sandstones, both of which are generally greenish-gray in color. The shales predominate but slightly, and are usually arenaceous. The sandstones are rather compact, fine-grained, hard, and flaggy, and vary from 2 to 6 inches in thickness. This series was found to contain

PLATE XXX.—Sandstone conglomerate (Berea?) at or near the base of the Pocono near Anthony.





PLATE XXXI.—Giant ripple-marks in basal Pocono conglomerate on Meadow Creek, 2.6 miles southeast of Neola.




PLATE XXXII.—Natural whelstones formed by jointing in Chenung sandstone on Kates Mountain.






PLATE XXXIII.—Close folding in Portage strata along Anthony Creek, north of Neola.






PLATE XXXIV.—Close folding in *Portage strata* along Anthony Creek, north of Neola.





PLATE XXXV.—Interfingering folds in Portage strata, north of Neola.

PLATE XXVIII.—Broad Ford Sandstone in C. & O. Railroad cut at Caldwell.



PLATE XXIX.—Basal Pocono sandstone conglomerate (Berea?) along Midland Trail (U. S. Route 60) 0.9 mile east of
adwvll.



assume certain aspects which Dr. Girty considers to be of a Devonian character, based on his study of the collections made by the writer (Reger) at these localities."

In Pocahontas County, Paul H. Price has made a number of collections of fossils from the Pocono and as reported by him⁴⁸, these fossils were identified by Dr. John L. Tilton, who considered them to be Mississippian. In his discussion of some of the fossils, Tilton remarks on the "wonderfully Chemung-like fossil assemblages."

Several collections of Pocono fossils were made in Greenbrier County in connection with the field work for this report and the fossils identified are listed in Chapter XIV. Although the fossils collected in Greenbrier County were not perfect nor complete specimens, they would have been unquestionably identified as Mississippian forms, if the Mississippian age of the entire Pocono Series had not been in doubt.

As reported by Reger⁴⁹, a collection of Pocono plants was made by Reger, Price, and Dr. David White, 2.2 miles southwest of the highway bridge across the Greenbrier River at Ronceverte, on the south side of the river, at an elevation of 1800' B. The collection was turned over to Dr. White and so far as known to the writers, no identifications of the fossils in the collection have been made.

CORRELATION, POCONO SERIES.

The Pocono Series as defined in Greenbrier County is plainly of the same general age as the beds described under the same series in other counties of West Virginia as well as the adjoining States of Maryland and Pennsylvania on the north, even though considerable change in conditions of deposition has taken place. Beds of the same apparent age, however, in southwestern Virginia and northeastern Tennessee have been described under such titles as Price Formation and Grainger. Reger⁵⁰ is of the opinion that the New Providence Group of Kentucky is of the same age as the Pocono, which was earlier pointed out by Butts in a discussion of the Mississippian Series of eastern Kentucky.

⁴⁸Price, Paul H., Pocahontas Report, W. Va. Geol. Sur., pp. 379-383; 1929.

⁴⁹Reger, David B., Mercer, Monroe, and Summers Report, W. Va. Geol. Sur., p. 511; 1926.

⁵⁰Ibid., p. 512.

DESCRIPTION OF MEMBERS, POCONO SERIES.

In some parts of Greenbrier County a lenticular sandstone is present immediately below the red Maecrady shales, that is regarded as marking the upper boundary of the Pocono Series. This sandstone is usually gray or brown in color, platy, and shaly, ranging in thickness from 0 to 66 feet as shown in the generalized section published on a preceding page.

MERRIMAC COAL.

In Greenbrier County, a lenticular coal was noted in the upper part of the Pocono that is believed to correlate with the **Merrimac** or "**Big Seam**" of Montgomery County, Virginia, where it has been mined on a commercial scale for several years. A great deal of time, energy, and money has been spent in prospecting this coal in Greenbrier County, with but little success. Although the occurrence of the seam is of great scientific interest, it does not appear to attain sufficient thickness, regularity, and purity in Greenbrier County to be of commercial value and further prospecting of this horizon should be discouraged.

Near Hokes Mill in southern Greenbrier County and adjoining parts of Monroe County, several coal test borings were drilled to test this coal and the results were very disappointing. The records of these borings (Nos. 16, 17, 18, and 19) are published in Chapter XI. The correlations shown in the records of these borings were determined by Mr. David B. Reger and it is noted that he recognized several beds such as "Squaw Sandstone," "Lindside Sandstone," and "Langhorne Coal." Since no method has been found for definitely identifying these beds on the surface in Greenbrier County, the correlations of these beds are not carried into the other parts of the county.

The following exposures of Merrimac Coal were noted in Greenbrier County:

Coal Exposure—No. 503 on Map II.

On west side of public road, 0.8 mile north of Hokes Mill; **Merrimac Coal**; elevation, 1640' B.

	Ft.	In.
Coal	1	6

Reger⁵¹ collected a sample (No. 636R) of coal near the above exposure and its chemical analysis is published under **No. 503** in the Table of Coal Analyses at the end of Chapter XI.

W. A. Napier Mine No. 1 (Abandoned)—No. 504 on Map II.

On west side of Greenbrier River, 2 miles north of Caldwell and 0.3 mile northeast of Coalbank School; Merrimac Coal; elevation, 1860' B.

	Ft.	In.
1. Sandstone, gray to brown, hard, micaceous, 10' to.....	15	0
2. Coal, impure, lenticular, 0' to.....	4	0
3. Sandstone, shaly.....	5	0

A sample (No. 77PH) was taken from No. 2 of the above section and its chemical analysis is published under **No. 504** in the Table of Coal Analyses at the end of Chapter XI. The above mine was operated for a time in 1928 and an estimated 150 tons of coal was removed.

Coal Exposure—No. 505 on Map II.

Along public road, 1.3 miles northeast of Julia and 0.7 mile northwest of Rorer; Merrimac Coal; elevation, 2250' B.

	Ft.	In.
Coal blossom.....	1	6

Coal Exposure—No. 506 on Map II.

Along public road, 0.85 mile northeast of Rorer; Merrimac Coal; elevation, 2470' B.

	Ft.	In.
Coal blossom, thickness not determined.....

Coal Prospect—No. 507 on Map II.

On west side of Greenbrier River, 0.95 mile east of Alum Springs and 0.7 mile west of Judyton, P. O., (Kelster Sta.); Merrimac Coal; elevation, 2085' B.

	Ft.	In.
1. Sandstone, cross-bedded, lenticular, with plant fossils and coal streaks.....	10	0
2. Coal, irregular, impure, 6 inches to.....	1	6
3. Shale, black, carbonaceous, fissile, thin-bedded, with plant fossils.....	10	0

⁵¹Reger, David B., Mercer, Moaroe, and Summers Report, W. Va. Geol. Sur., p. 516; 1926.

A sample (No. 98PH) was collected from No. 2 of the above section and its chemical analysis is published under **No. 507** in the Table of Coal Analyses at the end of Chapter XI.

Floyd Childers Coal Prospect—No. 508 on Map II.

Monroe County; near Greenbrier County line, 1.55 miles southeast of Salem Church; land formerly known as "Williams Place"; A. Bell Hoke owns mineral rights; **Merrimac Coal**; elevation, 2350' B.

	Ft.	In.
Coal, fallen shut, thickness reported.....	1	6

A sample (No. 102PH) was collected from the dump of the above prospect and its chemical analysis is published under **No. 508** in the Table of Coal Analyses at the end of Chapter XI.

A study of the analyses of the Merrimac Coal as published in the Table of Coal Analyses at the end of Chapter XI, together with the detailed exposures and prospects herein exhibited, indicates that little hope of finding valuable coal in this horizon can be entertained in Greenbrier County. The coal is so irregular in occurrence, so impure and thin, and so disturbed by folding that it could hardly be seriously considered as a commercial deposit and it is quite doubtful whether attempts to use it for local domestic purposes will ever be successful.

BROAD FORD SANDSTONE.

The **Broad Ford Sandstone**, coming near the top of the Pocono, is one of the prominent members of this series in Greenbrier County and is well exposed for many miles along the Greenbrier River. The lateral streams that flow into the main river have cut deep V-shaped valleys through the Pocono Series and now offer many excellent exposures of the Broad Ford member. This sandstone was named by Reger⁵² from its exposure near the village of Broad Ford at the line between Smyth and Tazewell Counties, Virginia.

In Greenbrier County this division of the Pocono Series is largely a sandy deposit, being massive in the upper part, but often split into benches, with the lower part becoming quite shaly. It is generally reddish-brown to gray, micaceous

⁵²Ibid., pp. 520-525.

feruginous, and has an upper bench which weathers into large concentric boulders, a characteristic that is traceable across southern West Virginia. It usually contains several zones of marine fossils, but in the localities where collections were made the fossils were so badly weathered that complete identifications were not possible. The Broad Ford, as well as the greater part of the Pocono Series, decreases in thickness to the northwest, and hence has its best development in the central and southern portions of the county. Along the Chesapeake and Ohio Railway, near the Greenbrier-Pocahontas County line, this sandstone is quite massive and forms steep precipitous cliffs west of the Greenbrier River. Its thickness, character, and stratigraphic position are shown in the Caldwell, Cold Knob—Hinkle Well, Spring Creek—North, and Spring—South Sections as published in Chapter V.

Certain portions of the Broad Ford Sandstone are suitable for building material and have been used for that purpose at several points in the county. The stone used in the construction of many of the Chesapeake and Ohio Railway bridges was quarried from this stratum.

In the general section which appears earlier in this chapter, a stage of variegated shales and flaggy sandstones is noted coming between the Broad Ford and Berea Sandstones. It is possible that this succession of beds should be included in the Broad Ford Sandstone. If this were done, however, some more inclusive term, such as Formation, would be necessary to properly designate it.

BEREA SANDSTONE.

In Greenbrier County, as in the counties to the north and south, the base of the Pocono Series is marked by a medium- to coarse-grained sandstone that is usually conglomeratic. This stratum has been termed the **Berea Sandstone** in the reports on adjoining counties, and that name is retained in this report.

The **Berea Sandstone** or **Berea Grit** was first named by Newberry⁵³ from its occurrence near the town of Berea in northeastern Ohio, where it has been quarried extensively. The Mississippian age of the Berea in Ohio has not been ques-

⁵³Newberry, John S., Report of Progress in 1869, pt. 1, pp. 21, 22 and 29, Ohio Geol. Sur.; 1870.

tioned and if the Pocono of southeastern West Virginia is Mississippian it is quite probable that its basal sandstone does correlate with the Berea of Ohio.

The character, thickness, and stratigraphic position of the Berea Sandstone are shown in the Caldwell Section, published in Chapter V, in the generalized section in this Chapter, and its appearance is well illustrated on Plates XXIX, XXX, and XXXI.

ECONOMIC ASPECTS, POCONO SERIES.

From an economic standpoint the Pocono Series is of minor importance, there being no coals of minable thickness, and the sandstones producing a soil that is better fitted for timber growth than for cultivation. As noted under the description of that member, the Broad Ford Sandstone is, in some localities, suitable for heavy masonry and has been used locally for that purpose. The shales are generally too sandy for brick or tile manufacture. Farther west in the State this series often holds large quantities of both oil and gas, the character of these strata being such as to make excellent reservoirs for their retention. In this county, however, there is little hope of finding either oil or gas in these rocks, as any of the lighter hydrocarbons that may have once existed in them has been permitted to escape, on account of their frequent exposure above drainage. A further discussion of oil and gas possibilities will be found in Chapter X.

CHAPTER VIII.

STRATIGRAPHY—DEVONIAN ROCKS.

GENERAL STATEMENT.

The rocks comprising the Devonian Period in Greenbrier County retain, in general, the same characteristics as found in New York and other northern Appalachian States, so that the generally accepted standard column of New York will be followed in this report. It is true that certain minor subdivisions have disappeared, while other members have considerably decreased in thickness, but at the same time the general group relationship is evident throughout. In a recent paper, Chadwick¹ has proposed a new system of classification of the Devonian rocks in New York and Pennsylvania and offers a revision of the range of the various fossils. The field work and mapping were finished in Greenbrier County before the appearance of Chadwick's paper. As a result the older classification of Devonian rocks is followed in this report without either rejecting or accepting Chadwick's classification. The Devonian of Greenbrier County has the following succession in descending order:

Upper Devonian: (Hampshire and Jennings of U. S. Geological Survey publications).

Catskill Series (0-400').

Chemung Series (2000-3000').

Hendricks Sandstone.

Shales and sandstones.

Portage Series (2000'±).

Shales, with thin sandstones.

Genesee Series (50-100').

Shale.

Middle Devonian: (Romney of U. S. Geological Survey publications).

Hamilton and Marcellus Series (500'±).

Shales, with thin limestones.

¹Chadwick, George Haleott, Faunal Differentiation in the Upper Devonian, G. S. A. Bull., Vol. 46, No. 2, pp. 305-342; 1935.

Lower Devonian:

Oriskany Series (80-90').

Huntersville Chert.

Ridgeley Sandstone.

Helderberg Series (300'±).

Becraft.

New Scotland.

Coeymaas (?).

Keyser.

Further comment on the nomenclature of this period will follow on succeeding pages under the description of the various subdivisions.

The Devonian of Greenbrier County will average approximately 6,500 feet in thickness, and comprises almost half of the outcropping rock column. Its outcrop is limited to the eastern side of the county and almost entirely to the territory east of the Greenbrier River, the only exception being the Catskill which outcrops along this stream and occasionally west of it. Good exposures are usually available for most portions of the section although much difficulty is encountered in measuring these beds as complete units at continuous exposures, because of the frequent folding and duplication of beds. Along Mays Draft, some 4.5 miles north of White Sulphur Springs, a total of 6,000 feet of Devonian rocks was measured starting at the base of the Pocono and extending down to the base of the Marellus Series. The thickness was measured by steel tape, using a Brunton clinometer, and corrections were made for the dip of the rocks. Ten dip readings were taken along the line of traverse, the rocks dipping to the northwest at an inclination of 20 to 50 degrees from the horizontal.

UPPER DEVONIAN ROCKS.

CATSKILL SERIES.

GENERAL ACCOUNT, CATSKILL SERIES.

The Catskill Series coming at the top of the Devonian and just beneath the Pocono Series, is composed of red shales interbedded with massive green or brown sandstones with occasional green and brown shales. The sandstones are very conglomeratic in some localities and east of Anthony Creek,

two massive conglomerates, each 30 to 40 feet thick, were noted in this series. The series reaches a maximum thickness of 400 feet near the Greenbrier-Pocahontas County line and thins away to zero thickness on Greenbrier Mountain. The Catskill was not noted along Howard Creek east of Caldwell, nor does it reappear south of this point.

Throughout most of their outcrop the shales and sandstones of the Catskill appear to be lenticular, changing from one to the other within narrow limits, so that definite correlation of individual beds for any distance is quite impracticable.

TOPOGRAPHIC EXPRESSION, CATSKILL SERIES.

In Greenbrier County there are several resistant sandstones in the Catskill Series and as a result the topographic expression of this series is very much like that developed on the overlying Pocono and underlying Chemung rocks. The Catskill rocks aid in forming Little Allegheny Ridge and Meadow Creek Mountain.

AREAL EXTENT, CATSKILL SERIES.

In areal extent the Catskill Series presents a narrow outcrop along and just east of the Greenbrier River, extending from the Pocahontas line southwestward to Greenbrier Mountain. The Catskill is present in only one other area, that being a narrow outcrop on Little Allegheny Ridge and Meadow Creek Mountain in the northeastern part of the county. The outcrop of this series is delineated on Map II and the areal extent together with that of the other Upper and Middle Devonian rocks is shown on Figure 13.

FIGURE 13

MAP


OF

GREENBRIER COUNTY

SHOWING

OUTCROPS OF

UPPER AND MIDDLE DEVONIAN ROCKS



CONTACTS, CATSKILL SERIES.

The contact of the Catskill Series with the overlying Pocono of the Mississippian has already been discussed under the description of the latter series, page 286. The contact at the base of this series where it rests on the Chemung has been the subject of much discussion. The generally accepted contact has been the dividing line between the red beds and the underlying green and brown fossiliferous sandstones and shales of the Chemung. At certain localities, however, red streaks are often found interlaminated with beds of Chemung character, while olive and green shales with typical Chemung fossils have been noted well up in the red shales. It is the opinion of some authorities and particularly paleontologists, that the contact should be placed at the last recurrence of fossils regardless of the presence of red shales. If this plan were followed the areal mapping of this contact in many counties would prove to be a hopeless task. Dr. I. C. White often expressed the opinion (oral expression) that the presence of marine fossils in the basal portion of the red beds was due to the local existence of lagoons where conditions remained favorable to marine life. It is now believed by some geologists that the typical non-marine Catskill sediments of the east are contemporaneous with at least a portion of the marine sediments of the Upper Devonian to the west. This interfingering effect of these marine and non-marine sediments is accounted for by a shifting strand line.

In Greenbrier County the bottom contact of the Catskill is placed at the top of a persistent, massive, often conglomeratic sandstone that occurs near the base of the typical red shales and near the top of those beds that are characteristic of the Chemung. This sandstone, which is correlated with the Hendricks Sandstone of Reger and Price², offers what is probably the most satisfactory boundary between these two series in Greenbrier County. Because this sandstone often contains fossils of Chemung age, it is placed in that series with the contact coming immediately above.

²Reger, David B., and Price, Wm. Armstrong, Tucker Report, W. Va. Geol. Survey, pp. 245-251, 1892.

Direction of Joints in the Pickaway Limestone.

Location Observed.	Direction.	Elevation.
Falling Springs District:		
1. Along U. S. Route 219, 1 mile southwest of Falling Springs (Renlek P. O.).....	N. 45° E.	1965' B.
Frankford District:		
2. Along U. S. Route 219, 2 miles north of Frankford and 0.45 mile west of Walnut Grove Church.....	N. 46° E.	2215' B.
3. Along U. S. Route 219, 1.1 miles north of Frankford and 1.25 miles west of Gilboa School.....	N. 45° E.	2290' B.
Lewisburg District:		
4. Along public road, 0.65 mile southwest of Maxwellton and 0.7 mile east of Fairview School	N. 45° E.	2290' B.
5. Along public road, 1 mile west of Central School and 1 mile northeast of Kramer School.....	N. 40° E.	1940' B.
6. Along Lewisburg-Fort Springs District line, 1.2 miles west of George School..	N. 40°-45° E.	2065' B.
7. Along U. S. Route 60, 0.65 mile northwest of city limits of Lewisburg.....	N. 45°-50° E.	2140' B.
Fort Springs District:		
8. Along public road, 0.5 mile northwest of Livesay School and 2 miles southwest of the city limits of Lewisburg..	N. 38°-40° E.	2075' B.
9. Along public road, 1 mile northeast of Curry School and 1.8 miles southwest of Livesay School.....	N. 40°-45° E.	2125' B.
10. Along public road, 1 mile northwest of Fort Springs.....	N. 40° E.	1800' B.
Irish Corner District:		
11. Near Acme Limestone Quarry, 0.6 mile west of Fort Springs.....	N. 40° E.	1675' B.
Monroe County, Second Creek District:		
12. Along U. S. Route 219, 1.05 miles northwest of Second Creek (town) and 0.75 mile southwest of Second Creek (stream)	N. 44°-46° E.	1885' B.
13. Along road 1 mile south of town of Pickaway, (type locality of Pickaway member)	N. 37°-42° E.	2215' B.

There appears to be little or no connection between the Piekaway joints and the structural features developed during the Appalachian Revolution. As shown on Figure 11 and in more detail on Map II, the regional structural trend, in Greenbrier County, is north 25 to 30 degrees east, while the average strike of the joints is about north 40 to 45 degrees east. As mentioned above, the Piekaway joints are, so far as known, confined to a single ledge.

Regional isopach maps drawn by R. C. Tueker, on the Greenbrier Series, and on the Mauch Chunk Series show that the iso-thickness lines extend in the same direction as do the Piekaway joints. It is believed that these iso-thickness lines indicate the direction of the Mississippian shore-line and that there probably was some connection between the direction of the shore-line and the Piekaway joints.

A possible explanation of the Piekaway joints is that they represent tension fractures resulting from differential subsidence of the sedimentary basin of Greenbrier time and that their alignment was controlled by the direction of this differential subsidence. This condition may have been repeated several times but in the case of the Piekaway ledge, the newly deposited material was of just the right character to form open fractures and before these fractures were obliterated by wave action or the deposition of more lime, they were filled with argillaceous and arenaceous material.

Two more factors that may have played a part in the formation of these joints are as follows: (1) The subsidence may have been accompanied by earthquakes and after the stress was set up, the earthquakes may have started the fractures. (2) Once started the cracks may have been enlarged by drying as there are indications of shallow water conditions during deposition of this part of the Greenbrier Series.

TAGGARD LIMESTONE.

The Taggard Limestone, named by Reger²⁵ from its occurrence on Taggard Branch, Monroe County, is present in Greenbrier County and retains the same general character as noted at its type locality, except that it was not considered

²⁵Reger, David B., Mercer, Monroe, and Summers Report, W. Va. Geol. Sur., pp. 476-479: 1926.

advisable to separate it from its associated shales. In the Alta and Julia P. O. Sections, published in Chapter V, this limestone is recorded at 35 and 25 feet thick, yellowish-gray to red, shaly and somewhat oolitic. It is also shown in the Renick Special Section, page 273.

From an economic standpoint the Taggard Limestone is of minor importance, being too impure and shaly for most commercial uses.

PATTON LIMESTONE.

The **Patton Limestone**, named by Reger²⁰, from its occurrence near Patton, Monroe County, is represented in Greenbrier County by a hard, blue limestone, containing occasional nodules of black chert. It is somewhat shaly and sandy at the top and bottom but the middle portion is generally freer from impurities than most of the other members of the series. Its character, thickness, and stratigraphic position are shown in the General Section, on a preceding page of this chapter, and in the Alta, Julia P. O., and Patton Sections published in Chapter V.

The commercial possibilities of this bed are discussed and chemical analyses given in Chapter XII. Lists of fossils collected from this horizon are published in Chapter XIV.

SINKS GROVE LIMESTONE.

The **Sinks Grove Limestone**, coming just below the Patton Limestone, was first named by Reger²¹ from its exposures in the vicinity of Sinks Grove, Monroe County. This same limestone is present in Greenbrier County although its development is much less prominent than that at its type locality. It is possible that the member was often mistaken for the overlying Patton Limestone or included with it, as at the majority of their exposures there is little evidence to distinguish them from one another. In general it is a massive, blue limestone, occasionally oolitic, and it may carry scattered nodules of black chert. Its thickness, character, and stratigraphic position are shown in the Alta, Julia P. O., and Patton Sections,

²⁰Ibid., pp. 480-483.

²¹Ibid., pp. 484-487.

as published in Chapter V and its possible commercial uses are discussed in Chapter XII. Lists of fossils collected from this horizon are published in Chapter XIV.

HILLSDALE LIMESTONE.

The **Hillsdale Limestone** of Reger²⁸, named from its occurrence just east of Hillsdale, Monroe County, is represented in Greenbrier County by a grayish-blue to dark, hard, massive limestone that usually contains numerous nodules of black chert (Plate XXVI) which may weather to a gray color. It contains marine fossils that are scanty in the chert but they are abundant in the limestone matrix. In many places the Hillsdale contains many silicified fossil corals (**Lithostrotion canadense**) which are now scattered over the Macerady outcrops where the limestone has been dissolved away.

The thickness, character, and stratigraphic position of the Hillsdale Limestone are shown in the Alta, Caldwell, Horse-shoe Bend School, Julia P. O., Patton, and Spring Creek Sections as published in Chapter V. The commercial possibilities of the member are discussed in Chapter XII, and in Chapter XIV there is a rather full discussion of the fossils found in this bed.

ECONOMIC ASPECTS, GREENBRIER SERIES.

The best agricultural soil of the county is found along the outcrops of the Greenbrier Series, and as a result its entire exposures are cleared and cultivated. In this respect the limestone belts offer quite a contrast to the almost totally uncleared Pocono outcrops. In some localities, however, where the topography is too steep to retain a tillable soil, its use is limited to grazing but in regions where the surface is comparatively level, no better farming lands can be found anywhere.

The rock from this series is used as material for road macadam, railroad ballast, agricultural lime, and for chemical uses. In Chapter XII, under the subject of "Limestone," will be found a further discussion of these economic features.

²⁸Ibid., pp. 487-490.

MACCRADY SERIES.

GENERAL ACCOUNT, MACCRADY SERIES.

The Maccrady Series, comprising those beds between the Greenbrier Series and the Pocono Series, is a distinct and well-defined stratigraphic division in the area of this report. This assemblage of rocks was originally named by Campbell²⁹ the "Pulaski Shale" from its exposure in the county of that name in Virginia, but as this title had been earlier applied to an Ordovician formation in New York, Stose³⁰ gave it the name "Maccrady Formation" from its exposure in Smyth County, Virginia. Since it has been the policy of the West Virginia Survey to avoid as far as practicable the term "Formation" in the application of names to major subdivisions, Reger³¹ has substituted the term Series for that of Formation, and the same usage will be followed in this report.

The Maccrady Series at its outcrops in Greenbrier County consists of deep-red shale and weakly bedded sandstone. Its thickness is quite variable, being thickest in the southeast part of its outcrop and thinnest in the north and northwest. It is estimated as 250 feet thick in the Caldwell Section and it appears to be about 60 feet thick near the Pocahontas County line. Other thicknesses between these extremes are shown in the Alta, Cold Knob—Hinkle Well, Horseshoe Bend School, Spring Creek—South, and Spring Creek—North Sections, as published in Chapter V.

TOPOGRAPHIC EXPRESSION, MACCRADY SERIES.

As with its stratigraphic position, the topography developed on the Maccrady outcrops is intermediate between that developed on the outcrops of the Greenbrier and on those of the Pocono, being more rugged than the former and less rugged than the latter. Being largely composed of shales that yield easily to weathering the Maccrady is usually marked by low smooth slopes.

²⁹M. R. Campbell, Geol. Soc. Am., Bull., Vol. V, pp. 171, 178; 1894.

³⁰G. W. Stose, Geology of the Salt and Gypsum Deposits of Southwestern Virginia, Bull. 530, U. S. Geol. Sur., pp. 232-255; 1913.

³¹David B. Reger, Mercor, Monroe, and Summers Report, W. Va. Geol. Sur., pp. 492-493; 1926.

AREAL EXTENT, MACCRADY SERIES.

Figure 12, page 285, shows the outcrop of the Macerady and Pocono Series and on Map II the extent of the Macerady outcrop is shown in more detail and on a larger scale. The best development of this series in Greenbrier County is in the vicinity of Ronceverte and Caldwell.

CONTACTS, MACCRADY SERIES.

The upper contact of the Macerady Series with that of the Greenbrier Series has been discussed on a preceding page under the discussion of the contacts of the latter series, where it was pointed out that a disconformity of considerable magnitude exists. The length of time represented by this disconformity can not be determined until the age of the Macerady is finally settled.

The contact of the Macerady and Pocono Series appears to be conformable in Greenbrier County although the change from massive sandstone to weakly bedded red shales is usually abrupt.

FOSSIL LIFE, MACCRADY SERIES.

The Macerady Series is not fossiliferous in Greenbrier County. In adjoining areas and in Virginia it is reported that the upper part of the Macerady is fossiliferous and the lower part non-fossiliferous, but Butts²² has pointed out the desirability of separating these fossiliferous beds from the non-fossiliferous beds. In Greenbrier County, as discussed by Tilton in Chapter XIV, there are a few beds at the base of the Greenbrier Series that might be classified as Macerady if one were to ignore the fossil evidence. The fact that these beds become more numerous and attain a greater total thickness toward the south is considered as added proof of the transgressive overlap of post-Macerady beds.

CORRELATION, MACCRADY SERIES.

In view of the preceding comment, the proper correlation of the Macerady Series with its equivalent in other States remains uncertain. Since the Macerady as herein described

²²Butts, Chas., Oil and Gas Possibilities at Early Grove, Scott County, Virginia, Bull. 27, Va. Geol. Sur., pp. 3-8: 1927.

is not fossiliferous, its age can be determined only by determining the bed above and below it. As pointed out above, the series is marked by a disconformity at the top so that the age of the overlying beds serves only as the youngest limit of the age of the Maerady. At its base the Maerady appears to be conformable with the Pocono and some of the outcrops suggest that the relationship between the two series may be that of different conditions of sedimentation. In other words, beds that are Maerady in one area might be the age equivalents of beds in other areas that are Pocono.

Stose³³, in the report where he first names the Maerady, says that it probably represents the lower part of the Mauch Chunk of Pennsylvania, but this idea can not be accepted as the Mauch Chunk Series is now known to belong above the Greenbrier Series while the Maerady belongs below.

ECONOMIC ASPECTS, MACCRADY SERIES.

In Greenbrier County, the Maerady Series has been of value only as a maker of agricultural soils, for which purpose it is admirably adapted, since not only its shales but also its sandstones readily disintegrate. Along the Holston River near its type locality in Smyth County, Virginia, some of the soft beds of this series are saturated or wholly replaced by valuable deposits of gypsum and rock salt which are now being mined extensively as described by Stose³⁴. There is no evidence that such deposits are present in Greenbrier County.

It is quite possible that some of the red and purple shales could be used for the manufacture of building brick or tile, since they are usually free from calcareous or organic matter and are quite plastic at some localities. Owing to their included iron they should burn to a rich red color.

³³Stose, George W., *Geology of the Salt and Gypsum Deposits of Southwestern Virginia*, Bull. 530, U. S. Geol. Sur., p. 233; 1913.

³⁴G. W. Stose, *Geology of the Salt and Gypsum Deposits of Southwestern Virginia*, Bull. 530, U. S. Geol. Survey, pp. 232-255; 1913; also see *Gypsum Deposits of the United States*, Bull. 697, U. S. Geol. Survey, pp. 283-298; 1920.

POCONO SERIES.

GENERAL ACCOUNT AND SECTION, POCONO SERIES.

The Pocono Series, belonging just beneath the Maerady and above the Catskill, where the latter is present, is considered the basal major subdivision of the Mississippian in Greenbrier County as well as in all the counties of the State and in Maryland, Pennsylvania, and portions of other States farther west and south. The series was named by Lesley³⁵ in 1877, its previous designation having been the "Vespertine" or "No. X" of Rogers, both of which were gradually abandoned as lacking a geographic association. In 1877 also it was described as Pocono by Stevenson, Ashburner, and Platt in other publications in evident agreement with Lesley's nomenclature.

As exposed in Greenbrier County, the Pocono consists of coarse, reddish-brown, micaceous sandstone, often cross-bedded and conglomeratic, with brown, bluish-gray, and occasional red or green sandy shales, together with some impure and lenticular coals. Marine and plant fossils occur at various horizons throughout the series.

The following generalized section illustrates the occurrence of this series in Greenbrier County:

General Section of the Pocono Series for Greenbrier County.

	Thickness.	Total
	Feet.	Feet.
1. Sandstone, gray and brown, platy, alternating with gray and dark sandy shales.....	0 to 66	66
2. Coal, Merrimac, slaty, impure, lenticular, with plant fossils.....	0 to 4	70
3. Sandstone, Broad Ford, reddish-brown to gray, occasionally olive to green, ferruginous, usually thick-bedded, but often shaly, weathering to large concentric boulders; carries at least two zones of marine fossils.....	50 to 175	245
4. Shale and sandstone, gray, green, or brown, and flaggy sandstones, alternating with green, olive, blue, or red and carbonaceous shales; upper part may be Broad Ford	100 to 210	455
5. Sandstone, Berea, gray or brown, coarse to conglomeratic, usually massive but occasionally separated into olive and brown shale, and thin platy gray or brown micaceous sandstone.....	50 to 145	600

³⁵Lesley, J. P., Preface to Report HH, Sec. Geol. Survey of Pa. pp. XXIV-XXVI; 1877.

TOPOGRAPHIC EXPRESSION, POCONO SERIES.

Containing several resistant sandstones, the Pocono is now found capping many of the ridges in Greenbrier County. This series invariably produces a rough and rugged topography and such areas are generally uncleared, and are commonly referred to as "brush country." Along the eastern border of the county the basal Pocono sandstones are found capping much of Allegheny Mountain, Little Allegheny Ridge, and Meadow Creek Mountain. Just east of the Greenbrier River rocks of the same series are found capping White Rock Mountain, Kates Mountain, Greenbrier Mountain, and Peach Orchard Ridge. The Greenbrier River is entrenched in the Pocono rocks for much of its length in Greenbrier County.

AREAL EXTENT, POCONO SERIES.

On Figure 12 may be seen the general extent of Pocono and Maecrady rocks and on Map II the outcrop is outlined in much greater detail and on a larger scale.

FIGURE 12

MAP


OF

GREENBRIER COUNTY

SHOWING

OUTCROPS OF

MACCRADY AND POCONO SERIES



CONTACTS, POCONO SERIES.

The upper contact of the Pocono Series with the Macerady Series was discussed on a preceding page in connection with the latter series, where it was pointed out that the contact appears to be conformable. The bottom contact of the Pocono is easily found in the northern part of the county where the red Catskill shales are present but its conformable or unconformable nature is not easily determined. In the southern part of the county the Pocono rests upon the Chemung, the Catskill being absent, and in many places the exact location of the contact is difficult to determine. If the Catskill is actually cut out by erosion, as conditions near Caldwell suggest, the contact is, of course, disconformable but such a relationship is not proved because the exact age of the conglomerates at Caldwell could not be definitely established despite the fact that many fossil collections were made.

FOSSIL LIFE, POCONO SERIES.

The following quotation from Reger³⁰ sets the stage for a discussion of both the fossil life and correlation of the Pocono:

"For nearly 100 years the rocks composing the Pocono Series, as now called, have been studied in Pennsylvania, Maryland, and the two Virginias and have been generally regarded as fresh-water deposits, although marine fossils have been observed at isolated localities where their occurrence has been looked upon as unusual and where little attempt has been made to trace them into adjacent territory. Dr. I. C. White and C. A. Ashburner recorded three occurrences in Bedford and Huntingdon Counties, Pennsylvania, in the Second Geological Survey of that State; but farther west in Fayette, Westmoreland, and Indiana Counties, Stevenson failed to see them, even describing some of the beds which now prove to be most interesting as 'Wholly characterless,' and most of the folios of the United States Geological Survey which covered the same ground many years later record no marine fossils, although Butts speaks of a *Lingula* and a fragment of a lamellibranch. In West Virginia a few isolated occurrences were noted by Prof. S. B. Brown, Dr. W. Armstrong Price, and the writer, a short paper having once been prepared for 'Science' by Dr. Price in which some of these exposures were noted and a few instances having been noted by him in the Tucker County Report of the West Virginia Survey. In general, however, the fossils have escaped attention throughout the State. In Virginia fossils have been found at a few points in the Price (Pocono) Formation, but apparently little attempt has been made to utilize them as correlation planes,

³⁰Reger, David B., Mereer, Monroe, and Summers Report, W. Va. Geol. Sur., pp. 508-510; 1926.

the statement being made in a quite recent report that no single sandstone bed, with the exception of the basal conglomerate, can be traced from place to place.

"After studying the Pocono Series in Mercer and Monroe Counties, West Virginia, and after following the outcrop from its type locality in the Pocono Mountains of Monroe County, Pennsylvania, southwestward across Pennsylvania, Maryland, West Virginia, Virginia, Kentucky, and Tennessee, the writer has found, much to his own confusion as well as to that of his predecessors, that abundant marine fossils exist in various zones of the series all the way from the Broad Top Coal Field of Huntington and Bedford Counties, Pennsylvania, westward into Blair, Westmoreland, Fayette, and other counties that border the eastern rim of the Appalachian basin and southward and southwestward across West Virginia by way of Preston, Tucker, Pocahontas, Greenbrier, Monroe, and Mercer Counties, to the Virginia State line. In the latter State they may also be followed from the coal fields of Montgomery County southwestward into Tennessee through the medium of the Price and Grainger beds and westward into Kentucky where part of the series is known as New Providence and where its fossils have had careful study.

"Such parts of the above study as properly pertain to Mercer and Monroe Counties, West Virginia, will be detailed under the 'Description of Members' on later pages of this Report but the more extensive studies will be reserved for a subsequent volume on the Mississippian.

"Before passing from the subject, however, it is well to note that some of the species found in the lower portions of the Pocono are types which have been regarded as confined almost exclusively to the Chemung Series of the Devonian, so that Dr. Glrty has not accepted them as belonging to the Mississippian Period. His viewpoint on some of these collections is quite natural since in many instances he did not see the localities in the field and had no evidence except that of the fossils themselves. In central and northern West Virginia, as well as throughout Pennsylvania, the distinctly red shales of the Catskill Series, varying from a few hundred to several thousand feet in thickness and always being easily recognized, intervene between the Chemung and Pocono, affording a lithologic sequence that can not be disregarded, so that the Pocono Series with its well-known Mississippian flora and its occasional beds of coal can always be identified. Under such conditions the presence of a fauna with certain Chemung aspects in the Pocono must be considered only as a recurrence of these species in younger strata. Such a recurrence need not be surprising, however, since the fauna of the Pocono, as already explained, has had only fragmentary study, and it would appear necessary to abandon the idea that certain types, including *Spirifer disjunctus*, perished before the close of the Devonian."

As indicated in the above quotation, the fossil life of the Pocono has not received the study it deserves in West Virginia and in the surrounding States. Chadwick³² has recently pub-

³²Chadwick, George Haleott, The Great Catskill Delta, The Pan-Amer. Geol., Vol. LX, No. 2; 1933: What is Pocono?, Amer. Jour. Sci., 5 ser., Vol. 29, No. 170, pp. 133-143; 1935: Faunal Differentiation in the Lower Devonian, G. S. A. Bull. Vol. 46, No. 2, pp. 205-242; 1935.

lished several papers that are in part or entirely on the Pocono of northern Pennsylvania. The sum total of his work, however, is that in that area the age of the "Pocono" not only varies but it is of Devonian age. Based on fossil plant evidence, David White³⁸ considers the Pocono to be Mississippian all the way from "East Mauch Chunk, on the slope of the Pocono Mountains, (in Pennsylvania) southward along the east side of the Appalachia Trough as far as Tennessee" Although Chadwick³⁹ says that he accepts White's thesis without question, save the use of the name Pocono, he implies that White should check the geologic range of his fossils. In the same paper Chadwick⁴⁰ also points out that I. C. White⁴¹ reports that there is no Pocono in Pocono Mountain or in Pocono Township or in fact in the whole Pocono plateau, except topping a few peaks and that the thesis and map of Norman Spenser Wagner⁴² fully confirms I. C. White's discovery that the "Pocono" does not exist on the Pocono plateau. In the same paper Chadwick also states that in Fayette County, Pennsylvania, the Pocono beds are Canadaway. David White⁴³ states that Reger⁴⁴ and Girty⁴⁵ have proved the Mississippian age of the Pocono in the Broadtop basin, Pa., but Chadwick⁴⁶ says that his "reading of Doctor Girty's interpretations has not been so unqualified." Along the same line it is interesting to note that Reger⁴⁷ reports:

"In this connection, however, it is well to remark that in northern West Virginia and on the Youghiogheny and Conemaugh Rivers of Pennsylvania where the Broad Ford Sandstone becomes quite shaly, the faunas of this and other members of the lower part of the Pocono

³⁸White, David, The Age of the Pocono, Amer. Jour. Sci., 5 ser., Vol. 27, No. 160, pp. 265-272; 1934; see also a discussion of Mississippian plants by White in the Mereer, Monroe, and Summers Report, W. Va. Geol. Sur., pp. 837-843; 1926.

³⁹Chadwick, George Halcott, What is Pocono?, *ibid.*, see especially the foot-note, p. 133.

⁴⁰*Ibid.*, see p. 142.

⁴¹White, I. C., 2nd Geol. Sur. Pa., G6, pp. 89-90; 1882.

⁴²See Chadwick's foot-note, *ibid.*, p. 143.

⁴³Reference, foot-note 38, p. 270.

⁴⁴Reger, David B., Pocono Stratigraphy in the Broadtop basin of Pennsylvania; Bull. G. S. A., Vol., 39, pp. 397-410; 1927.

⁴⁵Girty, G. H., Pocono fauna of the Broadtop coal field, Pa., U. S. Geol. Sur., Prof. Paper 150E, p. 127; 1928.

⁴⁶*Ibid.*, p. 141.

⁴⁷Reger, David B., Mereer, Monroe, and Summers Report, W. Va. Geol. Sur., p. 525; 1926.

PLATE XXVI.—Chert nodules weathered in relief in Hillsdale Limestone on Mill Creek, 1.6 miles south of Ashbury.





PLATE XXVII.—Quarrying road material from the Broad Ford Sandstone in a road cut east of Caldwell.




PLATE XX.—Stylolites in basal Greenbrier Limestone along Midland Trail (U. S. Route 60) west of Lewisburg.



PLATE XXI.—Mud cracks in Taggard Limestone along Midland Trail (U. S. Route 60) one mile west of Alta.





PLATE XXII.—Greenbrier Limestone stripped of cover for quarrying at the Acme Limestone Company Quarry near Port Spring. Note typical Pickaway joints.






PLATE XXIII.—Typical joints in Pickaway Limestone near Union, Monroe County. Cross-section view of bed.



PLATE XXIV.—Typical joints in Pickaway Limestone near Union, Monroe County. Top of bed shown.



PLATE XXV.—Typical joints in Pickaway Limestone near Union, Monroe County.

chert nodules were noted in the lower part of this member. Its thickness, character, and stratigraphic position are shown in the Acme Limestone Quarry, Alta, Julia Post-Office, and Renick Sections as published in Chapter V and lists of fossils collected from this horizon are published in Chapter XIV. Its economic possibilities will be considered in Chapter XII.

Within the Pickaway member there is one ledge, near the top, that is characteristically jointed, normal to the bedding-plane. The thickness of the ledge affected by the joints is usually between 3 and 8 feet but at the Acme Limestone Quarry the jointed ledge was observed to be 15 feet thick (Plate XXII). No similar joints were found in beds above or below this particular ledge. The following special section shows the succession of beds above and below the jointed bed:

Renick Special Section.

Falling Springs District; begins 1 mile southwest of Renick P. O. and measured along the road to Spring Creek; arrangement in descending stratigraphic order.

Thickness. Total.		Feet. Foot.		Greenbrier Series (125+)	
10	10+	10	25	Limestone, Union, oolite, fossiliferous.....	10
				Limestone, yellowish-gray, weathers banded ribbon effect	10
15	15	15	55	Limestone, dark-gray, granular, ealite streaks, part oolite, large fossils, lower part maroon and siliceous	30
				Limestone, greenish-yellow, weathers	
				white, argillaceous.....	10
				Limestone, dark-blue, weathers yellow, characteristic joints.....	5
				Limestone, yellowish-blue to dark-blue, impure.....	35
				Limestone, red and greenish-yellow.....	3
				Limestone, greenish-blue.....	2
				Limestone, gray-blue, granular, siliceous, very hard.....	15
125	20	20	125	Concealed to creek.....	15
				Taggard	20
				Pickaway	50
				105
				55

In many places the impure limestone, noted in the above section immediately over the jointed ledge, grades into a calcareous shale.

*The partings in this ledge lack many of the characteristics that are usually inferred by the use of the term joint. For the want of a better term, however, these partings are called joints in this report.

chert nodules were noted in the lower part of this member. Its thickness, character, and stratigraphic position are shown in the Aeme Limestone Quarry, Alta, Julia Post-Office, and Renick Sections as published in Chapter V and lists of fossils collected from this horizon are published in Chapter XIV. Its economic possibilities will be considered in Chapter XII.

Within the Pickaway member there is one ledge, near the top, that is characteristically jointed²⁴ normal to the bedding-plane. The thickness of the ledge affected by the joints is usually between 3 and 8 feet but at the Aeme Limestone Quarry the jointed ledge was observed to be 15 feet thick (Plate XXII). No similar joints were found in beds above or below this particular ledge. The following special section shows the succession of beds above and below the jointed bed:

Renick Special Section.

Falling Springs District; begins 1 mile southwest of Renick P. O. and measured along the road to Spring Creek; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Greenbrier Series (125'+)		
Limestone, Union, oolitic, fossiliferous.....	10+	10
Limestone, yellowish-gray, weathers banded ribbon effect	15	25
Limestone, dark-gray, granular, calcite streaks, part oolitic, large fossils, lower part maroon and siliceous	30	55
Limestone, greenish-yellow, weathers white, argillaceous.....10'	} Pickaway ... 50	105
Limestone, dark-blue, weathers yellow, characteristic joints..... 5		
Limestone, yellowish-blue to dark-blue, impure.....35		
Limestone, red and greenish-yellow.. 3'		
Limestone, greenish-blue..... 2	} Taggard 20	125
Limestone, gray-blue, granular, siliceous, very hard.....15		
Concealed to creek.....		

In many places the impure limestone, noted in the above section immediately over the jointed ledge, grades into a calcareous shale.

²⁴The partings in this ledge lack many of the characteristics that are usually inferred by the use of the term joint. For the want of a better term, however, these partings are called joints in this report.

The joints are filled with an impure calcareous cement that disintegrates more easily than does the limestone proper and as a result, weathering gives the ledge a conspicuous and distinctive appearance (see Plates XXIII, XXIV, and XXV). The unweathered rock will break across the joints almost as easily as along them so that it was possible to chip away most of the rock on either side of one of the joints, leaving a piece of rock that was about 60 per cent. joint filling. Chemical tests show that the sample has the following composition:

	Per cent.
Silica (SiO_2).....	37.85
Ferric Iron (Fe_2O_3).....	5.98
Alumina (Al_2O_3).....	14.70
Lime (CaO).....	16.38
Calcium Carbonate (CaCO_3).....	29.23*
Magnesia (MgO).....	2.82
Magnesium Carbonate (MgCO_3).....	5.89*
Potash (K_2O).....	3.67
Soda (Na_2O).....	0.86
Titanium Oxide (TiO_2).....	0.81
Phosphoric Acid (P_2O_5).....	1.01

*Calculated from the oxides.

100.00

By comparing the above analysis with analyses of samples taken from the entire ledge (as published in Chapter XII), it is seen that the material filling the joints is largely clay and quartz minerals that were probably added after the bed was deposited. It is therefore believed that the joints or tension fractures were in existence prior to the deposition of the next younger bed and that these open joints were filled with mud and sand during the deposition of the younger bed. Such conditions would suggest drying or mud-cracking as the cause of the tension joints but the joints do not have the polygonal pattern characteristic of mud-cracks (compare Plate XXI with Plate XXIV).

Individual joints are rarely over 10 feet long and are neither perfectly straight nor exactly parallel. One of the remarkable things about these joints, however, is the constancy of their average direction over a distance of some 30 miles along their outcrop. Figure 11 shows a number of locations at which the direction of the Pickaway joints was measured and the following table gives the data for these localities:




Figure 11.—Map showing localities where the direction of Piek-away joints was measured, and near-by major structural features.

to state definitely that such an unconformity exists. The Hinton Group is composed of approximately 70 per cent. shale, 25 per cent. sandstone, and 5 per cent. (or less) limestone.

The upper half of the Hinton Group is composed of shales and thin sandstones. The shales are predominantly red or variegated and some beds may be calcareous, while the sandstones are usually greenish-gray, thin-bedded and shaly. In the general vicinity of Kieffer, two or more thin, very impure coals were noted in this part of the Hinton Group.

AVIS LIMESTONE.

The **Avis Limestone** of Reger^a, formerly termed the **Hinton Limestone** by Krebs^b, is one of the most persistent and easily recognized members within the Hinton Group in Greenbrier County. It is usually steel gray in color although the top may be stained yellow on weathering. The limestone is sometimes divided into two benches, being separated by a thin bed of calcareous shale. Its character, thickness, and stratigraphic position are shown in the Kieffer, Roach Run, Alderson, and Cold Knob—Hinkle Well Sections, published in Chapter V.

This limestone has been quarried along the Midland Trail just east of Little Clear Creek and its possibilities as a quarry rock together with chemical analyses are discussed in Chapter XII.

Between the Avis Limestone and the Stony Gap Sandstone there are from 300 to 500 feet of red or variegated shales, interbedded with greenish-gray or red sandstones. Some of the beds are strongly calcareous and locally they may grade into limestones.

STONY GAP SANDSTONE.

The **Stony Gap Sandstone** of Reger^c, or **Hinton** of Stevenson, is present in Greenbrier County and forms the basal member of the Hinton Group. This sandstone was recognized

^aIbid., pp. 347-351.

^bC. E. Krebs, Raleigh County and Western Portions of Mercer and Summers Counties Report, W. Va. Geol. Sur., pp. 75, 76, and 88; 1916.

^cDavid B. Reger, Mercer, Monroe, and Summers Report, W. Va. Geol. Sur., pp. 371-378; 1926.

many years ago as an important key rock and was called the Hinton Sandstone by Dr. John J. Stevenson from its exposure near Hinton, Summers County, but apparently little recognition was given it. Later Campbell^o, applied the term "Hinton Formation" to a major group of rocks in the New River Valley, and its usage has become so well fixed in the geologic literature of West Virginia that it appears unwise to return to the earlier application, which would possibly lead to confusion. Reger has accordingly renamed this horizon the Stony Gap Sandstone from its occurrence at the village of that name in Mercer County, where it is well exposed. At its type locality it is described as being a light-gray or white, massive, coarse, and extremely hard and quartzitic ledge, varying in thickness from 35 to 85 feet.

In Greenbrier County this sandstone retains its same general character, being a gray to white, medium-grained, massive, hard, and quartzitic sandstone, but attaining no thickness greater than 50 feet. Its position can be noted in the measured sections containing the Mauch Chunk Series and located in detail from Map II as it forms the basal member of the Hinton Group which is thereon delineated.

So far as known no use has been made of this stratum for any purpose, but owing to its resistant character, its purity, and its pleasing appearance, it should be suitable for building stone and other local uses.

DESCRIPTION OF MEMBERS, BLUEFIELD GROUP.

The Bluefield Group is the largest subdivision of the Mauch Chunk Series and in Greenbrier County it is composed of 60 to 65 per cent. shale, 30 per cent. sandstone, and 5 to 10 per cent. limestone. In appearance the upper two-thirds of this group is quite similar to the upper groups of the Mauch Chunk but the bottom third is intermediate in appearance between the rest of the Mauch Chunk and the underlying Greenbrier Series. On ordinary hillside exposures it is sometimes difficult to tell where the Mauch Chunk-Greenbrier contact belongs.

^oM. R. Campbell, Pocahontas Folio, No. 26. U. S. Geol. Sur.; 1896.

The upper 550 to 600 feet of the Bluefield Group is composed of shales and sandstones. The shales are mostly red but some green or brown beds were noted. Many of the beds are calcareous and some may locally grade into limestone. The sandstones are greenish-gray or reddish-brown, usually thin-bedded, shaly, and fine-grained.

DROOP SANDSTONE.

The **Droop Sandstone** was named by Reger¹⁰ from its occurrence on Droop Mountain, Pocahontas County. In Greenbrier County this sandstone is usually grayish-brown, massive, medium-grained, and hard. It is frequently cross-bedded and ripple-marked and sometimes carries carbonized plants. Its thickness, character, and stratigraphic position are noted in the Alum Run, Alta, Alderson, Butler Mountain, Hawver School—East, Cold Knob—Hinkle Well, and Richlands—Two Miles North Sections, published in Chapter V. In thickness this sandstone exceeds all other sandstones in the Mauch Chunk, frequently attaining a thickness in excess of 60 feet.

Due to the fact that it is much more resistant than the beds immediately above and below it, the Droop Sandstone is often found capping the ridges. Under such conditions, weathering often removes much of the iron in the sandstone, leaving a nearly pure silica sand that appears to have the properties of a glass-sand. In some localities the Droop Sandstone is strongly cemented with secondary silica and appears to be durable enough for road material. So far as known this sandstone has not been quarried in Greenbrier County for either purpose.

TALCOTT AND ADA SHALES.

A shale bed that is believed to represent both the **Talcott and Ada Shales** of Reger¹¹ was noted in the Renick Valley Section. A yellow to olive-green sandy shale 55 feet thick was noted immediately under the sandstone last described. Elsewhere these beds were not identified.

¹⁰Reger, David B., Mereer, Monroe, and Summers Report, W. Va. Geol. Sur., pp. 415-418; 1926.

¹¹Ibid., pp. 418-426.

REYNOLDS LIMESTONE.

The **Reynolds Limestone**, of Reger¹², named from its occurrence in Monroe County near Reynolds School, is a shaly, blue to yellowish-blue limestone, 15 to 40 feet thick in Greenbrier County. It is very fossiliferous and lists of fossils identified in collections from this horizon are published in Chapter XIV. Its thickness, character, and stratigraphic position are shown in the Alta, Blue Sulphur Springs, Alderson, Butler Mountain, Briery Knob, Renick, Hawver School—East and Hawver School—West Sections, published in Chapter V. Due to the usual proximity of the outcrop of the Reynolds Limestone to that of the very pure limestones of the Greenbrier Series, it is of little economic value, although locally it may furnish a small amount of road material or agricultural lime.

Between the limestone just described and the underlying Webster Springs Sandstone, occurs a shale as shown in the General Section, which may be the equivalent of the Bickett Shale of Reger¹³.

WEBSTER SPRINGS SANDSTONE.

The **Webster Springs Sandstone** of Reger¹⁴ is represented in Greenbrier County by 10 to 50 feet of shaly, grayish-brown sandstone. Its character, thickness, and stratigraphic position are shown in the Cold Knob—Hinkle Well, Renick, and Renick Valley Sections, as published in Chapter V.

GLENRAY LIMESTONE.

The **Glenray Limestone** of Reger¹⁵ is represented in Greenbrier County by 10 to 60 feet of more or less impure limestone. It is usually a bluish-gray, siliceous, thick-bedded, very fossiliferous limestone, belonging 100 to 150 feet above the base of the Mauch Chunk Series. Its stratigraphic position is shown in the Alta, Alderson, Blaker Mills, Blue Sulphur Springs,

¹²Ibid., pp. 426-430.

¹³Ibid., pp. 430-431.

¹⁴Reger, David B., Webster County Report, W. Va. Geol. Sur., pp. 227-228; 1920.

¹⁵Reger, David B., Mercer, Monroe, and Summers Report, W. Va. Geol. Sur., pp. 432-437, 1926.

Hawver School—East, Hawver School—West, Richlands—Two Miles North, and Reniek Sections, as published in Chapter V.

A discussion of the commercial possibilities of the Glenray Limestone is published in Chapter XII and lists of fossils collected from this horizon are published in Chapter XIV.

LILLYDALE SHALE.

The **Lillydale Shale** of Reger¹⁶ is represented in Greenbrier County by a dark to greenish-gray, concretionary, micaceous shale that is usually somewhat carbonaceous. This somewhat fossiliferous bed is believed to be the same as the "**Pencil Cave**" of the oil well drillers of central and northern West Virginia. Its thickness, character, and stratigraphic position are noted in the Alta, Alderson, Butler Mountain, Briery Knob, Reniek, Reniek Valley, Richlands—Two Miles North, Savannah School, and Unus Sections as published in Chapter V. Lists of fossils collected from this shale are published in Chapter XIV.

A shaly lenticular sandstone that may correspond to the **Edray Sandstone** of Reger¹⁷, was noted in the Briery Knob, Reniek, Richlands—Northwest, and Richlands—Two Miles North Sections, as published in Chapter V. In general it is a thin, poorly defined stratum in Greenbrier County.

ECONOMIC ASPECTS, MAUCH CHUNK SERIES.

From an economic standpoint the Mauch Chunk Series does not have much to offer which can be readily exploited. The coals are all too thin and impure for even local domestic use. So far as known it contains no precious ore or metals. The shales could be used for the manufacture of brick and tile, but owing to an almost universal occurrence of this material, the demand would be limited to local use. The limestone of this series is of little value except as a soil maker as compared to the underlying Greenbrier Series. The soil from this series seems best adapted for timber growth and grazing land. One sandstone, the Droop, offers a good prospect as a glass-sand.

¹⁶Ibid., pp. 437-443.

¹⁷Ibid., pp. 443-445.

GREENBRIER SERIES.

GENERAL ACCOUNT AND SECTION, GREENBRIER SERIES.

The Greenbrier Series, comprising the middle portion of the Mississippian and coming directly under the Manah Chunk Series and immediately over the Maccrady Series, is composed almost entirely of limestone rocks. The name was derived, apparently, from the Greenbrier River, along which its best and greatest exposures occur, but by whom the title was first applied is not known. It is possible the name "Maxville" of Andrews¹⁸ is entitled to priority, but like many other instances, the term Greenbrier has become so fixed in the literature of this and adjoining States that it seems unwise to supplant it by the Ohio title. Furthermore this formation in the latter State represents only a small portion of the series at its type locality in West Virginia, and no definite correlation between the two has been made.

The base of this series in West Virginia has been quite definitely established as resting upon the Maccrady red and purple shales in the southern counties; and on the Pocono sandstones, which offer a still greater contrast, in northern West Virginia where the former shaly beds have disappeared.

The Greenbrier Series in the area under discussion has a thickness that varies from approximately 475 to 750 feet, with a rapid thinning to the northeastward. Its maximum thickness here offers a contrast to its much greater thickness in adjoining counties to the south where Reger¹⁹ has been able to trace many of the minor subdivisions over considerable areas and has given them suitable titles. These subdivisions while somewhat attenuated have been recognized in Greenbrier County and will be retained, so far as applicable, in this report. The subdivisions have been based mainly on lithologic characteristics.

The following general section was prepared from several measured sections and local notes and indicates the character of the series in the area of this report:

¹⁸E. B. Andrews, Ohio Geol. Sur., Report Progress, 1869, pp. 80, 84; 1870.

¹⁹David B. Reger, Mereer, Monroe, and Summers Report, W. Va. Geol. Sur., pp. 449-451; 1926.

General Section of the Greenbrier Series for Greenbrier County, West Virginia.

	Thiekness. Feet.	Total. Feet.
Limestone, Alderson, dark-gray, sandy, with crystalline streaks; very hard, oeeasionally oolitic, with numerous fosslls, bryozoa (<i>Archimedes</i>), brachlopods, erlnolds, (especlally <i>Pterotocrinus</i>), eorals, and a few pelecypods	50 to 150	150
Shale, Greenville, brown to dark, fissile, cal-careous, lenticular, with marine fosslls; abundant <i>Chonetes</i> , fish tooth.....	40 to 0	150
Limestone, Union, gray to dark, weathering white, hard, shaly at top; oolitic in part; contains profuse marine fosslls; <i>Pentremites</i> , <i>Archimedes</i> , gastropods, bryozoa.....	200 to 150	300
Limestone, Pickaway, dark, hard, brittle, with oeeasional red streaks, but with only sparing marine fosslls.....	50 to 135	435
Limestone, Taggard, gray, oolitic, fosslliferous, associated with red shale.....	10 to 35	470
Limestone, Patton, somewhat shaly at top, but hard, pure, and weathering gray at base; usually contains 5 to 10 feet of light-gray oolite; marine fosslls; oeeasional nodules of black chert.....	150 to 90	560
Limestone, Sinks Grove, blue, hard, siliceous, weathering yellow at top and gray at base; often contains nodules of black chert; also contains marine fosslls, brachlopods, bryozoa, crinolds, and gastropods.....	40 to 90	650
Limestone, Hillsdale (St. Louis age as corre-lated in Kentucky), grayish-blue, hard, mas-sive; profuse marine fosslls including <i>Litho-strotion canadense</i> (<i>basaltiforme</i>), <i>L. prolf-erum</i> ; contains nodules of gray and black chert	30 to 100	750
Maccrady Series.....

TOPOGRAPHIC EXPRESSION, GREENBRIER SERIES.

In Greenbrier County there is a large area in which the outcropping rocks are limestones of the Greenbrier Series. In much of this area, a typical "karst" topography has been developed that is characterized by the presence of numerous sink-holes, a relatively low relief, and the general absence of an interconnecting valley system. This relatively low relief developed on the rocks of the Greenbrier Series is believed to be due in part to the absence of valley cutting and in part to the development of an intermediate erosion surface. Most of the streams crossing the Greenbrier outcrop have estab-


lished subterranean courses, which accounts for the apparent absence of valley cutting in the area.

Where outliers of basal Mauch Chunk remain, the underlying limestones have been protected from chemical erosion and as a result these outliers are usually found capping a ridge or knob. The effect of Mauch Chunk outliers is well illustrated by Falling Spring Mountain and Weaver Knob.

AREAL EXTENT, GREENBRIER SERIES.

The areal extent of the Greenbrier Series in Greenbrier County can be seen at a glance on Figure 10, while Map II shows the outcrops in much greater detail. The entire thickness of the limestone rocks is exposed along U. S. Routes 219 and 60, where they may be studied in detail.

FIGURE 10
MAP
OF
GREENBRIER COUNTY
SHOWING
OUTCROPS OF
GREENBRIER SERIES



CONTACTS, GREENBRIER SERIES.

The contact of the Greenbrier and Mauch Chunk Series is conformable as discussed on a foregoing page under the description of the latter series.

At the base of the Greenbrier the contact with the Maerady is much more marked and it is clear that an unconformity exists. Below the lowest massive limestone bed, there often occurs a calcareous shale that appears to be re-worked Maerady material. This shale may laterally grade into impure limestone that carries St. Louis fossils as described by the late Professor Tilton in Chapter XIV. Apparently there are no beds of Spergen (Salem) or Warsaw age in Greenbrier County.

FOSSIL LIFE, GREENBRIER SERIES.

In Greenbrier County this series is more or less fossiliferous throughout. A large number of collections were made and these have been identified by Professors Dana Wells and John L. Tilton. Lists of the fossils identified from each collection are published in Chapter XIV.

CORRELATION, GREENBRIER LIMESTONE.

The Greenbrier Series was mapped in this county using the same unit boundaries that were used in the Survey Report on Mercer, Monroe, and Summers Counties to the south and in the Pocahontas County Report to the north. As mapped the series starts at the top with the Alderson Limestone and extends down to, and includes, the lowest massive limestone bed of the Hillsdale member.

During the course of the field work a number of fossil collections were made with special reference to the Greenbrier-Maerady contact. Subsequent to the completion of the field work on this part of Greenbrier County, detailed study of the fossils in these collections by the late Professor John L. Tilton indicates that 5 to 40 feet of calcareous shale that had been mapped as Maerady belongs in the Greenbrier Series. The paleontologic evidence involved is presented, in more detail, in Chapter XIV. It is hardly necessary to point out that the inclusion of such a small thickness of beds in the Greenbrier Series does not materially affect the areal extent of the series as shown on Map II.

DESCRIPTION OF MEMBERS, GREENBRIER SERIES.

ALDERSON LIMESTONE.

The **Alderson Limestone** was named by Reger²⁰ from its occurrence in the vicinity of Alderson, Monroe County, where it is described as a dark-gray calcareous formation, weathering to an earthy yellow color, with a thickness which varies from 75 to 325 feet, and having an abundance of marine fossils. Attention is called to the variation in bedding at its type locality, there being some zones which are highly siliceous and which make a hard and durable limestone, and others which are fairly pure and crystalline, while still others are shaly and readily disintegrate. In Greenbrier County, somewhat the same character is retained except in a lesser degree. This member represents the succession of beds coming between the dark Lillydale Shale of the Mauch Chunk Series and the underlying Greenville Shale. In the general section at the beginning of this chapter it is shown as being dark-gray and sandy, with crystalline streaks, very hard, and containing numerous marine fossils, the most conspicuous of which are **Pentremites** which weather out in great abundance and which are locally called "petrified hickory nuts."

The thickness, character, and stratigraphic position of the Alderson Limestone are shown in the Alta, Alum Run, Aeme Limestone Quarry, Alderson, Blaker Mills, Butler Mountain, Briery Knob, Hawver School—East, Renick, Renick Valley, Richlands—Northwest, Savannah School, and Unns Sections, published in Chapter V. Lists of fossils collected from this member are published in Chapter XIV and the use of this member as a quarry rock is discussed in Chapter XII.

GREENVILLE SHALE.

The **Greenville Shale**, named by Reger²¹ from its occurrence near Greenville, Monroe County, where it is a black, fissile, and carbonaceous deposit, belonging, when present, between the Alderson and Union Limestones, and being quite lenticular and containing marine fossils, is present in Green-

²⁰David B. Reger, Mereer, Monroe, and Summers Report, W. Va. Geol. Sur., pp. 462-466; 1926.

²¹Ibid., pp. 466-7.

brier County. This shale is brown to dark green, fissile, and calcareous, containing numerous marine fossils, and an occasional shark tooth. Its thickness, character, and stratigraphic position are shown in the Aeme Limestone Quarry, Alta, Briery Knob, and Richlands—Northwest Sections. Fossils collected from this horizon are listed in Chapter XIV.

UNION LIMESTONE.

The **Union Limestone**, belonging just under the Greenville Shale, is probably the most important and persistent member of the Greenhrier Series in Greenbrier County. It was named by Reger²² from its occurrence at Union, Monroe County, where it is a gray, hard limestone weathering white, and being often crystalline, usually pure, frequently having an oolitic structure and containing numerous marine fossils, its thickness varying from 100 to 275 feet. In Greenbrier County the same general character is retained, its nature being that of a gray to dark, hard limestone, which weathers white, is shaly at the top, and usually oolitic. Marine fossils are scattered throughout but so retained in the matrix that collections are not readily made.

The thickness, character, and stratigraphic position of the Union Limestone are shown in the Aeme Limestone Quarry, Alta, Butler Mountain, Renick, Renick Valley, Richlands—Northwest, and Julia Post-Office Sections, as published in Chapter V and lists of fossils collected from this member are published in Chapter XIV. This member is a source of lime for chemical use, agricultural lime, and road material and its use for these purposes will be discussed in Chapter XII.

PICKAWAY LIMESTONE.

The **Pickaway Limestone**, named by Reger²³ from its occurrence in Monroe County, near Pickaway, and described as a very dark, hard, and sandy deposit immediately below the Union Limestone, varying in thickness from 175 to 400 feet, is present in Greenbrier County. It is usually blue to yellow in color, shaly at the top and massive at the base. Occasionally

²²Ibid., pp. 467-472.

²³Ibid., pp. 473-476.



PLATE XVIII.—Cross-bedding in the Webster Springs Sandstone (Mauch Chunk) one mile southwest of Modoc P. O.,
falling Spring Mountain.



PLATE XIX.—Greenbrier Limestone topography near Lewisburg. Mauch Chunk hills in background.

lations be largely based on the lithologic characteristics of the different beds and quite naturally the boundaries of these lithologic units may not correspond to the boundaries of paleontologic units in other areas. In some areas where the Mississippian has been divided into paleontologic units, its total thickness is measured in hundreds of feet and its units in tens of feet, whereas, in West Virginia the Mississippian may be measured in thousands of feet and its units may be hundreds of feet thick.

Mr. R. C. Tucker has recently compiled a chart showing the range of the fossils thus far reported from the Mississippian rocks of West Virginia, and this chart shows that very few of the fossils are confined to a single bed or even to a small group of beds. As a result the writers believe that it would be unwise to attempt to name interstate age equivalents in any but the most general terms.

It is clear that all of the Mississippian beds above the St. Louis Limestone as described by Weller¹ in Illinois, are represented in Greenbrier County and that the equivalent of the St. Louis Limestone is the basal member of the Greenbrier Series, as herein described. The upper limit of St. Louis beds or the limits of the equivalent of the St. Genevieve Limestone or even the limits of the Chester Group can not be stated with any degree of accuracy. It may be said, however, that there is no evidence to support the somewhat prevalent idea that the base of the Mauch Chunk Series corresponds roughly to the base of the Chester Group of Illinois. There is some evidence that both the lower beds of the Mauch Chunk Series and the upper part of the Greenbrier Series are of Chester age.

The Maecrady Series in this area is apparently non-fossiliferous, and its exact age in Greenbrier County is not known. Lithologically it resembles both the Mauch Chunk of upper Mississippian age and the Catskill of upper Devonian age.

The Pocono Series has always been considered Mississippian by the West Virginia Geological Survey, but it must be admitted that there is little to prove the age of the series in

¹Weller, Stuart, The Mississippian Brachiopoda of the Mississippi Valley Basin, Ill. St. Geol. Sur., 1914.

Greenbrier County. Professor Wells states that the fossils collected from the Pocono of Greenbrier County suggest its Mississippian age, but unfortunately the specimens are not complete enough to permit an unqualified statement.

MAUCH CHUNK SERIES.

GENERAL ACCOUNT AND SECTION, MAUCH CHUNK SERIES.

The Mauch Chunk Series, the upper division of the Mississippian, underlies the Pottsville Series of the Pennsylvanian. Its greatest thickness is along the Greenbrier-Summers County line where the series is approximately 2,800 feet thick. The least thickness of the series at the outcrop is at the Greenbrier-Pocahontas County line where it is approximately 1,900 feet thick. It is probable that a well drilled near the county line on North Fork of Cherry River would not find more than 1,400 feet of Mauch Chunk rocks and one drilled at Russellville would probably not find over 1,000 feet in this series. From the foregoing figures it is seen that the Mauch Chunk Series thins to the northwest at a very rapid rate. This thinning is a combination of a loss of thickness of individual beds and a loss of some of the Bluestone beds at the Pottsville-Mauch Chunk disconformity.

The rocks of the Mauch Chunk Series are composed of shales, sandstones, limestones, and a few impure coals. The proportion of one type of rock to another varies rapidly from place to place. Rocks of nearly every color common to sedimentary rocks may be found in this series but deep red or greenish-gray rocks are predominant.

The following general section illustrates the nature of the Mauch Chunk stratigraphic column in Greenbrier County:

General Section of the Mauch Chunk Series for Greenbrier County.

	Thickness. Feet.	Total. Feet.
Bluestone Group (80' to 675')		
Shales, red, with some green beds, occasional micaceous sandstone; may contain one or more thin, shaly limestone beds; contains two thin lenticular coaly shales.....	80-675	675
Princeton Group (20' to 80')		
Sandstone, greenish-gray, or stained reddish-brown by limonite; often a mass of pebbles and these are characteristically poorly sorted; occasional plant fossil.....	20- 80	755
Hinton Group (500' to 850')		
Shales, red, variegated, interbedded with green to red argillaceous sandstone, some beds highly calcareous; contains two or more thin coaly shales, near Kieffer.....	170-270	1025
Sandstone, gray to brown, often shaly, calcareous	0- 20	1045
Shale, calcareous, often quite sandy.....	0- 20	1065
Limestone, Avis, steel gray, may be stained yellow by weathering, shaly, very fossiliferous.....	10- 30	1095
Shales, red, variegated, interbedded with greenish-gray to red sandstones, some beds highly calcareous	290-460	1555
Sandstone, Stony Gap, greenish-gray, white or reddish-brown, massive, often cross-bedded, medium-grained, resistant to weathering.....	30- 50	1605
Bluefield Group (900' to 1200')		
Shales, mostly red, some green, some brown, interbedded with greenish-gray or reddish-brown sandstones; contains two or more thin shaly limestones.....	550-600	2205
Sandstone, Droop, gray, white or brown, medium-grained, massive, often strongly cross-bedded, sometimes carries carbonized plants.....	50-100	2305
Shale, yellow, olive, sandy.....	90-140	2445
Limestone, Reynolds, blue on fresh exposure, weathers yellow, usually impure, shaly, very fossiliferous	15- 40	2485
Shale, yellow, sandy, with streaks of red shale.....	70- 40	2525
Sandstone, Webster Springs, grayish-brown, medium-grained, shaly.....	10- 50	2575
Limestone, Glenray, gray, hard, siliceous or shaly, very fossiliferous.....	10- 60	2635
Shale, red to yellow, sandy.....	30- 40	2675
Shale, Lillydale, greenish-gray to yellow at top, dark at base, fissile, somewhat sandy in places; sometimes carries a lenticular sandstone (Edray) that may occur at top, base or within the shale	75-130	2805
Greenbrier Series		

TOPOGRAPHIC EXPRESSION, MAUCH CHUNK SERIES.

In localities where there are no external modifying influences, such as the presence of overlying Pottsville beds or distortion by folds, the topography of the Mauch Chunk Series usually resolves itself into a series of haphazardly arranged ridges, each of which is capped by a hard sandstone and as a result has a more or less flat crest. From the edges of these crests the descent is usually abrupt until another durable sandstone interrupts the steep slope and forms a shelf. The same succession of steep, shaly slopes and sandstone benches may be repeated several times until the deep valley floor is reached. These valleys are usually narrow because of the apparently youthful cycle of the major streams, and raggedly V-shaped because of the benching of the hillsides.

AREAL EXTENT, MAUCH CHUNK SERIES.

Figure 9 shows at a glance the outcrop of this series in Greenbrier County, while on Map II the same outcrops are delineated in much greater detail. By this figure and map it is evident that approximately 25 per cent. of the surface rocks of the county are of the Mauch Chunk Series. A further examination of Figure 9 and Map II reveals that this series is limited to the portion of the county west of the Greenbrier River and comprising all that area west of the main Greenbrier Limestone belt with the exception of the areas covered by the Pottsville Series as seen on Figure 8 and the area of older rocks along the Williamsburg Anticline.



LATE XIV.—Gullying in the red shale of the Hinton Group of the Mauch Chunk Series, near Rupert. Photo, taken in
See Plate XV for picture in 1931.




PLATE XV.—Gullying in the red shale of the Hinton Group of the Mauch Chunk Series, near Rupert. Gullies have deepened two to three feet in one year. Photo. taken in 1931. See Plate XIV for picture taken in 1930.






PLATE XVII.—Subaerial scouring in the Hinton Group of the Mauch Chunk along Midland Trail (U. S. Route 60). 1.2
s southeast of Crawley.

FIGURE 9

MAP


OF

GREENBRIER COUNTY

SHOWING

OUTCROPS OF

MAUCH CHUNK SERIES



CONTACTS, MAUCH CHUNK SERIES.

The contact of the Mauch Chunk Series with the overlying Pottsville Series and the unconformity that exists between them have been discussed under the description of the latter series. At the base of the Mauch Chunk Series there is not the marked contrast with the underlying Greenbrier Limestone Series as there is with the overlying Pottsville, but the contact is one of gradual change rather than an abrupt break. Considering the two series as a whole there is a large difference, the Mauch Chunk consisting mainly of red shales and sandstones with occasional thin streaks of coal and with the basal portion carrying comparatively thin limestones and shales, while the Greenbrier Series is made up almost entirely of massive limestones. At the contact, however, the two series blend together lithologically as well as paleontologically.

FOSSIL LIFE, MAUCH CHUNK SERIES.

In the Mauch Chunk Series the fossils have changed materially from that at its type locality of reptile tracks and vertebrate remains, to a fauna composed almost entirely of marine shells with an occasional fish tooth, along with a variety of fossil plants. The fossils are distributed throughout the series but increase in number toward the base. No attempt was made to get a complete assemblage from this series, but collections were made at exposures where the fossils were well weathered out. These collections were studied by Professor Dana Wells and their identification will be found in Chapter XIV under the heading Notes on Paleontology. Several loose specimens of *Stigmaria* were collected but none of these appears under this heading. The collections from this series were made primarily from the Avis, Reynolds, and Glenray members.

CORRELATION, MAUCH CHUNK SERIES.

The Mauch Chunk Series of this report is the representative of the Mauch Chunk of Pennsylvania, (No. XI of the earlier Rogers' classification), except that in that State certain calcareous beds are included in the Mauch Chunk that appear to be the equivalent of the part of the Greenbrier Series of

West Virginia. To the southwest the Mauch Chunk correlates, in part, with the Pennington Shale of Virginia but apparently the Pennington includes nothing below the Stony Gap Sandstone and therefore does not include the Bluefield Group which is almost half of the Mauch Chunk in this county.

Reger² has made a very detailed study of the Mauch Chunk Series in Mercer, Monroe, and Summers Counties and in the report cited in the foot-note he described and named a large number of individual beds. In planning the field work for the report on Greenbrier County, it was deemed inadvisable to attempt the detailed work that would make the correlation of all individual beds possible. As a result, only the group boundaries and a few of the more prominent and continuous members are noted in measured sections in Chapter V and in the description of the series in this Chapter.

DESCRIPTION OF MEMBERS, BLUESTONE GROUP.

The lithologic characteristics of individual beds of the Bluestone Group vary rapidly from one place to another so that detailed correlation without almost continuous exposures is very difficult. It is quite clear, however, that from southeast to northwest successively older horizons are in contact with the basal beds of the Pottsville Series.

A coal seam, or more properly, a coaly shale, was noted in this group, that may represent the Hunt Coal of Reger³. One foot of coaly shale, occurring about 80 feet below the base of the Pottsville, was noted on the south end of Little Sewell Mountain and on Big Clear Creek Mountain; what appears to be the same bed was noted 30 feet below the base of the Pottsville. The elevation of these coal exposures as well as the succession of beds above and below them may be seen in the Little Sewell Mountain—South End, and the Big Clear Creek Mountain Sections, published in Chapter V.

In the general vicinity of Rockeliff and Kieffer there are several exposures of a coaly shale that belongs about 100 feet above the Princeton Conglomerate. This coaly shale may be the

²Reger, David B., Mercer, Monroe, and Summers Report, W. Va. Geol. Sur., pp. 291-444; 1926.

³Ibid., pp. 316-317.

equivalent of the Pipestem Coal of Reger⁴. The bed is never more than a few inches thick and is of no economic value.

Approximately 90 per cent. of the group is shale. Most of the shales are deep red in color but a few beds are green, yellow, brown, or dark gray. Some of the shales are calcareous. The sandstone making up the remaining 10 per cent. of the group is usually green, fine-grained, thin-bedded and shaly.

DESCRIPTION OF MEMBERS, PRINCETON GROUP.

PRINCETON SANDSTONE.

The **Princeton Sandstone**, or Princeton Conglomerate of Campbell⁵, is a prominent marker in many parts of Greenbrier County. On each of the two forks of Cherry River in the northern part of the county it is the most prominent bed of the exposed Manah Chunk. In that region, as in most places, it is strongly cemented with limonite and carries the characteristic large pebbles. The sand and pebbles are usually very poorly sorted and this characteristic, when used with some caution, makes it possible to distinguish this sandstone from any other in the region. Near Kieffer the Princeton is almost entirely composed of pebbles. About one mile west of Rupert characteristic drift boulders from this bed may be observed along the Midland Trail.

The character and stratigraphic position of the Princeton Sandstone are shown in the Goddard Mountain, Sims Station, Little Sewell Mountain—West Side, Little Sewell Mountain—South End, Cherry Low Place, Little Rocky Run, Kieffer, Roach Run, Cold Knob—Hinkle Well, Briery Knob, and Alderson Sections, published in Chapter V, and its outcrop is delineated on Map II.

DESCRIPTION OF MEMBERS, HINTON GROUP.

In Greenbrier County there is some evidence of a loss of some of the upper beds of the Hinton Group, by a disconformity. The lithology of the various upper beds is so similar, however, that without additional detailed field work it would be unwise

⁴Ibid., pp. 323-324.

⁵Campbell, M. R., Pocahontas Folio, No. 26, U. S. Geol. Sur., 1896.

CHAPTER VII.

STRATIGRAPHY—MISSISSIPPIAN ROCKS.

GENERAL STATEMENT.

The rocks of the Mississippian Period outcrop in a broad band, trending in a northeast-southwest direction across the center of Greenbrier County. In descending order these rocks are subdivided as follows:

Manch Chunk Series:	Feet.
Bluestone Group.....	80 to 675
Princeton Conglomerate.....	20 to 80
Hinton Group.....	500 to 850
Bluefield Group.....	900 to 1200
Greenbrier Series.....	475 to 700
Maccrady Series.....	60 to 250
Pocono Series.....	200 to 600
Totals	2235 to 4355

The above minimum-maximum thicknesses only apply to the outcropping rocks. It is reasonably certain that a well drilled in the northern or extreme western part of the county would find thicknesses that are less than the minimum figures given above. The description of the groups now follows in descending stratigraphic order.

CORRELATION, MISSISSIPPIAN PERIOD.

In view of the present available information along with conflicting opinions as to the relative ages of different groups, a proper and satisfactory correlation of the lithologic units of the Mississippian, with their equivalents in other areas, will not be obtained until each is studied in its entirety. If one is to compile a geologic and economic report on a large area within a reasonable length of time, it is necessary that corre-

CHAPTER VII.

STRATIGRAPHY—MISSISSIPPIAN ROCKS.

GENERAL STATEMENT.

The rocks of the Mississippian Period outcrop in a broad band, trending in a northeast-southwest direction across the center of Greenbrier County. In descending order these rocks are subdivided as follows:

Mauch Chunk Series:	Feet.
Bluestone Group.....	80 to 675
Princeton Conglomerate.....	20 to 80
Hinton Group.....	500 to 850
Bluefield Group.....	900 to 1200
Greenbrier Series.....	475 to 700
Macerady Series.....	60 to 250
Pocono Series.....	200 to 600
Totals	2235 to 4355

The above minimum-maximum thicknesses only apply to the outcropping rocks. It is reasonably certain that a well drilled in the northern or extreme western part of the county would find thicknesses that are less than the minimum figures given above. The description of the groups now follows in descending stratigraphic order.

CORRELATION, MISSISSIPPIAN PERIOD.

In view of the present available information along with conflicting opinions as to the relative ages of different groups, a proper and satisfactory correlation of the lithologic units of the Mississippian, with their equivalents in other areas, will not be obtained until each is studied in its entirety. If one is to compile a geologic and economic report on a large area within a reasonable length of time, it is necessary that corre-

CHAPTER VII.

STRATIGRAPHY—MISSISSIPPIAN ROCKS.

GENERAL STATEMENT.

The rocks of the Mississippian Period outcrop in a broad band, trending in a northeast-southwest direction across the center of Greenbrier County. In descending order these rocks are subdivided as follows:

Mauch Chunk Series:	Feet.
Bluestone Group.....	80 to 675
Princeton Conglomerate.....	20 to 80
Hinton Group.....	500 to 850
Bluefield Group.....	900 to 1200
Greenbrier Series.....	475 to 700
Macerady Series.....	60 to 250
Poeono Series.....	200 to 600
Totals	2235 to 4355

The above minimum-maximum thicknesses only apply to the outcropping rocks. It is reasonably certain that a well drilled in the northern or extreme western part of the county would find thicknesses that are less than the minimum figures given above. The description of the groups now follows in descending stratigraphic order.

CORRELATION, MISSISSIPPIAN PERIOD.

In view of the present available information along with conflicting opinions as to the relative ages of different groups, a proper and satisfactory correlation of the lithologic units of the Mississippian, with their equivalents in other areas, will not be obtained until each is studied in its entirety. If one is to compile a geologic and economic report on a large area within a reasonable length of time, it is necessary that corre-

On the north side of Joe Knob, 1.05 miles southeast of mouth of Smokehouse Branch; Little Fire Creek Coal?; elevation, 3384' L.

		Ft.	In.
Coal and bone (slate roof).....	0'	1"	
Coal	1	2	
Slate	0	1½	
Coal	0	2½	
Coal and slate.....	0	3½	
Coal (slate floor).....	1	0	10½

The stratigraphic position of the Little Fire Creek Coal is shown in the records of Borings Nos. 5I and 11.

PINEVILLE SANDSTONE.

The **Pineville Sandstone** of Hennen⁵⁸, named from its occurrence near the town of Pineville, Wyoming County, is a prominent sandstone over much of the region of its outcrop. It is generally massive, grayish-white, coarse-grained, with a variable thickness and occasionally its position in the column is occupied by shale. Its thickness and stratigraphic position are shown in the Little Clear Creek, Sims Station, and Big Clear Creek Mountain Sections and in the records of Borings Nos. 5A, 5B, 5G, 6, 11, 13, and 14.

NO. 9 POCAHONTAS COAL.

The **No. 9 Pocahontas Coal** of White⁵⁹ and Hennen⁶⁰, belonging immediately below the sandstone last described was not observed at outcrop but was noted in the records of Borings Nos. 5B, 5C, 5I, 11, and 14. It is generally only a few inches thick and as it occurs only a few feet above the No. 8 Pocahontas Coal, it can be distinguished from No. 8 only if both coals are present.

NO. 8 POCAHONTAS COAL.

The **No. 8 Pocahontas Coal** of White⁶¹ and Hennen⁶² is the basal member of the New River Group as classified in the Gen-

⁵⁸Hennen, Ray V., Wyoming-McDowell Report, W. Va. Geol. Sur., pp. 211-12; 1915.

⁵⁹White, I. C., Vol. II(A), W. Va. Geol. Sur., pp. 102 and 177, 1908.

⁶⁰Hennen, Ray V., Wyoming-McDowell Report, W. Va. Geol. Sur., pp. 212-213, 1915.

thin streaks of coal appears to represent the Little Five Creek Coal on the north end of Sims Mountain (Coal Exposure No. 373 on Map II), shown in the Sims Mountain—North End Section published in Chapter V.

Wm. Bennett Mine—No. 374 on Map II.

On Wm. Bennett land, on Little Sowell Mountain, 2.25 miles south-west of Rupert and 1.55 miles northeast of Meadowvale School; Little Five Creek Coal; elevation, 3285' B.

	Ft.	In.
1. Sandstone, brownish-white, massive, to top of hill	65	0
2. Shale, dark	4	5
3. Coal, bony, hard	0' 4"	
4. Coal, banded	0	11
5. Coal, hard	0	4
6. Coal, soft, (fire clay floor)	1	2
	2	
	2	9

A sample (No. 134PH) was collected from Nos. 3, 4, 5, and 6 of the above section, the analysis of which is published under No. 374 in the Table of Coal Analyses at the end of Chapter XI. The opening was not driven in very far, so the analysis may not truly represent the clean coal.

On the south end of Little Sowell Mountain a blossom of coal was noted at the Little Five Creek horizon as shown in the Little Sowell Mountain—South End Section, published in Chapter V. This exposure is No. 375 on Map II. As shown in the Big Clear Creek Mountain Section in Chapter V, a few inches of coal was noted at exposure No. 375A on Map II, on Big Clear Creek Mountain, that apparently represents the Little Five Creek Coal.

Gauley Coal Land Co. Prospect No. A410—No. 376 on Map II. On the southeast side of Job Knob Branch, 0.7 mile northeast of mouth; Little Five Creek Coal?; elevation, 3583' L.

Ft. 1
In. 6

Gauley Coal Land Co. Prospect No. A253—No. 377 on Map II.

On north side of Joe Knob, 1.05 miles southeast of mouth of Smokehouse Branch; Little Five Creek Coal?; elevation, 3561' L.

Ft. 1
In. 6

Coal 1' 2"

at the top of the Pocahontas Group, was noted at various points in Greenbrier County. In general, it is hard, medium-grained, usually micaceous, bluish-gray to brown, with a thickness ranging between 10 and 40 feet. It is not usually well exposed at outcrop and as a result it is difficult in many places to separate the New River and Pocahontas Groups. In the general vicinity of Duo the Flattop Mountain Sandstone apparently coalesces with the Pierpont Sandstone, cutting out the beds that normally occur between the two sandstones. The character and stratigraphic position of the Flattop Mountain Sandstone are exhibited in the Sims Station Section and in the records of Borings Nos. 5B, 6, 11, 12, 13, and 14. The top of this sandstone is generally 400 to 450 feet below the Sewell Coal.

RIFT SHALE⁶⁶, No. 7 POCAHONTAS COAL⁶⁷, PIERPONT SANDSTONE⁶⁸, ROYAL SHALE⁶⁹, No. 6 POCAHONTAS COAL⁷⁰.

In Greenbrier County and the adjoining parts of Nicholas and Fayette Counties, that part of the Pottsville Series between the base of the Flattop Mountain Sandstone and the top of the Eekman Sandstone often carries three coal beds and may contain as many as five or more. In some places the number of seams depends upon whether a succession of coal, shale, and coal, is a single bed with a parting, or two coals with an intervening shale member. It was observed that at different places first one and then the other of these coals may show the best section. The exact correlation of these seams, over any considerable area, is very difficult and in some cases the correlations are little more than a guess. The correlation of the zone, however, can be established with a reasonable

References to the type localities of the above beds are:

⁶⁶Hennen, Ray V., Wyoming-McDowell Report, W. Va. Geol. Sur., p. 217, 1915.

⁶⁷White, I. C., Vol. II(A), W. Va. Geol. Sur., pp. 102-4, 1908; and Hennen, Ray V., Wyoming-McDowell Report, W. Va. Geol. Sur., pp. 217-18, 1915.

⁶⁸Hennen, Ibid., pp. 218-19.

⁶⁹Krebs, C. E., Raleigh Report, W. Va. Geol. Sur., pp. 366-7, 1916.

⁷⁰White, I. C., Vol. II(A), W. Va. Geol. Sur., pp. 103-4, 1908; and

FLATTOP MOUNTAIN SANDSTONE. The Flattop Mountain Sandstone of White⁶¹ and Hennen⁶², named from its occurrence on the summit of Flattop Mountain.

The character and stratigraphic position of the various members are shown in the General Section of the Pottsville Series published in an earlier page of this Chapter. In Chapter V numerous measured sections show the character of the sediments at various points.

of shale than does the New River Group.

general, the Pocahontas Group contains a greater percentage follows the standard column for southern West Virginia. In sequence of beds can be resolved into a general column that disappearance and to the lenticularity of the coal beds. The hontas Group is extremely difficult, due to its thinning and previous page, the tracing of individual beds of the Pocahontas Group is extremely difficult, due to its thinning and As discussed under "Correlation, Pottsville Series", on a sections and in the records of coal test borings.

is often necessary to ignore this boundary in the measured not clearly defined in many parts of Greenbrier County and it contact of this Group with the overlying New River Group is part of the county but is absent in the northern part. The maximum thickness of slightly over 300 feet in the southwestern Mauch Chunk red shales of the Mississippian, attains a maximum thickness of slightly over 300 feet in the southwestern beginning at the top with the Flattop Mountain Sandstone and extending down through the rock column to the top of the The Pocahontas Group or Lower Pottsville of White⁶³,
DESCRIPTION OF MEMBERS, POCAHONTAS GROUP.

Gold Knob road at Coal Exposure No. 378B on Map II.

appears to be the blossom of the same coal was noted on the records of Borings Nos. 5A, 5B, 5C, and 14. In addition what 378A on Map II), and its stratigraphic position is shown in the in the Big Clear Creek Mountain Section (at Exposure No. partings, but it is usually quite impure. Its blossom is noted this coal may attain a thickness of four feet or over, including general Section given on an earlier page. In Greenbrier County

Pocahontas Coal can usually be recognized.

The **Rift Shale**, belonging between the Flatop Mountain Sandstone and the No. 7 Pocahontas Coal, is no doubt present in Greenbrier, but the exact equivalent of the Rift Shale of the type locality is unknown.

The designation of a coal as the **No. 7 Pocahontas Coal** in this Chapter as well as in Chapter XI, means that the coal occurs between the Flatop Mountain Sandstone and the Beckman Sandstone and that it is apparently above the No. 6 Pocahontas Coal. Numerous measured sections of No. 7 Pocahontas Coal and several chemical analyses are given in Chapter XI. Its stratigraphic position is shown in the Big Clear Creek Mountain and Little Sewell Mountain Sections and in the records of Borings Nos. 5A, 5B, 5C, 13, and 14. A coal that is provisionally correlated as the No. 7 Pocahontas Coal is mined locally at Charmco and on Big Clear Creek Mountain. This seam is known locally as the "Beckley" and is sometimes referred to as the "Dirty Seam" due to the fact that it often has a "high" ash content. The so-called "Dirty Seam" is believed to have a "high" ash content only because it is compared to the extremely pure coal from seams above and below it. Analyses of this "dirty" coal show an ash content of from 6 to 9 per cent, which, in many areas, would be considered a low-ash content.

The **Pierpont Sandstone** was noted in the records of Borings Nos. 6, 11, 12, 13, 14, and 15, all near or east of Duo. In the general vicinity of Charmco, the position of this sandstone is usually occupied by shale that may contain one or more coals.

The **Royal Shale** was noted at several points in connection with the No. 6 Pocahontas Coal. It is dark to black and carries the fossil shell *Lingula*, which is common to most black shales in the Pottsville.

The **No. 6 Pocahontas Coal** is believed to be the most persistent of the coals in this zone. It is mined commercially at the Greenbrier Five Creek Coal Company "Midland" mine

Pocahontas Coal can usually be recognized. The Rift Shale, belonging between the Flat-top Mountain Sandstone and the No. 7 Pocahontas Coal, is no doubt present in Greenbrier, but the exact equivalent of the Rift Shale of the type locality is unknown.

The designation of a coal as the No. 7 Pocahontas Coal in this Chapter as well as in Chapter XI, means that the coal occurs between the Flat-top Mountain Sandstone and the Ekloman Sandstone and that it is apparently above the No. 6 Pocahontas Coal. Numerous measured sections of No. 7 Pocahontas Coal and several chemical analyses are given in Chapter XI. Its stratigraphic position is shown in the Big Clear Creek Mountain and Little Sewell Mountain Sections and in the records of Borings Nos. 5A, 5B, 5C, 13, and 14. A coal that is provisionally correlated as the No. 7 Pocahontas Coal is mined locally at Charney and on Big Clear Mountain. This seam is known locally as the "Beckley" and is sometimes referred to as the "Dirty Seam" due to the fact that it often has a "high" ash content. The so-called "Dirty Seam" is believed to have a "high" ash content only because it is compared to the extremely pure coal from seams above and below it. Analyses of this "dirty" coal show an ash content of from 6 to 9 percent, which, in many areas, would be considered a low-ash content.

The Pierpont Sandstone was noted in the records of Borings Nos. 6, 11, 12, 13, 14, and 15, all near or east of Duo. In the general vicinity of Charney, the position of this sandstone is usually occupied by shale that may contain one or more coals.

The Royal Shale was noted at several points in connection with the No. 6 Pocahontas Coal. It is dark to black and carries the fossil shell *Lingula*, which is common to most black shales in the Pottsville.

The No. 6 Pocahontas Coal is believed to be the most persistent of the coals in this zone. It is mined commercially at the Greenbrier Five Creek Coal Company "Midland" mine

The No. 5 Pocahontas Coal and No. 4 Pocahontas Coal occur so close together and are so similar in Greenbrier County that they are difficult to differentiate. On Big Clear Creek Mountain a coal was noted at Exposure No. 468 on Map II that is believed to represent the No. 5 Pocahontas Coal. A thickness of 0.2 foot of coal was shown at this exposure as published in the Big Clear Creek Mountain Section in Chapter V.

Since the No. 4 Pocahontas Coal is believed to be the more continuous of these two coals, when only one seam is exposed at this horizon, it has been designated as the No. 4 Pocahontas Coal. The following exposures and prospects were noted in Meadow Bluff District:

Coal Exposure No. 469 on Map II is published in connection with the Sims Station Section in Chapter V.

Coal Exposure No. 470 on Map II is published in connection with the Sims Mountain—North End Section in Chapter V.

Coal Prospect No. 471 on Map II.

On the west side of Goddard Mountain, 1.8 miles southeast of East Rainelle; No. 4 Pocahontas Coal; elevation, 2950' B.

Coal	0'	1"
Shale	0	3
Coal	0	1
Shale	0	4
Coal (slate floor)	0	10

1

7

Coal Exposure No. 472 on Map II is published in connection with the Little Sewell Mountain—West Side Section in Chapter V.

Gauley Coal Land Co. Coal Prospect 605A—No. 473 on Map II.

On east side of Mill Creek, 2.45 miles southeast of Charmino and 2.1 miles north of Rupert; No. 4 Pocahontas Coal; elevation, 2908' L.

Ft. 2

In. 6

Coal and slate

of Laurel Creek Seam by mine operators and residents. Numerous measured sections, the results of chemical analyses, and an estimate of the available tonnage are published in Chapter XI. The stratigraphic position of the No. 6 Pocahontas Coal is shown in the Little Clear Creek, Sims Station, Little Sewell Mountain—West Side and South End, Big Clear Creek Mountain, and Sims Mountain—North End Sections, and in the records of Borings Nos. 5C, 6, 11, 12, 13, 14, and 15. Figure 21 shows the probable minable area of the No. 6 Pocahontas Coal and its outcrop is outlined in blue on Map II.

The chemical analyses of the No. 6 Pocahontas seam show it to be an excellent fuel. The volatile matter is low, the ash content is very low, the fusion point of the ash is high, and the B. T. U. is high; all of which are highly desirable qualities for a domestic fuel. This seam is destined to play a more and more important role in the production of coal in Greenbrier County.

The type was set up on Chapter XI (Commercial Coal) before the Chapter on Stratigraphy of the Pottsville was written. Due to an oversight, one of the coal exposures of No. 6 Pocahontas Coal marked on Map II was omitted from that Chapter and as a result it is necessary to include a record of the exposure here:

Coal Exposure No. 414A on Map II.

Meadow Bluff District; on public road, on Sims Mountain, 0.4 mile southwest of Sims School and 1.4 miles east-southeast of Sims (R. R. Station); No. 6 Pocahontas Coal; elevation, 3000' B.

Coal blossom, thickness undetermined.

ECKMAN SANDSTONE.

The Eekman Sandstone of Hennen⁷¹, named from its occurrence at the town of Eekman, McDowell County, is a lenticular, brown to gray, sandstone in Greenbrier County. The bed has no distinguishing characteristics and as a result it was rarely identified in measured sections or cores. Its thickness and stratigraphic position are shown in the General Section and in the records of Borings Nos. 5A and 11.

No. 480 on Map II.

On the north side of Little Clear Creek Mountain, 0.55 mile southwest of mouth of Old Field Branch; No. 4 Pocahontas Coal?; elevation, 3402' L.

Coal	1	
	Ft.	In.
		8

Gauley Coal Land Co. Coal Prospect No. A407—

No. 481 on Map II.

On the north side of Little Clear Creek Mountain, 1.15 miles east of mouth of Smokehouse Branch and 0.8 mile southwest of mouth of Old Field Branch; No. 4 Pocahontas Coal?; elevation, 3398' L.

Coal	1	
	Ft.	In.
		9

Gauley Coal Land Co. Coal Prospect No. A400—

No. 482 on Map II.

On south side of Little Clear Creek Mountain near head of Little Clear Creek, 2.1 miles northeast of mouth of Kuhn Branch; No. 4 Pocahontas Coal; elevation, 3481' L.

Coal (sandstone roof; slate floor)	1	
	Ft.	In.
		7

UPPER POCAHONTAS SANDSTONE.

The Upper Pocahontas Sandstone of Hennen¹, named from its occurrence at Pocahontas, Virginia, appears to be represented in the vicinity of Charcoal and Rainelle. It is generally massive, medium- to coarse-grained, gray to brown, and lenticular. Its character and stratigraphic position are shown in the Sims Mountain—North End Section and in the records of Borings Nos. 11, 13, and 14.

NO. 3 POCAHONTAS "RIDER" COAL.

The No. 3 Pocahontas "Rider" Coal of Hennen¹, was questionably identified at a few localities in Greenbrier County. One of the few exposures was at No. 483 on Map II on the Cold Knob road, where a coal blossom of undetermined thickness was noted at an elevation of 4080' B. Its character and stratigraphic position are shown in the General Section and in the record of Boring No. 5C.

¹Hennen, Ray V., Wyoming-McDowell Report, W. Va. Geol. Surv.

On the east side of Mill Creek, 2.6 miles southeast of Charmeo and 1.89 miles north of Rupert; No. 4 Pocahontas Coal; elevation, 2939' L.

		Ft.	In.
Coal (sandstone roof).....	0'	2"	
Fire clay	3	6	
Coal	0	6	
Bone	0	1	
Coal (fire clay floor).....	1	0	5 3

Gauley Coal Land Co. Coal Prospect 604—No. 475 on Map II.

On the east side of Mill Creek, 2.35 miles southeast of Charmeo and 1.9 miles northwest of Rupert; No. 4 Pocahontas Coal; elevation, 2907' L.

		Ft.	In.
Coal (sandstone roof).....	0'	6"	
Sandstone	4	0	
Coal	0	8	
Bone	0	4	
Coal (fire clay floor).....	2	6	8 0

Gauley Coal Land Co. Coal Prospect 600A—No. 476 on Map II.

On south end of Blg Clear Creek Mountain, 2.35 miles southeast of Charmeo and 1.6 miles northwest of Rupert; No. 4 Pocahontas Coal; elevation, 2907' L.

	Ft.	In.
Coal and slate.....	1	4

Coal Exposure No. 477 on Map II is published in connection with the Big Clear Creek Mountain Section in Chapter V.

Gauley Coal Land Co. Coal Prospect 600—No. 478 on Map II.

On west side of Blg Clear Creek, 1.45 miles north of Rupert; No. 4 Pocahontas Coal; elevation, 3050' L.

	Ft.	In.
Coal and bone.....	2	7

Gauley Coal Land Co. Coal Prospect No. A409— No. 479 on Map II.

On the north side of Little Clear Creek Mountain, 0.6 mile south of mouth of Old Field Branch; No. 4 Pocahontas Coal?; elevation, 3376' L.

	Ft.	In.
Coal (slate roof).....	2'	3"
Slate	2	6
Coal	0	3
Slate	0	4
Coal (fire clay floor).....	1	4 1/2

The No. 2 Pocahontas Coal of Lathrop and White⁸⁰ was tentatively identified in the Sims Station Section and in the records of Borings Nos. 11, 13, and 14. The horizon is marked by black shale and may carry a few inches of coal.

In a few of the coal test borings, a sandstone was reported below what was believed to be the No. 2 Pocahontas Coal. This sandstone may represent the Vivian Sandstone of Henn⁸⁰.

NO. 1 POCAHONTAS COAL.

A coal seam that is believed to represent the No. 1 Pocahontas Coal of Lathrop and White⁸¹ was observed in the vicinity of Charcoal. The following prospect was noted:

Ed. Grafton Coal Prospect—No. 501 on Map II.

On north side of Meadow River, 0.35 mile northwest of Charcoal; No. 1 Pocahontas Coal?; elevation, 2,170' B.
 Coal, very dull lustre, partly concealed, reported by L. E. McClung to have a total thickness of...
 3
 4
 In. Ft.

A sample (No. 140PH) was collected from the portion exposed (about 2 feet thick) for chemical analysis. The analysis is published under No. 501 in the Table of Coal Analyses at the end of Chapter XI.

The character and stratigraphic position of this seam are shown in the Sims Mountain—North End Section (at Exposure No. 500 on Map II), in the Charcoal Section (at Exposure No. 502 on Map II), and in the records of Borings Nos. 11, 12, and 13.

⁸⁰Lathrop, W. A., "The Virginias," p. 97, June, 1884; and White, I. C., W. Va. Geol. Survey, Vol. II, pp. 689-690, 1903; and Vol. II(A), pp. 103-104, 1908.
⁸¹Henn, Ray V., Wyoming-McDowell Report, W. Va. Geol. Surv., pp. 232-234, 1915.
⁸²Lathrop, W. A., "The Virginias," p. 97, June, 1884; and White,

The **No. 3 Pocahontas Coal** of Lathrop and White⁷⁶, named for its occurrence at Pocahontas, Virginia, was observed at numerous points in southwestern Greenbrier County. From a stratigraphic standpoint, it is believed to be the lowest minable coal bed in the territory of this report. This seam has been mined for local use at a few points, but at present (1936), none of these mines are in regular operation. Many measurements of No. 3 Pocahontas Coal, results of analyses, and an estimate of the available tonnage, are published in Chapter XI. The character and stratigraphic position of this seam are shown in the Goddard Mountain, Little Sewell Mountain—West Side, Little Sewell Mountain—South End, Sims Station, Big Clear Creek Mountain, and Little Clear Creek Sections published in Chapter V, and in the records of Borings Nos. 13, 14, and 15, published in Chapter XI. Figure 23 shows the probable minable area of No. 3 Pocahontas Coal and the position of the horizon of the seam on Map II is easily found by reference to the green structure contours and to the table of intervals published in Chapter IV.

LOWER POCAHONTAS SANDSTONE.

The **Lower Pocahontas Sandstone** of Hennen⁷⁷, is thick-bedded, medium-grained, and lenticular in Greenbrier County. Its character and stratigraphic position are shown in the General Section and in the records of Borings Nos. 11, 13, and 14.

NO. 2 "A" POCAHONTAS COAL.

The **No. 2 "A" Pocahontas Coal** of Hennen⁷⁸, was not observed at outcrop but was tentatively identified in the records of Borings Nos. 11 and 14, where it is only a few inches thick.

⁷⁶Lathrop, W. A., "The Virginias," p. 97, June, 1884; and White, I. C., Bull. 65, U. S. Geol. Sur., pp. 203-4, 1891; Vol. II, W. Va. Geol. Sur., pp. 689-690, 1903; and Vol. II(A), W. Va. Geol. Sur., pp. 103-104, 1908.

⁷⁷Hennen, Ray V., Wyoming-McDowell Report, W. Va. Geol. Sur.,

From an economic standpoint, the Pottsville Series is the most important subdivision of the exposed rock column of Greenbrier County. It contains five minable coal seams and at least three other seams show a minable section in certain parts of the county. These are, in descending order, the Sewell, Little Raleigh, Beekley, Fire Creek, No. 6 Pocahontas, and No. 3 Pocahontas. Of these, the Sewell Coal is by far the most important, although the No. 6 Pocahontas Coal is rapidly gaining in importance.

Aside from the coal, however, the rocks of the Pottsville Series contain few materials of economic importance. Many of the sandstones are suitable for various types of masonry structures, but the lack of a near-by market limits their use for this purpose at the present time. Others show sufficient purity to be a source of silica sand suitable for the several uses to which such silica is adapted. The series contains no true fire clays of any consequence in this county.

Jaeger Coal are shown in the Quinwood Section and in the records of Borings Nos. 5E and 8. Its interval above the Sewell Coal varies from 215 to 245 feet.

LOWER JAEGER SANDSTONE.

The Lower Jaeger Sandstone of Hennen², named for its occurrence near Jaeger, McDowell County, appears to be poorly represented in Greenbrier County. When present it is a grayish-brown shaly sandstone rarely more than 15 feet thick. Its character and stratigraphic position are shown in the Quinwood Section and in the record of Boring No. 7.

LOWER JAEGER SHALE.

The Lower Jaeger Shale of Hennen², named from its occurrence near Jaeger, McDowell County, is represented by a dark-brown to gray sandy shale in Greenbrier County. It ranges from 15 to 35 feet in thickness and its stratigraphic position is shown in the records of Borings Nos. 5E, 9, and 10.

HARVEY (CONGLOMERATE) SANDSTONE.

The Harvey (Conglomerate) Sandstone of Campbell², named from the town of Harvey (now Bolt P. O.), Raleigh County, is represented in Greenbrier County by a medium- to coarse-grained, grayish-white or light-brown sandstone. It is a lenticular bed, its interval being often occupied in part or occasionally entirely by sandy shale. The sandstone proper ranges from 20 to 60 feet in thickness, being fine- to medium-grained, gray or grayish-brown and hard. Its stratigraphic position is noted in the Quinwood Section and in the records of Borings Nos. 5L, 5E, 8, 9, and 10. In its more massive phase this sandstone is occasionally apparently coalesced with the Guyandot Sandstone, cutting out the intervening Sandy Huff Shale and the Castle Coal.

SANDY HUFF SHALE.

The Sandy Huff Shale of Hennen², named for its exposure at the mouth of Sandy Huff Branch, McDowell County, is rep-

²Ibid., p. 191.

²Ibid., p. 191-2.

²Campbell, M. R., Raleigh Folio, No. 77, U. S. Geol. Surv., 1902.

The **Hughes Ferry Coal** of White²³, named from its occurrence on the north side of Gauley River, just above the Hughes Ferry bridge, 2.8 miles south of Summersville, Nicholas County, and believed by Hennen²⁴ to represent the **Iaeger Coal** of White²⁵, seldom reaches two feet in thickness in Greenbrier County. It is recorded in the Quinwood Section as two feet thick, but impure, and in the records of Borings Nos. 5E, 7, 8, and 10 as being less than one foot six inches thick. It belongs 275 to 300 feet above the Sewell Coal.

MIDDLE IAEGER SANDSTONE.

The **Middle Iaeger Sandstone** of Hennen²⁶, named from its occurrence at Iaeger, McDowell County, is a shaly lenticular sandstone in Greenbrier County. In its more massive phase, it is a gray to brown, medium- to coarse-grained sandstone, rarely over 30 feet in thickness. Its character and stratigraphic position are shown in the Quinwood Section and in the record of Borings Nos. 5E, 7, and 10. The bottom half of the usual 60-foot interval between the Hughes Ferry Coal and the Lower Iaeger Coal is occupied by a sandy shale.

LOWER IAEGER COAL.

The **Lower Iaeger Coal** of Hennen²⁷, named from its occurrence at Iaeger, McDowell County, is represented in Greenbrier County by an impure coal that varies in thickness with the amount of impurities included in the measurement. It is too thin, impure, and irregular to be classified as a minable seam. The following is one of the few observed exposures of this coal:

Coal Blossom—No. 3 on Map II.

On the Russellville-Nuttersville road, 1.3 miles east of Russellville;
Lower iaeger Coal; elevation, 2365' B.

	Ft.	In.
Coal, poorly exposed.....	0	6

²³White, I. C., Vol. II(A), W. Va. Geol. Sur., pp. 252-253, 1904.

²⁴Hennen, Ray V., Fayette Report, W. Va. Geol. Sur., p. 299, 1919.

²⁵White, I. C., Vol. II(A), W. Va. Geol. Sur., pp. 251-252, 1908.

²⁶Hennen, Ray V., Wyoming-McDowell Report, W. Va. Geol. Sur.,

SEWELL "B" COAL.

The Sewell "B" Coal of Hennen³², named from its occurrence in Wyoming and McDowell Counties, was noted at only one locality in Greenbrier County. In the Quinwood Section it is noted as being six inches thick. Its interval above the Sewell Coal at this point is about 60 feet.

SEWELL "A" COAL.

The Sewell "A" Coal of Hennen³³, named from its association with the Sewell Coal in Wyoming and McDowell Counties, was noted at various points in the territory covered by this report. It is usually one to two feet in thickness, although occasionally thicker, and in many respects it resembles the Sewell Coal in appearance.

While it is generally too thin to be classed as a minable seam, this coal will no doubt eventually furnish some fuel for local use. Its stratigraphic position is shown in the Duo and Quinwood Sections and in the records of Borings Nos. 5E, 5J, 5L, 5M, 7, 8, 9, and 10.

The following two prospects appear to represent the Sewell "A" Coal:

Gauley Coal Land Company Prospect No. 97—

No. 7 on Map II.

On the west bank of Elijah Branch, 1.3 miles northwest of Duo; authority, Gauley Coal Land Company; Sewell "A" Coal?; elevation, 3519' L.

Coal	2	Ft.	8
In.			

Gauley Coal Land Company Prospect—No. 9 on Map II.

On the north side of Beech Ridge, 1.1 miles northeast of Clearco; authority, Gauley Coal Land Company; Sewell "A" Coal; elevation, 3577' L.

Coal	0'	2"		
Coal	0	1		
Slate	2	3		
Coal	2			
In.				

resented at various points throughout the territory of this report. When present, the shale compensates for the varying thickness of the Harvey Sandstone, is dark-gray in color, somewhat sandy, and is frequently cut out by the Harvey Sandstone. Its character and stratigraphic position are shown in the Quinwood Section and in the records of Borings Nos. 8, 9, and 10.

CASTLE COAL.

The **Castle Coal** of Hennen³², named from its occurrence near the town of Castle, Wyoming County, was identified at various points in Greenbrier County. In general it appears to be a high quality of coal but is too thin for mining, rarely reaching two feet in thickness. In the Quinwood Section it is about 115 feet above the Sewell Coal, north of Anjean in Borings Nos. 5E, 5L, and 5M, it is 100 to 120 feet above the Sewell Coal, and east of Dno. what appears to be the same coal is shown in the records of Borings Nos. 7, 8, 9, and 10 as 144 to 167 feet above the Sewell Coal. As mentioned above, the Harvey Sandstone occasionally cuts out this coal.

GUYANDOT SANDSTONE.

The **Guyandot Sandstone** of Campbell³³, named for its occurrence in Wyoming County, is also noted in Greenbrier County, being massive, grayish-white and coarse-grained. It is somewhat lenticular, its interval sometimes being occupied by sandy shale. When present, it ranges in thickness from 30 to 50 feet. Its position is noted in the Quinwood Section and in the records of Borings Nos. 5E, 5L, 5M, 7, 8, 9, and 10. As noted above, the Guyandot Sandstone is sometimes apparently coalesced with the Harvey Sandstone.

SKELT SHALE.

The **Skelton Shale**, of Reger³⁴, named from its occurrence near the village of Skelt, Webster County, was tentatively identified in the Quinwood Section where it is black and 6½ feet thick.

³²Ibid., pp. 193-4.

As pointed out by Price¹¹ the most outstanding example of erratic boulders in coal, yet reported, occurs in the Sewell Coal of Greenbrier County. The following table, taken from the paper referred to above, gives a description of forty boulders. The key number in the left-hand column of the table refer to Plate XII, taken from the same paper:

Erratic Boulders in the Sewell Coal.

The Sewell Coal is the same as the “Ganley Seam” of Dowell County, and is believed by Reger¹⁰ to be the same as the Sharon Coal of Pennsylvania.

marks. mate distances in feet above or below other stratigraphic page 137, a table of intervals is published showing its approximate contours on Map II are based on it and the outcrop of the Sewell Coal is a very important “key-rock.” The green structure contours on Map II are based on it and the outcrop of the In Greenbrier County, as in the adjoining counties, the (particularly the Sewell Coal) of southeastern Nicholas County. able to the public much additional information on the coals been published and their publication in this report makes available to the public much additional information on the coals Nicholas County. Many of these records have not previously contains the records of a number of coal test borings drilled in of the Sewell Coal in Greenbrier County, Chapter XI also contains detailed information concerning the thickness and distribution are published in Chapter XI. In addition to a vast amount of area of Sewell Coal, and an estimate of its available tonnage, data together with Figure 17 showing the approximate minable measured and sampled at numerous mines and prospects. These Nos. 5E, 5J, 5K, 5L, 5M, 6, 7, 8, 9, 10, and 11. This coal was in the Duo and Charneo Sections and in the records of Borings steam and domestic fuel. Its stratigraphic position is shown low in volatile matter the coal has an enviable reputation as a content of sulphur, ash, and phosphorus. Being comparatively

The **Lower Guyandot Sandstone** of Hennen³⁷, named from its occurrence near Wilmore, McDowell Comnty, is a massive, coarse-grained, grayish-white sandstone in northern Greenbrier Comnty but over much of Meadow Bluff District its position in the column is, in whole or in part, occupied by sandy shale. Its character and stratigraphic position are shown in the Duo Section and in the records of Borings Nos. 5I, 6, and 8.

On Fork Mountain, in the vicinity of the abandoned coal mines No. 224 and No. 225, this sandstone has, in places, apparently "cut out" or mashed out the Sewell Coal. In areas where this sandstone is thick it is advisable for coal companies to thoroughly prospect the property before spending any large sums in opening mines.

HARTRIDGE BLACK SHALE.

The **Hartridge Black Shale** of Reger³⁸, named from its occurrence at the mining village of Hartridge, Randolph County, was observed at a number of localities in Greenbrier County. As a rule it is a dark to black, argillaceous, laminated deposit, with abundant plant fossils. Its stratigraphic position is shown in the Duo Section, in the records of Borings Nos. 6, 7, 8, 9, 10, and 11, and it is noted in connection with a number of the special sections of the Sewell Coal in Chapter XI.

This shale is often rich in flora and fauna. Fossil Collections Nos. 6, 13, 85, 142, 143, 145, 146, and 149 were collected from this horizon and in addition to the plant fossils, and *Naiadites elongata*, previously reported in this number, Price found fish remains. These are listed under Collection No. 146 in Chapter XIV.

SEWELL COAL.

The **Sewell Coal** of White³⁹, named from its occurrence on Sewell Mountain, Fayette Comnty, is by far the most important member of the Pottsville present in Greenbrier Comnty and has long been mined extensively on a commercial scale. It

³⁷Ibid., pp. 175 and 196-7.

³⁸Reger, David B., Barbour-Upshur-Western Randolph Report, W. Va. Geol. Sur., pp. 288-290; 1918.

³⁹White, R. H., 1885, p. 65, Pl. C.

Table I.—Boulders from Sewell Coal in Greenbrier, County, West Virginia.

Mine.	Weight Pounds	Dimensions (Inches)			Kind of Rock	Remarks
		Greatest Length	Width	Height		
Argarette ...	161½	19	15	14	Quartzite, gray.	From lower half of coal.
Argarette ...	58	14	12	8	Sandstone, gray, medium-grained. quartzite	From lower half of coal.
Argarette ...	25	11	8	6	Quartzite, gray.	Kldney-shaped
lie	22	10½	7½	5½	Quartzite, gray.	From roof shales.
Argarette ...	10½	9½	4	5	Sandstone, dark, carbonaceous.	Rectangular with rounded edges.
Argarette ...	7½	5½	4	4	Quartzite, dark to white.	Vein-quartz? very compact.
Argarette ...	3¼	5	4¾	3	Sandstone, grayish-white.	Medium grain size.
lie	17¼	7½	6	5½	Quartzite, dark-gray.	Rectangular block with rounded edges.
Argarette ...	17½	11	7	6	Sandstone, gray, fine-grained.	Estimated original weight, 21 pounds.
lie	12¼	7	7	5½	Quartzite, gray.	Polished
lie	10½	10½	3¾	4¾	Quartzite, gray.	Broken in transportation.
Argarette ...	10¾	10½	7	4	Sandstone, conglomerate, gray.	Cut in quartz vein.
lie	10¼	9	4½	4	Quartzite, grayish-white.	Estimated original weight, 12 pounds.
lie	12	8½	4	3¾	Quartzite, gray.	Well polished.
Argaretto ...	8½	6½	5	4½	Quartzite, gray.	Subangular
lie	7¾	6½	6	4	Quartzite, gray.	Elliptical, well polished.
lie	5	5	4	4	Quartzite, gray to maroon.	Subangular, three flat faces.
lie	4½	7½	4½	3	Quartzite, gray.	Bottom side flat.
Argarette ...	4½	5	4	4	Quartzite, gray.	Subangular (see 15, 17), striations.
Argarette ...	4	Quartzite, altered.	Spherical, cut by quartz vein, sericite quartz, and chlorite.
lie	3½	5½	4½	2½	Conglomerate, metamorphosed.	Quartz grains, 5 mm. to very fine. Chlorite, sericite, and secondary quartz, pyrite.

in the amounts to any level in the accumulating material. Such an environment could be explained by a slight uplift near the mouth of a river that formerly had a rather steep gradient."

WELCH SANDSTONE.

The Welch Sandstone of Hennen² named from its occurrence near the town of Welch, McDowell County, is present in the territory of this report and quite often it apparently coalesces with the underlying Upper Raleigh Sandstone, cutting out the Welch Coal. It is usually grayish-white, medium-to coarse-grained, lenticular and ranges in thickness from 20 to 45 feet. Its interval in the column is sometimes occupied by sandy shale. Its stratigraphic position is shown in the Little Rocky Run Section and in the records of Borings Nos. 5M, 6, 9, and 11.

WELCH COAL.

The Welch Coal of White³, named from its occurrence near Welch, McDowell County, was noted at several points in Greenbrier County. In general it is a soft, columnar coal, resembling the Sewell bed in appearance and is rarely over two feet in thickness. North of Quinwood, on Price Fork, and on the headwaters of Hominy Creek in the edge of Nicholas County, this seam may have a thickness of 30 inches or more and it is reported that prospectors have confused its outcrop with that of the overlying Sewell Coal. The coal appears to be of excellent quality and while the seam is too thin and erratic in occurrence to be classified as minable, it will, no doubt, eventually furnish some fuel for local use. As noted in the description of the overlying sandstone, the Welch Coal is often cut out by the Welch Sandstone. The stratigraphic position of the Welch Coal is shown in the Duo Section and in the records of Borings Nos. 5K, 5M, 9, and 11.

UPPER RALEIGH SANDSTONE.

The Upper Raleigh Sandstone of White⁴, named from its occurrence in Raleigh County and being the upper division

²Hennen, Ray V., Wyoming-McDowell Report, W. Va. Geol. Sur., pp. 198-9, 1915.

Subsequent to the writing of the paper cited above, the writers observed literally hundreds of these boulders at the various mines near Quinwood, the Leekie mines near Anjean, the Raine mine near Duo, and at the Cleareo mine near Cleareo. The air-line distance between the Cleareo mine and the Leslie mine is approximately ten miles, thus any theory suggested to account for the occurrence of the boulders must permit widespread distribution.

The following discussion of the transportation of the erratics is a quotation taken from pages 72 and 73 of the paper cited above:

"To account for the presence of boulders in coal, the view previously expressed by most geologists is that they were held in the roots of trees and rafted to their present position. To assume that all of these boulders, especially the larger ones (see Flg. 3, Nos. 1, 2, 3), could have been carried to their present location without considerable quantities of other foreign material, calls for a stretch of the imagination. That some may have been so transported is not doubted. It does not seem logical, however, that a stream sufficiently large to raft trees that could carry some of the boulders here noted would be found in a coal-forming environment. A stream of such size would certainly "wash out" the peat bog itself. Furthermore, the presence of boulders at various horizons in the coal would necessitate the presence of the stream throughout the entire time of the accumulation of the coal-making material.

"The second method of transportation, which is looked upon with favor by some but strenuously objected to by others, is ice. The prevailing opinion seems to be that Pennsylvanian temperatures were not sufficiently low for the formation of ice. Considerable evidence, however, has been advanced to show that during a part of the Pennsylvanian, and specially in the higher altitudes, ice was present for a portion of the year, a view in which the writer concurs. The present boulders, however, do not show the characteristics common to those transported by ice, such as faceted faces or striations. It does not follow, however, that river or shore ice may not have carried these boulders from beaches or along the banks of streams into the Pottsville basin. This, however, would be expected prior to or following the coal accumulation, and could account for the boulders only in the underclay or the overlying sediments. It has already been pointed out that the boulders do not occur at any one particular horizon in the seam, but may be found at any level from the underclay to the roof shales. It should be stated, however, that the majority are found in the lower part of the seam.

"If we may assume immediately preceding the coal accumulation a stream with a very low gradient, along which boulders had been deposited by transporting agents, gradually being encroached upon by coal vegetation, it would be possible for trees by overturning to draw the erratics up into the peat bog. It is known that succeeding

several points for local consumption but are not
there are no actively operating mines in the county. Green-
brier is the only county in the State in which the Little Raleigh
Coal is known to be of minable thickness.

Under "Commercial Coal," Chapter XI, numerous mea-
surements of the thickness of the Little Raleigh Coal are given.
In that Chapter will also be found chemical analyses of coal
from this seam, an estimate of the probable available tonnage
and Figure 19 shows the probable area of minable Little
Raleigh Coal. The stratigraphic position of the seam is shown
in the Charcoal Section and in the records of Borings Nos.
5, 5A, 5C, 5H, 5I, 5K, 5M, 6, and 11. The position of the out-
crop of this seam is not shown on Map II but it may easily be
plotted by reference to the Sewell Coal contour lines, using
the table of intervals published in Chapter IV, page 137.

LOWER RALEIGH SANDSTONE.

The Lower Raleigh Sandstone of White²⁰, or the lower
division of the Raleigh Sandstone of Campbell²⁰, in Greenbrier
County, often attains a development almost equal to that of the
Upper Raleigh ledge, which it resembles in both texture and
physical appearance. Its thickness and stratigraphic position
are shown in the Russellville Section and in the records of
Borings Nos. 5, 5A, 5H, 5K, 5M, 6, 11, and 12.

BECKLEY "RIDER" COAL.

The Beckley "Rider" Coal of Krebs²¹, named from its asso-
ciation with the Beckley Coal in Raleigh County, was tenta-
tively identified in the general vicinity of Anjean. The oc-
currence of this coal is very erratic and its correlation is very
uncertain. It was not observed at outcrop but its stratigraphic
position is shown in the records of Borings Nos. 5C, 5D, 5I,
and 5H.

BECKLEY COAL.

The Beckley Coal of Campbell²², named from its occur-
rence near the city of Beckley, Raleigh County, was opened
at numerous points in Greenbrier County. It is generally

²⁰Ibid., pp. 198-9.

²¹Campbell, M. R., Raleigh Folio, No. 77, U. S. Geol. Surv., 1902.

²²Ibid., pp. 198-9.

of the Raleigh Sandstone of Campbell⁴⁷; is believed by Reger to be the same as the **Sharon Sandstone** of Pennsylvania. It is generally massive, grayish-white to brown, medium- to coarse-grained, occasionally pebbly and forms great cliffs around the mountainsides along its outcrop. It has often acted as a buffer in preserving from erosion a large acreage of coal and wide benches with the Upper Raleigh Sandstone outcropping at the edge are common. Its thickness ranges from 50 to 75 feet and its top varies from 20 to 60 feet below the Sewell Coal. Its character and stratigraphic position are shown in the Charmco, Dno. and Little Rocky Run Sections and in the records of Borings Nos. 5A, 5H, 5M, 6, 9, and 11. The sandstone contains a larger amount of coarse material and is more often conglomeratic in the northeastern part of its outcrop in the county than in the southwest part of the county.

LITTLE RALEIGH "A" COAL.

The **Little Raleigh "A" Coal** of Krebs⁴⁸, named from its occurrence in Raleigh County, appears to be represented at a few points in the county. It is generally impure, less than one foot in thickness, and comes 10 to 20 feet above the Little Raleigh Coal. Its character and stratigraphic position are shown in the Charmco Section and in the records of Borings Nos. 5C, 9, and 11.

LITTLE RALEIGH COAL.

The **Little Raleigh Coal** of White⁴⁹, named from its occurrence in Raleigh County, occurs in the basal part of the 10 to 30 feet of shale that usually separates the Upper and Lower Raleigh Sandstones. It is quite persistent over most of Greenbrier County and in some areas it is definitely of minable thickness. It is generally multiple-bedded, soft, and columnar and ranges in thickness from a few inches to slightly over four feet, usually carrying slate partings when the greater thickness is approached. This coal has been mined at

⁴⁷Campbell, M. R., Raleigh Folio, No. 77, U. S. Geol. Sur.; 1902.

⁴⁸Reger, David B., Barbour, Upshur, and Western Randolph Report, W. Va. Geol. Sur., pp. 292-293; 1918.

⁴⁹Krebs, C. E., Raleigh Report, W. Va. Geol. Sur., pp. 322 and 361;

position and character of the Quinnimont shale in the Sims Mountain—North End Section and in the records of Borings Nos. 5A, 5B, 5D, 5F, 5G, 5H, 5I, 5K, 6, 12, 13, and 14.

QUINNIMONT SHALE.

The Quinnimont Shale of Campbell¹, named for its occurrence near the town of Quinnimont, Fayette County, occupies, together with the sandstone last described, the interval between the Beckley and Five Creek Coals. In general it is dark gray and sandy with a variable thickness that in part compensates for the variations in thickness of the overlying sandstone. Due to its erratic thickness and sandy character it was rarely identified by name, in the measured sections or cores.

FIRE CREEK COAL.

The Fire Creek Coal of White² was named from its occurrence in the vicinity of Five Creek and Quinnimont, Fayette County, where it has long been mined on a commercial scale. In general it is multiple-bedded, soft, and ranges from a thin film to seven feet in thickness. There are no mines operating in this seam in Greenbrier County at the present (1936) time. On Little Clear Creek Mountain there is a large area of Five Creek Coal with a thickness in excess of five feet but over much of the county it rarely exceeds three feet. The stratigraphic position of the Five Creek Coal is shown in the Sims Station and Big Clear Mountain Sections and in the records of Borings Nos. 5, 5A, 5B, 5D, 5F, 5H, 5I, 13, and 14. In Chapter XI, numerous measured sections are given, as well as chemical analyses and an estimate of the available tonnage. Figure 20 shows the probable area of minable coal and the position of the outcrop of this seam delineated on Map II.

The Fire Creek Coal, as herein correlated, should not be confused with the seam that many of the residents of the county call "Five Creek." In western Greenbrier and eastern Fayette Counties considerable coal is being produced from the No. 6 Poahontas Coal. It is this seam that is usually called "Five Creek" by the residents of the area.

multiple-bedded, soft, columnar and at some points it bears a striking resemblance to the Sewell bed in appearance. This similarity in appearance has led some prospectors to believe that the uppermost coal prospected on Little Clear Creek Mountain is the Sewell Coal. A comparison of the records of Borings No. 11 and No. 13 indicates clearly that such is not the case. The Sewell Coal occurs in boring No. 11 at a depth of 53 feet and the top of the Mauch Chunk is shown at a depth of 828 feet. In boring No. 13 the top of the Mauch Chunk reds is shown at a depth of only 505½ feet and the coal in question was opened some nine feet above the top of the boring. The correlation indicated by these borings was verified by the junior author by tracing the outcrop of the various sandstones from the location of Boring No. 11, to Grassy Knob, thence along Old Field Mountain to Little Clear Creek Mountain.

Numerous measured sections, results of chemical analyses, and an estimate of the available tonnage of the Beekley Coal are published in Chapter XI. The probable area of minable Beekley Coal is shown on Figure 19, and its stratigraphic position is shown in the records of Borings Nos. 5, 5A, 5B, 5C, 5D, 5F, 5G, 5H, 5K, 6, 11, 12, and 14. Its outcrop is not delineated on Map II, but it is easily plotted thereon by use of the green structure contours and the table of intervals published in Chapter IV.

In the vicinity of Anjean the occurrence of this seam is quite erratic or its interval below the Sewell Coal is extremely variable. No. 7 Pocahontas Coal, which will be described on a subsequent page, is locally (erroneously) called the "Beekley" Coal.

QUINNIMONT SANDSTONE.

The **Quinnimont Sandstone** of White⁵³, named from its occurrence near the town of Quinnimont, Fayette County, was noted at a number of points in Greenbrier County. It is generally a hard, gray, massive, medium-grained sandstone and it is particularly hard and quartzitic in the vicinity of Anjean. Its thickness is quite variable. The stratigraphic

The **Little Fire Creek Coal of White**⁵⁶, named from its association with the coal last described, is represented in Greenbrier County by a multiple-bedded, soft, columnar coal that varies in thickness from a few inches to slightly over two feet. It is frequently absent or represented by black shale. On Boggs Knob and Little Sewell Mountain small truck mines have been opened in this seam. As noted above it is quite irregular in occurrence and thickness and this together with the small area in which the seam appears to average even two feet thick prevents its classification as minable. This coal will, no doubt, continue for some time to furnish a small amount of fuel for local use. The following openings in the Little Fire Creek Coal were noted in Meadow Bluff District:

Meadow River Lumber Company Mine—No. 371 on Map II.

On west side of Boggs Knob, 2 miles southeast of Sims; Little Fire Creek Coal; elevation, 3255' B.

		Ft.	In.
Coal, bony (slate roof).....	0' 4"		
Coal, clean, columnar (shale floor) 2 0		2	4

A sample (No. 87PH) was taken from the above section, the analysis of which is published under No. 371 in the Table of Coal Analyses at the end of Chapter XI.

Hennen⁵⁷, visited the same mine about 1918 and measured and sampled the coal. He reports the following:

		Ft.	In.
Coal, bony, 6" to.....	0' 8"		
Coal, soft.....	1 11	2	7

A sample (No. 925H) was collected by him, the results of which are republished under No. 371 in the Table of Coal Analyses at the end of Chapter XI.

On the north side of Boggs Knob this coal has a thickness of two feet at **Coal Prospect No. 372 on Map II**, with an elevation of 3200' B.

The Five Creek Coal, as herein correlated, should not be confused with the seam that many of the residents of the Fayette Counties considerable coal is being produced from the No. 6 Poehontas Coal. It is this seam that is usually called "Fire Creek" by the residents of the area.

The Fire Creek Coal, as herein correlated, should not be confused with the seam that many of the residents of the Fayette Counties considerable coal is being produced from the No. 6 Poehontas Coal. It is this seam that is usually called "Fire Creek" by the residents of the area.

FIRE CREEK COAL.

The Fire Creek Coal of White² was named from its occurrence in the vicinity of Fire Creek and Quinimont, Fayette County, where it has long been mined on a commercial scale. In general it is multiple-bedded, soft, and ranges from a thin film to seven feet in thickness. There are no mines operating in this seam in Greenbrier County at the present (1936) time. On Little Clear Creek Mountain there is a large area of Fire Creek Coal with a thickness in excess of five feet, but over much of the county it rarely exceeds three feet. The stratigraphic position of the Fire Creek Coal is shown in the Sims Station and Big Clear Creek Mountain Sections and in the records of Borings Nos. 5, 5A, 5B, 5D, 5F, 5H, 5I, 11, 13, and 14. In Chapter XI, numerous measured sections are given, as well as chemical analyses and an estimate of the available tonnage. Figure 20 shows the probable area of minable coal and the position of the outcrop of this seam is delineated on Map II.

QUINIMONT SHALE.

The Quinimont Shale of Campbell³, named for its occurrence near the town of Quinimont, Fayette County, occupies together with the sandstone last described, the interval between the Beekley and Fire Creek Coals. In general it is dark gray and sandy with a variable thickness that in part compensates for the variations in thickness of the overlying sandstone. Due to its erratic thickness and sandy character it was rarely identified by name, in the measured sections or cores.

FIGURE 8

MAP

OF


GREENBRIER COUNTY

SHOWING

OUTCROPS OF

POTTSVILLE SERIES

- KANAWHA GROUP
- NEW RIVER GROUP
- POCAHONTAS GROUP



The Pottsville Series of the Pennsylvanian, representing the base of this System and lying just over the Mauch Chunk Series of Mississippian age, comprises the youngest formation of the region. The Pottsville Series was first named and described by Pennsylvania geologists from its occurrence at Pottsville, eastern Pennsylvania, where it is composed of numerous conglomeratic sandstones accompanied by anthracite coal seams. Later it was subdivided by Dr. I. C. White into the Upper Pottsville or Kanawha Group, the Middle Pottsville or New River Group, and the Lower Pottsville or Poehontas Group. Custom has sanctioned the use of the geographic names last mentioned because of their relation to the Kanawha and New River coal fields of southern West Virginia and Virginia. The Pottsville Series is represented in western Greenbrier County by the basal members of the Kanawha Group, the New River Group, and the Poehontas Group.

At the base of the Pottsville Series there is an unconformity, general and as extensive as the series itself. North and west of southern West Virginia, in addition to the thinning of the beds between the coal seams, a greater and greater number of the basal members of the Pottsville Series are absent. In the north and northwest part of the territory of this report the Poehontas Group is entirely absent, and it is doubtful if all the basal members of the New River Group are present¹.

The following quotation from Price² summarizes the history of the deposition of the Pottsville rocks:

"At the close of the Mauch Chunk time there existed a broad low coastal plain, bordering a vast expanse of shoals, ferruginous mud-flats, with ripple-marks, mud-cracks, rain-prints, and in some localities fossil tracks. This was followed by an orogenic movement producing subsidence under loading, with stability at intervals, sufficient for a growth of vegetation to form coals. The early subsidence was most pronounced along the east shore with a westward transgression of the sea."

¹It may be stated here that the report of E. V. d'Inville for the Gauley Coal Land Company made in 1900 was very complete for the Gauley or Sewell Coal seam over much of Greenbrier and Nicholas Counties, but as stated by him, the account of other seams was "wholly

evidenece of an ancient drowned valley of Mauch Chunk time, or, interpreted another way, it is evidenece of an aneient monadnock on the Mauch Chunk peneplain. It is the only example in southeastern West Virginia to come to the writers' attention, of major topographie relief of that period.

An exeellent diseussion of the nomenelature of the Pottsville Series is given by Reger³. The member names used in southern West Virginia are used in this report as shown in the following general section.

General Section, Pottsville Series, Greenbrier County.

	Thiekness. Feet.	Total. Feet.
Kanawha Group (250')		
Fire elay, impure, and shale (not observed in Greenbrier)	5 to 10	10
Sandstone, Upper Gilbert (not observed in Greenbrier)	0 to 10	10
Coal, Glenalum Tunnel (not observed in Greenbrier)	0	10
Sandstone, Lower Gilbert, massive, gray.....	30 to 80	90
Coal, Gilbert "A" (not observed in Greenbrier)	0	90
Shale, Gilbert, dark, laminated.....	0 to 4	96
Coal, Gilbert, soft, columnar (not observed in Greenbrier)	2 to 4	100
Shale, sandy.....	0 to 19	119
Sandstone, Dotson, massive, gray.....	20 to 65	184
Coal, Douglas "A", (not observed in Greenbrier)	0 to 1	185
Shale, sandy, dark.....	5 to 13	198
Coal, Douglas, often slaty (not observed in Greenbrier)	0 to 2	200
Sandstone, Lower Dotson, masslve, gray.....	10 to 25	225
Shale, Douglas, dark, sandy, laminated.....	5 to 13	238
Coal, Lower Douglas, soft, often slaty (not observed in Greenbrier).....	0 to 2	240
Shale, gray and sandy.....	0 to 10	250
New River Group (940')		
Sandstone, Upper Nuttall, masslve to heavy and curront-bedded, grayish-white to brown....	70 to 50	300
Shale, dark, sandy.....	0 to 20	320
Coal, laeger "B", multiple-bedded, soft.....	1 to 0	320

	Feet.		Feet.
Coal, Little Fire Creek, multiple-bedded, soft, columnar	0 to	2	1132
Sandstone, Pineville, massive to current-bedded	30 to	50	1182
Shale, sandy.....	20 to	0	1182
Coal, No. 9 Pocahontas, multiple-bedded, soft, columnar	2 to	0	1182
Shale and sandstone mixed.....	15 to	4	1186
Coal, No. 8 Pocahontas, impure, soft, columnar	0 to	4	1190
Pocahontas Group (306')			
Sandstone, Flattop Mountain, massive to current-bedded, medium-grained, micaceous, bluish-gray to brown.....	10 to	40	1230
Shale, Rift, dark-gray, with argillaceous and siliceous layers.....	10 to	0	1230
Coal, No. 7 Pocahontas, multiple-bedded, soft, columnar	0 to	4	1234
Shale, gray and sandy.....	0 to	5	1239
Sandstone, Pierpont, massive to current-bedded, medium-grained, hard, micaceous, bluish-gray to light-gray.....	40 to	20	1259
Shale, sandy, alternating with sandstone.....	0 to	10	1269
Shale, Royal, buff, sandy, with fresh- or brackish-water fossil fauna.....	5 to	0	1269
Coal, No. 6 Pocahontas, multiple-bedded, soft, columnar	0 to	5	1274
Shale, sandy.....	0 to	5	1279
Sandstone, Eckman, massive to current-bedded, medium-grained, buff, to bluish-gray.....	40 to	20	1299
Coal, No. 5 Pocahontas, soft, columnar.....	0 to	1	1300
Shale, sandstone, and dark shale, with plant fossils abundant.....	0 to	20	1320
Coal, No. 4 Pocahontas, multiple-bedded, soft, columnar	0 to	2	1322
Shale, sandy.....	0 to	5	1327
Sandstone, Upper Pocahontas, massive to heavy-bedded, medium-grained to coarse.....	0 to	30	1357
Coal, No. 3 Pocahontas "Rider".....	2 to	0	1357
Shale, dark, with plant fossils abundant, and fresh- or brackish-water fossil fauna.....	0 to	10	1367
Coal, No. 3 Pocahontas, multiple-bedded, soft, columnar	0 to	5	1372
Shale, gray, and sandy.....	10 to	0	1372
Sandstone, Lower Pocahontas, generally massive, medium-grained.....	0 to	25	1397
Shale, gray and sandy.....	10 to	0	1397
Coal, No. 2 Pocahontas, multiple-bedded, soft...	0 to	2	1399
Shale, gray.....	0 to	15	1414
Sandstone, Vivian, massive, bluish-gray, medium-grained, lenticular	0 to	30	1444
Coal, No. 1 Pocahontas, generally single-bedded, soft, columnar.....	0 to	2	1446

	Feet.	Feet.
Sandstone, Lower Nuttall, massive, medium-grained, gray to brown.....	50 to 100	420
Coal, laeger "A", slaty.....	1 to 0	420
Shale, Upper laeger, dark.....	50 to 40	460
Coal, Hughes Ferry, single-bedded.....	1 to 2	462
Shale, sandy.....	0 to 5	467
Sandstone, Middle laeger, grayish-white, medium-grained	10 to 45	512
Shale, sandy	40 to 10	522
Coal, Lower laeger, double-bedded.....	0 to 2	524
Fire clay shale.....	0 to 1	525
Sandstone, Lower laeger, gray and brown.....	5 to 15	540
Shale, Lower laeger, dark-gray.....	15 to 35	575
Sandstone, Harvey Conglomerate, medium-grained to coarse, grayish-white to brown, lenticular	60 to 20	595
Shale, Sandy Huff, dark-gray.....	0 to 25	620
Coal, Castle, single-bedded, soft, columnar.....	2 to 0	620
Sandstone, Guyandot, massive, grayish-white, coarse-grained	30 to 50	670
Shale, Skelt, sandy, and dark.....	0 to 5	675
Coal, Sewell "B", slaty, impure.....	0 to 3	678
Shale, sandy	20 to 30	708
Coal, Sewell "A", double-bedded, soft, columnar	0 to 2	710
Sandstone, Lower Guyandot, massive, coarse-grained, grayish-white.....	10 to 30	740
Shale, Hartridge, dark, with plant fossils carrying fresh- or brackish-water fossil shells	0 to 5	745
Coal, Sewell, generally double-bedded, soft, columnar	2 to 7	752
Shale, gray, sandy, lenticular.....	40 to 5	757
Sandstone, Welch, massive to current-bedded, grayish-white	20 to 45	802
Shale, dark, argillaceous, lenticular.....	0 to 3	805
Coal, Welch, multiple-bedded, soft, columnar...	0 to 2	807
Shale, gray, sandy.....	0 to 5	812
Sandstone, Upper Raleigh, heavy to current-bedded, grayish-white to brown.....	75 to 50	862
Coal, Little Raleigh "A", impure.....	0 to 1	863
Shale, sandy, lenticular.....	0 to 25	888
Coal, Little Raleigh, multiple-bedded, soft, columnar	4 to 2	890
Shale, sandy, lenticular.....	15 to 5	895
Sandstone, Lower Raleigh, massive to current-bedded, lenticular.....	50 to 100	995
Coal, Beekley "Rider".....	0 to 2	997
Shale, dark-gray, argillaceous, lenticular.....	0 to 20	1017
Coal, Beekley, multiple-bedded, soft, columnar	0 to 3	1020
Sandstone, Quinnimont, lenticular.....	0 to 70	1090
Shale, Quinnimont, dark-gray, siliceous to argillaceous, laminated, lenticular.....	40 to 5	1095
Coal, Fire Creek, "Quinnimont", multiple-	0 to 7	1100

at many localities within the area. These, however, are rather small in extent and do not warrant description.

FOSSIL LIFE.

In the Pottsville Series throughout southern West Virginia, fossil plants are abundant and well preserved in the shales associated with the coals, and often in the sandstones. They have been widely studied by many authorities⁴.

In contrast to the plant life is the scarcity of marine, brackish- or fresh-water fauna. As pointed out by Lueke⁵, erroneous conclusions as to conditions of deposition may be drawn from the lack of fossils.

Fossil shells of the genus *Lingula* have been reported from roof shales of almost every persistent coal of the New River and Pocahontas Groups. In Greenbrier County Price has noted fish teeth, scales, and coprolites in the roof shales of the Sewell Coal.

CORRELATION, POTTSVILLE SERIES.

As pointed out under the "General Account," the detailed subdivisions of that part of the Pottsville Series remaining in Greenbrier County follow the established nomenclature for southern West Virginia. Synonymous names and a reference to the type locality will be given in the description of each member on subsequent pages.

The problem of the proper correlation of individual beds within the Pottsville Series in Greenbrier County is very difficult. The chief causes of the difficulties and some of the specific areas in which they apply may be summarized as follows: (1) The rapid thinning of the Pottsville measures in a north and northwest direction. (2) Paucity of fossil fauna; as noted above, the Pottsville is devoid of any significant fossil fauna. (3) Similarity of the interval between many of the coals and similarity of the lithologic characteristics of

⁴See Vol. V(A), Part II, W. Va. Geol. Sur., 1913, for a discussion of many of these plants by David White.

POCAHONTAS EXPOSITION.

The topography of the Pottsville Series in the area, as in all other parts of the State in which the series outcrops, is, in a large degree, rough, rugged, and mountainous. The thick, massive sandstones and conglomerates, cut across by streams, leave standing huge cliffs which make bold shoulders along their valleys and from which much talus accumulates on the slopes. This is reflected by the coal-test borings in that they always report from 10 to over 40 feet of "surface" or "boulders and clay." In regions not cut across by roads, this talus material masks the bed-rock, and coal prospecting must be done by coring or by digging deep trenches. Invariably the series produces a very poor soil unfit for cultivation, so that the land is seldom cleared.

CONTACTS AND UNCONFORMITIES.

The contact of the New River Group of the Pottsville Series with the overlying Kanawha Group is at the top of the prominent Upper Nuttall Sandstone. This is a good horizon at which to make the division because the sandstone is very massive and persistent, and there is little evidence of widespread disconformity.

The contact of the New River Group with that of the underlying Pocahontas Group is not so well marked in this region. It is at the base of the No. 8 Pocahontas Coal and at the top of the Flattop Mountain Sandstone.

In this area, as in other parts of the State, there is evidence of a marked unconformity at the contact of the Pottsville Series with that of the Mauch Chunk. That a considerable period of time elapsed from the close of the latter period before the deposition of Pottsville sediments was begun, as mentioned under the "General Account" above, is also evidenced by the marked contrast in the conditions accompanying sedimentation, the soft, red shales of the Mauch Chunk being succeeded by the heavy, coarse, gray to grayish-white and current-bedded sandstones and coal seams of the Pottsville.

Slight local disconformities, revealed by the temporary

The **Lower Gilbert Sandstone** of Hennen and Reger⁷ was tentatively identified near Hanging Rock, just east of the common corner of Nicholas, Webster, and Greenbrier Counties. At the one point observed, it is a massive, grayish-white, coarse-grained sandstone. It appears to cap several knobs near the locality mentioned.

The **Gilbert "A" Coal** of Hennen⁸, named for its occurrence in McDowell County, was not observed in Greenbrier County.

GILBERT SHALE.

The **Gilbert Shale** of Hennen⁹, named from its occurrence in Wyoming County, was observed at the same locality as the sandstone described above. It was poorly exposed and could not be examined in detail.

The **Gilbert Coal** of Hennen and Reger¹⁰ was not observed in Greenbrier County but it is no doubt present over a small area in the extreme northern part of the county. It is described as minable in the reports for the adjoining counties but due to lack of information it is not so recognized here.

DOTSON SANDSTONE.

The **Dotson Sandstone** of Campbell¹¹, named from its occurrence at Wyoming Station (formerly Dotson), Mingo County, was noted in the extreme northern part of the county. At the few points it was observed it was a massive gray sandstone with a thickness ranging between 20 and 65 feet.

The **Douglas "A" Coal** and the **Douglas Coal** of Hennen¹², were not observed in Greenbrier County.

⁷Hennen, Ray V., and Reger, D. B., Logan and Mingo Report, W. Va. Geol. Sur., p. 219; 1914.

⁸Hennen, Ray V., Wyoming and McDowell Report, W. Va. Geol. Sur., p. 167, 1915.

⁹Ibid., p. 168.

¹⁰Hennen, Ray V., and Reger, David B., Logan and Mingo Report, W. Va. Geol. Sur., pp. 221-222; 1914.

¹¹Campbell, M. R., Tazewell Folio, No. 44, U. S. Geol. Sur., 1898.

apply to all of the Pottsville beds throughout the area of their outcrop in the county. (+) Inaccuracies in the topographic base map. This was particularly confusing on the headwaters of Brown Creek and just west of Charuco. (5) Lenticular beds and variations in the interval between coals. These factors apply more or less to all of the Pottsville rocks in Greenbrier County but they are particularly confusing in correlating the Pocahontas coals near Charuco and in correlating the Beckley and Fire Creek Coals near Anjean.

The coal test borings drilled for the Bellwood Coal Company in Fayette County provide excellent illustrations of the lensing of the coal beds, in the interval between the No. 9 and No. 6 Pocahontas Coals. Special attention is called to the record of Boring No. 151 in which is shown four seams of coal two feet or more in thickness and one other coal an inch thick in an interval of only 71 feet. Attention is also called to the record of Boring No. 148 which was drilled less than 0.2 mile north of No. 151. In the same interval that carried five coals in No. 151, the record of No. 148 shows only two coal beds and only one of those is over two feet in thickness.

Under such conditions correlation of individual seams can not always be made with certainty.

DESCRIPTION OF MEMBERS, KANAWHA GROUP.

The Kanawha Group of White, comprising the upper portion of the Pottsville Series, is the youngest group of stratified rocks remaining in Greenbrier County, and is represented by its basal portion, with a maximum thickness of 250 feet. Except for small isolated patches on the tops of some of the mountains, the rocks of this group are confined to the extreme northern part of the county. Exposures of the rocks of this group, other than the sandstone ledges, were seldom seen and the character of the intermediate horizons is therefore quite indefinite.

underlies the New River Group, is absent by means of an unconformity and the New River Group rests directly on the Mauch Chunk. In thickness the series ranges from about 600 feet in the northern part of the county to about 950 feet at the Fayette-Greenbrier County line. In common with the rest of the Pottsville the greatest thinning is toward the northwest.

Of the four minable coals in the group, the Sewell Coal is by far the most important from both an economic and stratigraphic standpoint. In Greenbrier County this coal bed is the most persistent member of the Pottsville and is invaluable in unraveling the stratigraphy of the western third of the county. A description of the Sewell seam and of the three other minable seams—Little Raleigh, Beekley, and Fire Creek—are given on subsequent pages of this Chapter and in Chapter XI.

UPPER NUTTALL SANDSTONE.

The Nuttall Sandstone of Campbell and White¹⁷, later termed the Upper Nuttall Sandstone by Hennen¹⁸, named for its occurrence along New River, Fayette County, between Nuttallburg and Gauley Bridge, where it is a conspicuous cliff-forming ledge, is present over a small area in Greenbrier County. It is a medium-grained, gray to brown, massive sandstone varying in thickness from 50 to 70 feet. It is generally found only on the mountain tops, some of the best exposures being on Shellecamp Ridge, Little Beech Knob, and Buck Knob. Its stratigraphic position is shown in the General Section and in the Quinwood Section. The interval from the base of the Sewell Coal to the top of the Upper Nuttall Sandstone ranges from about 450 feet in the northern end of the county to slightly over 500 feet near Duo.

IAEGER "B" COAL.

The Iaeger "B" Coal of Hennen¹⁹, belonging in the interval between the Upper and Lower Nuttall Sandstones, appears to

¹⁷Campbell, M. R., Raleigh Folio, No. 77, U. S. Geol. Sur., Dec. 1901.

White, I. C., Bull. 65, U. S. Geol. Sur., p. 200, 1891; Vol. II, W. Va. Geol. Sur., pp. 616 and 665, 1903; and Vol. II(A), W. Va. Geol. Sur., pp. 253-254, 1908.

¹⁹Hennen, Ray V., Fayette Report, W. Va. Geol. Sur., p. 295, 1919.

The Lower Dotson Sandstone of Hennen¹³, named from its relationship to the Dotson Sandstone, appears to be present on a few of the high knobs in Meadow Bluff District and was noted at several places north of the North Fork of Cherry River. In appearance it is similar to the other Pottsville sandstones in that it is massive, gray, and coarse-grained. The thickness of this sandstone ranges from 10 to 25 feet.

DOUGLAS SHALE.

The Douglas Shale of Hennen¹⁴, named from its occurrence near the town of Douglas, McDowell County, where it comes just below the Lower Dotson Sandstone (erroneously termed Lower Nuttall in the Report cited) and described as bearing marine or brackish-water fossils, is present but apparently non-fossiliferous in Greenbrier County. It was noted on several high knobs in Meadow Bluff District but was not examined in detail. The total thickness of shale between the overlying Lower Dotson Sandstone and the top of the Nuttall Sandstone appears to be about 25 feet. This may have included the Lower Douglas Coal of Hennen¹⁵ which was not definitely recognized in Greenbrier County.

DESCRIPTION OF MEMBERS, NEW RIVER GROUP.

The New River Group of Fontaine¹⁶, named from its development along New River in Fayette and Raleigh Counties, West Virginia, comprises approximately two-thirds of the Pottsville Series of Greenbrier County. It is defined as including the beds between the top of the Upper Nuttall Sandstone and the base of the No. 8 Pocahontas Coal. In the northern part of the county the Pocahontas Group, that normally

¹³Hennen, Ray V., Fayette Report, W. Va. Geol. Sur., pp. 274-275; 1919.
¹⁴Hennen, Ray V., Wyoming and McDowell Report, W. Va. Geol. Sur., pp. 183-4; 1915.
¹⁵Ibid., pp. 184-185.
¹⁶Fontaine, Wm. M., The "Great Conglomerate" on New River, West Virginia, Amer. Jour. Sci., Third Series, Vol. VII, 1874, pp. 459-



PLATE X.—Pottsville Sandstone (Guyandot) on Little Rocky Run of South Fork of Cherry River.

be represented in Greenbrier County by a very impure coal of black shale that is usually less than one foot in thickness. Its horizon is noted in the Quinwood Section and in Borings No. 7.

LOWER NUTTALL SANDSTONE.

The Lower Nuttall Sandstone of Hennen²⁰, named for its occurrence along New River, Fayette County, where it occurs a few feet below the Upper Nuttall Sandstone and is a prominent cliff-forming ledge, is often quite shaly in Greenbrier County. As a result its outcrop was seldom noted, being recorded in only the Quinwood Section. At that locality it is a gray to brown, medium-grained sandstone, 30 feet thick. It is recorded in Borings Nos. 5E, 7, 8, and 10 with a thickness of 50 to 95 feet, the lower part of the bed being somewhat shaly.

Since its top belongs immediately below the Jaeger "B" Coal, its interval above the Sewell Coal is 400 to 450 feet.

JAEGER "A" COAL.

The Jaeger "A" Coal of Hennen²¹, named from its occurrence in McDowell County, where it comes only a few feet under the Lower Nuttall Sandstone, was not observed as a surface exposure, but is noted in Borings Nos. 7, 8, and 10, being only a few inches in thickness. It belongs 330 to 350 feet above the Sewell Coal and 40 to 50 feet above the Hughes Ferry Coal.

UPPER JAEGER SHALE.

The Upper Jaeger Shale of Hennen²², named from its occurrence in McDowell County, where it occupies the interval between the Jaeger "A" Coal and the Hughes Ferry (Jaeger) Coal, is represented in Greenbrier County by a dark to gray, sandy shale 40 to 50 feet thick. Its thickness and stratigraphic position are shown in the Quinwood Section and in records of Borings Nos. 8 and 10.

²⁰Hennen, Ray V., Fayette Report, W. Va. Geol. Surv., p. 297, 1919.



PLATE XI.—View of operations in the Sewell Coal just east of Quinwood.




PLATE XIII.—Fossil plant showing attached rootlets. Pottsville Sandstone (Upper Raleigh) at Duo.



PLATE XII.—Erratic boulders from the Sewell Coal, Greenbrier County. No. 28 is a granite that was broken by the miners. Photo. by Paul H. Price.

		Feet.	Feet.
Shale, Patton, weathered horizon of very irregular sandy chert with red clay (erosion surface?, 5-10')..	10		140
Limestone, light-blue, abundant light-gray chert, some nodules quite large, numerous battered bryozoa and erinoid stems.....	30'		
Shale, olive-green and sandy (1925' B.)	10	Sinks Grove	60
Limestone, blue-black, massive, good	5		
Limestone, weathers sandy.....	5		
Limestone, light-blue, weathers white, clay seams.....	5		
Limestone, light-yellow, argillaceous	5		
Limestone, light-gray, massive, black chert (top, 1900' B.).....	5'	Hillsdale	85
Limestone, gray-blue, massive, one band of black nodular chert.....	12		
Shale, yellow, sandy, limy.....	3		
Limestone, light-blue, yellowish, sandy, shaly, with chert at base....	15		
Limestone, much weathered, partly concealed, with chert remaining in place, to Second Creek.....	50		
			285

MEASURED SECTIONS, FORT SPRINGS DISTRICT.

Fort Springs is a small triangular-shaped district lying just north of Irish Corner District. The outcropping rocks range from the Bluefield Group of the Mauch Chunk to and including the upper part of the Pocono Series.

Hawver School Section—West.

Fort Springs—Blue Sulphur District; starting $\frac{1}{2}$ mile west of Hawver School on Muddy Creek Mountain; measured with aneroid eastward to the road forks at 2440' L., then with the road down the west side of the mountain to the top of the Alderson Limestone. Measurements shown for that portion above 2440' are greater than true vertical and for that portion below 2440' the measurements are less than true vertical. Arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Bluefield Group (415'+).		
Sandstone, white, massive, Droop (in part) (top, 2640')	25	25
Shale, yellow, olive, sandy.....	90	115
Limestone, impure, shaly, fossiliferous, Reynolds (Coll. 114) (top, 2525').....	15	130
Shale, olive, sandy.....	35	165
Shale, red sandy.....	35	200
Limestone, blue, gray, impure, heavy-bedded, Glenray (top, 2440' L.).....	40	240
Shale, yellow, olive, sandy, and concealed.....	175	415

Fort Springs District: starting $\frac{1}{2}$ mile west of Hawver School on Muddy Creek Mountain; measured with aneroid eastward to the road forks at 2440' L., then with the road down the east side of the mountain to the road forks at B. M. 1788'. Measurements for that portion above the Greenbrier Series are greater than true vertical while the measurements from the top of the Greenbrier Series down are less than true vertical. Arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Bluefield Group (435' +).		
Sandstone, white, massive Droop, in part, (top, 2640' B.).....	25	25
Shale, yellow, olive, sandy.....	90	115
Limestone, impure, shaly, banded, Reynolds (Coll. 114), (top, 2525' B.).....	15	130
Shale, olive, sandy.....	35	165
Shale, red, sandy.....	35	200
Limestone, Glenray (2440' B.). Offset eastward along road to the same horizon.		
Limestone, Glenray, (Coll. 105), (top, 2355' B.). Offset eastward along the road to the same horizon....	45	245
Limestone, 5' exposed, Glenray, (Coll. 104), (top, 2140' B.)		
Shale, yellow, olive, sandy, and concealed.....	190	435
Greenbrier Series (150' +)		
Limestone, shaly.....40' } Alderson		
Limestone, massive, large } (Colls. 19 and 102)....	85	520
erminoid stems.....5 } (top, 1940' B.)		
Limestone, yellow, shaly.....40		
Limestone, blue, massive, Alderson.....	65	585
Limestone, Union, (top, B. M. 1788').....		

MEASURED SECTIONS, LEWISBURG DISTRICT.

This small, more or less rectangular-shaped district, affords very few good exposures for measuring sections. The surface rocks include the basal part of the Mauch Chunk, the Greenbrier, the Macerady, and the upper part of the Pocono Series.

Richlands Section—Northwest.

Lewisburg District; section $\frac{1}{4}$ mile northwest of Richlands; starting at the top of a knoll and measured descending southward to stream; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series (55' +)		
Shale, green, yellow and sandy.....	40	40
Sandstone, Edray, brown, cross-bedded.....	15	55
Greenbrier Series (95' +)		
Limestone, Alderson, cross-bedded and siliceous at top, blue, more pure, massive near center, shaly at base.....	65	120
Shale, Greenville, dark to yellow, fissile, fossiliferous	15	135

Lewisburg District; starting on Miller Ridge 2 miles north of Richlands and measured along the road southward; arranged in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Bluefield Group (540' +)		
Sandstone, Droop, white, massive, caps Miller Ridge		
—exposed (base, 2755' B.).....	20	20
Shale, red, brown, and concealed.....	270	290
Limestone, lenticular, shaly.....20'	60	350
Shale, brown, some reds.....30		
Limestone, fossiliferous.....10		
Shale, yellow, brown, sandy.....	150	500
Sandstone, Edray, gray, brown, cross-bedded.....	15	515
Shale, Lillydale, dark to green, fissile, concretionary	25	540
Greenbrier Series		
Limestone, Alderson, shaly.....		

MEASURED SECTIONS, FRANKFORD DISTRICT.

Frankford District borders on Lewisburg District, being more or less centered on the town of Frankford. The surface rocks range in age from the lower Mauch Chunk to the Upper Devonian.

Savannah School Section.

Frankford District; beginning at road forks on Carroll Hill and traversing southeast along county road to forks ½ mile northwest of Savannah School; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Bluefield Group (85' +)		
Concealed	10	10
Sandstone, Webster Springs, brown, partly concealed	10	20
Shale, Lillydale?, olive-green, sandy, micaceous, fissile	65	85
Greenbrier Series—Alderson Member (111')		
Limestone, gray, siliceous, Pentremites, Archimedes, (top, 2300' B.).....	10	95
Shale, dark, carbonaceous.....	5	100
Limestone, bluish-yellow, shaly, Archimedes.....	10	110
Shale, yellowish-blue, calcareous, sandy, streak of red shale.....	15	125
Limestone, bluish-gray, massive, hard, fossiliferous..	10	135
Limestone, bluish-green, weathers yellow, fenestel- loids, Archimedes, Pentremites, Composita, Spiri- fers	10	145
Shale, greenish-blue, weathers yellow, calcareous, fenestelloids, crinoid stems, Composita, Spirifer (base, 2240' B.).....	25	170
Shale, green, yellow, sandy.....	15	185
Shale, black.....	2	187
Limestone, massive, Archimedes, Pentremites.....	6	193
Limestone, yellowish-gray, chalky, plants (Coll. 109)	3	196

Fort Springs District; starting $\frac{1}{2}$ mile west of Hawver School on Muddy Creek Mountain; measured with aneroid eastward to the road forks at 2440' L., then with the road down the east side of the mountain to the road forks at B. M. 1788'. Measurements for that portion above the Greenbrier Series are greater than true vertical while the measurements from the top of the Greenbrier Series down are less than true vertical. Arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Bluefield Group (435' +).		
Sandstone, white, massive Droop, in part, (top, 2640' B.).....	25	25
Shale, yellow, olive, sandy.....	90	115
Limestone, impure, shaly, banded, Reynolds (Coll. 114), (top, 2525' B.).....	15	130
Shale, olive, sandy.....	35	165
Shale, red, sandy.....	35	200
Limestone, Glenray (2440' B.). Offset eastward along road to the same horizon.		
Limestone, Glenray, (Coll. 105), (top, 2355' B.). Offset eastward along the road to the same horizon....	45	245
Limestone, 5' exposed, Glenray, (Coll. 104), (top, 2140' B.)		
Shale, yellow, olive, sandy, and concealed.....	190	435
Greenbrier Series (150' +)		
Limestone, shaly.....40'	Alderson (Colls. 19 and 102).... (top, 1940' B.)	85
Limestone, massive, large erinoid stems.....5		
Limestone, yellow, shaly.....40		
Limestone, blue, massive, Alderson.....	65	585
Limestone, Union, (top, B. M. 1788').....

MEASURED SECTIONS, LEWISBURG DISTRICT.

This small, more or less rectangular-shaped district, affords very few good exposures for measuring sections. The surface rocks include the basal part of the Mauch Chunk, the Greenbrier, the Maerady, and the upper part of the Pocono Series.

Richlands Section—Northwest.

Lewisburg District; section $\frac{1}{4}$ mile northwest of Richlands; starting at the top of a knoll and measured descending southward to stream; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series (55' +)		
Shale, green, yellow and sandy.....	40	40
Sandstone, Edray, brown, cross-bedded.....	15	55
Greenbrier Series (95' +)		
Limestone, Alderson, cross-bedded and siliceous at top, blue, more pure, massive near center, shaly at base.....	65	120
Shale, Greenville, dark to yellow, fissile, fossiliferous	15	135

with the Pocono and the basal part of the Greenbrier Limestone, make up the list of outcropping rocks. There are very few exposures in the area suitable for measuring sections.

Caldwell Section.

White Sulphur District; beginning just east of the junction of Monroe Run and Howard Creek and traverse east along the C. & O. Railroad tracks; arrangement in descending stratigraphic order.

	Thickness, Feet.	Total. Feet.	
Greenbrier Series (42'+)			
Limestone, Hillsdale, blue, hard, black chert nodules along and across the bedding (Coll. 20).....	15	15	
Limestone, light-blue, laminated, weathers yellow, brachiopods and bryozoa (Coll, 21).....	15	30	
Shale, yellow, fissile, limy.....	8	38	
Limestone, yellow, weathered.....	4	42	
Maccrady Series (250')			
Red and buff shales and sandstones (estimated).....	250	292	
Pocono Series (600')			
Sandstone, buff, shaly.....	4	296	
Shale, yellow to olive, sandy, pyramidal joints.....	10	306	
Coal blossom and black shale.....	0.5	306.5	
Shale, dark	8	314.5	
Sandstone, brown, lentleular (0.3.5').....	3.5	318	
Shale, gray, sandy.....	12	330	
Shale, dark, carbonaceous (0.2').....	2	332	
Shale, nodular, concretionary, (mud-flat conditions)..	5	337	
Shale, gray.....	2	339	
Sandstone, gray, massive, micaceous, makes cliff..... 50'	} Broad Ford	175	514
Shale, grayish-brown, very sandy.... 25			
Sandstone, grayish-brown, conglomeratic, shaly, clay galls, pyrite concretions (Colls, 3, 23, 28, 29)..100			
Concealed, incompetent beds, (Sunbury Shale?).....	100		614
Sandstone, olive-brown, flaggy (Coll. 31).....	25		639
Sandstone, olive-brown, more massive but somewhat flaggy	35		674
Concealed, with incompetent beds.....	75		749

Spring Creek Section—South.

Frankford District; measured descending the hill south of Spring Creek, to the mouth of Spring Creek; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Greenbrier Series		
Limestone, cherty, Hillsdale (base, 2060' B.).....
Maccrady Series (125')		
Shale, olive-brown, sandy.....	10	10
Shale, red.....	115	125
Pocono Series (85'+)		
Sandstone, Broad Ford.....	85	210

Spring Creek Section—North.

Frankford District; measured ascending the hill north of Spring Creek and rearranged in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Greenbrier Series		
Limestone, cherty, Hillsdale (base, 2020' B.).....
Maccrady Series (105')		
Shale, red.....	105	105
Pocono Series (60'+)		
Sandstone, gray to brown.....	5'	} Broad Ford 60
Shale, olive-green, sandy, fissile.....	5	
Sandstone, grayish-brown to green, cross-bedded, blocky.....	50	
		165

Unus Section.

Frankford District; measured along the county road ½ mile northwest of Unus P. O.; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Blucfield Group (100')		
Shale, dark, weathers brown....90' } Lillydale	100	100
Shale, black.....10' } ('Top, 2370' B.)		
Greenbrier Series—Alderson Member (150')		
Limestone, yellowish-blue, siliceous, impure Archimedes (top, 2270' B.).....	35	135
Shale, yellow, green, red, sandy, calcareous.....	25	160
Limestone, blue, shaly, weathers yellow, abundant fossils	90	250
Limestone, Union, blue, massive (B. M. 2120').....

MEASURED SECTIONS, WHITE SULPHUR DISTRICT.

White Sulphur is a large district in the southeast corner of the county. Every major division of the Devonian rocks in

trix zone).....	20	2802
Concealed (computed).....	538	3340

Portage Series (602'+)

Shale and gray 4" flagstones (Coll. 45).....	4	3344
Concealed (computed).....	338	3682
Shale, dark, fissile, weathers brown, gray, hard flags (Colls. 46, 47).....	10	3692
Concealed	50	3742
Shale, yellow, brown, fissile, (Coll. 48, near middle)	200+	3942

Eckle School Section.

White Sulphur District; measured along the north side of the road, traversing south from Eckle School; arranged in descending stratigraphic order.

	Thickness. Feet.	Total. Feet
Marcellus Series (in part) (40')		
Shale, black, crumpled (Coll. 135).....	40+	40
Oriskany Series (88')		
Sandstone, grayish-blue, hard, calcareous. <i>Orbiculoidea roederi</i> , <i>Rhipidomella musculosa</i> , <i>Hippari- onyx proximus</i> , <i>Anoplia nucleata</i> , <i>Camarotoechia oriskania</i> , <i>Spirifer cumberlandiae</i> , <i>Spirifer murchisoni</i> , <i>Anoplothea dichotoma</i> , <i>Platyceras gebhardi</i> (Coll. 125).....	8'	
Chert, blue, black to gray, tough, minute joints, hackly.....	65	
Sandstone, Ridgeley, grayish-brown to white, frag- ments of "wheat grain" conglomerate, <i>Spirifer arcuosus</i> , <i>Spirifer murchisoni</i> , <i>Rhipidomella mus- culosa</i> (Coll. 152).....	15	128

Heldberg Series (90'+)

Limestone, gray to blue, specks of limonite, weathers sandy, hard, cross-bedded, <i>Schuchertella wool- worthana</i> , <i>Rensselaria subglobosa</i> var. <i>avus</i> , <i>Rensselaria</i> sp., <i>Spirifer concinus</i> (Coll. 123).....	50'	
Limestone, bluish-gray, limonite specks, knobby, <i>Rhipidomella ob- lata</i> , <i>Dalmanites pleuroptyx</i> (Coll. 124)	20	
Limestone, dark-blue, knobby, irreg- ular bedded bands of hard black chert, <i>Favosites conicus</i> , <i>Spirifer</i>		

Becraft
(Coll. 151).. 90 218

streaked with white quartz pebbles	4'		
Sandstone, grayish-brown, fine-grained, massive.....	15		
Sandstone, gray, quartz pebbles....	2		
Sandstone, gray, fine-grained.....	6		
Conglomerate, loosely cemented, various sizes of white quartz pebbles	35		
Shale, brown, arenaceous.....	6	Berea	
Conglomerate	5	Conglomerate...	143 892
Shale, gray to buff, with thin sandstone flags.....	9		
Sandstone, light-gray, many pebbles	4		
Sandstone, gray, weathers brown..	11		
Shale, gray, brown, arenaceous....	20		
Sandstone, 12" to 18" flags, with quartz pebbles at base.....	15		
Shale, brown, fissile, sandy.....	10		
Conglomerate, 0' 8" to 1'.....	1		

Chemung Series (2448')

Shale, brown, fissile (Colls. 4, 30, 24).....	20	912
Sandstone, gray, massive.....	8'	
Sandstone, gray, shaly.....	12	
Sandstone, dark-gray, massive.....	15	
Sandstone, conglomerate, gray, with quartz pebbles.....	2	
Sandstone, gray, shaly.....	10	959
Concealed	200	1159
Sandstone, yellowish-brown, flaggy and shaly. (Coll. 25 near base).....	68	1227
Sandstone, hard, massive (Coll. 26).....	20	1247
Sandstone, grayish-brown, micaceous (Coll. 32).....	24	1271
Sandstone, and sandy shale, gray, olive, and brown (Coll. 33).....	10	1281
Sandstone, olive, and somewhat shaly (Coll. 34).....	10	1291
Sandstone, gray, green, brown, shaly, exfoliated, weathering (Coll. 35 at base).....	80	1371
Sandstone, gray, olive and brown, olive, sandy shale (Coll. 36 at base).....	100	1471
Concealed (estimated).....	400	1871
Shale and sandstone, gray and green flags and brown sandy shale; exposed along Midland Trail at The Pines (Coll. 37).....	100	1971
Concealed (computed).....	511	2482
Shale, olive and brown, sandy, sandstone flags and green, olive and brown shales.....	35	2517
Sandstone, grayish-green and brown sandy shale (Coll. 39).....	15	2532
Sandstone, gray, tough, flaggy and shaly (Colls. 40, 41)	50	2582

Chert, tough, blue-gray haekly.....	45'	Hunters- ville Chert	70	570
Sandstone, fine, green, weathers brown (sample 1M).....	5			
Chert, dark-gray.....	20			
Sandstone, medium-grained, light- brown, porons (sample 2M).....	5'	Ridgeley Sandstone ..	15	585
Sandstone, medium-grained, light- brown, porous (sample 3M).....	5			
Sandstone, medium-grained, light- brown, porous (sample 4M).....	5			

Helderberg Series (300')

Limestone and chert, Becraft.....			60	645
Sandstone, hard, fine-grained, white to light-gray, almost a quartzite (sample 5M).....	5'	New Scot- land Member —Healing Springs Sandstone ..	25	670
Sandstone, hard, fine-grained, white to light-gray, some porous spaces (sample 6M).....	5			
Sandstone, quartzite, weathered, limonite stained and porous from leaching (sample 7M).....	5			
Sandstone, and concealed (not sam- pled)	10			
Concealed and shaly limestone (Coeymans if present).....	35'			
Limestone, sandy to shaly.....	20			
Limestone, gray, platy, calcite streaks	20			
Limestone, blue-gray, massive, cal- cite streaks	40			
Limestone, blue-gray.....	20			
Concealed and gray limestone.....	65			
Sandstone, Clifton Forge, fine- grained, hard, porous and limonite stained from weathering, strongly cemented by silica, (sample 8M)....	5	Keyser Lime- stone Member	215	885
Sandstone, Clifton Forge, fine- grained, hard, porous and limonite stained from weathering (sample 9M)	5			
Sandstone, Clifton Forge, fine- grained, hard, some parts porous and limonite stained from weath- ering (sample 10M).....	5			

Salina and Niagara Series (400')

Limestone, light-gray, thin-bedded, cut by an intri- cate network of calcite veins.....	130	1015
Limestone, bluish-gray, thin, platy, cut by calcite veins	90	1105

Bobs Ridge Section.

White Sulphur District; measured on the east side of the gap where Howard Creek cuts through Bobs Ridge; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Oriskany Series (70'+)		
Chert, Huntersville.....	50+	50
Sandstone, white, coarse, iron-stained fossil pits.....10'	} Ridgeley 20	70
Concealed10		
Helderberg Series (130'+)		
Limestone, gray, crystalline, sandy, limonite specks, few fossils.....25'	} Becraft Member 90	160
Limestone, gray, crystalline, crinoid stems, massive, sandy.....15		
Limestone, gray, to black, cherty, <i>Streptelasma strictum</i> , <i>Schuchertella woolworthana</i>50		
Sandstone, Healing Springs, white to brown, massive, quartzite, 8 samples taken for mineralogical study (in 5' sections)	} New Scotland Member 40	200

MEASURED SECTIONS, ANTHONY CREEK DISTRICT.

This large district, in the northeast part of the county, contains the oldest rocks outcropping in the territory covered by this report. The outcropping rocks range from the Greenbrier Limestone of the Mississippian down to the Red Medina of the Silurian. In spite of the size of the district and the great thickness of rocks exposed, there are very few exposures at which it is worth while measuring a section. In most of the area the rocks have been so folded and mashed that a true thickness can not be obtained.

In the following two sections, sample numbers marked 1M, 2M, etc., indicate that specimens were collected for mineralogical examination:

Alvon Section—West Side.

Anthony Creek District; measured along the north side of Anthony Creek; traversing southeastward and starting at a point 0.7 mile northwest of Alvon; corrected for dip and arranged in descending stratigraphic order.

Thickness. Feet.	Total. Feet.
---------------------	-----------------

Concealed	25	1285
Clinton Series (in part; 60')		
Sandstone, fine-grained, white, quartzitic (sample 11M).....	5	
Sandstone, fine-grained, hard, white to brown (sample 12M).....	5	
Sandstone, fine-grained, hard, ilmenite stains, small cavities lined with quartz crystals (sample 13M).....	5	
Sandstone, fine-grained, hard, white, weathers brown, some parts porous (sample 14M).....	5	
Sandstone, fine-grained, hard, quartzitic, white to brown (sample 15M)	5	
Sandstone, fine-grained, hard, white to brown (sample 16M).....	5	
Sandstone, fine-grained, white, quartzitic (sample 17M).....	5	
Sandstone, fine-grained, hard, white, weathers brown, less weathered parts contain a little calcite and pyrite (sample 18M).....	5	
Sandstone, fine-grained, hard, gray, contains some calcite and pyrite (sample 19M).....	5	
Sandstone, fine-grained, hard, white to brown (sample 20M).....	5	
Sandstone, fine-grained, very hard, quartzitic, white (sample 21M).....	5	
Sandstone, fine-grained, hard.....	5	
	Keefer	
	Sandstone ..	60 1345

Burr Valley Section.

Pocahontas County, Little Levels District; measured along the road traversing southeastward, starting at a point 1.1 miles south-southeast of Burr School and 0.7 mile northeast of Burr; corrected for dip and arranged in descending stratigraphic order.

	Thickness.	Total.
	Feet.	Feet.
Oriskany Series (93'±)		
Chert, yellow, sandy.....	4	
Chert, gray to black.....	8	
Sandstone, green, fine to very fine-grained, weathers brown, sample 1M)	1	
Chert and concealed.....	60	
	Huntersville	
	Chert	73 73

For convenient reference the thickness of the exposed stratified rocks of Greenbrier County, as determined by the measured sections of this Chapter, is compiled in the following table, showing not only the thickness of the various series but also the totals for the different grand divisions, or periods, down to the lowest depths to which there are exposures or borings. A line of dots (.....) under a series indicates that it was not exposed or in some cases not examined, where the section was measured. A question mark (?) indicates that the series was present and was examined but could not be differentiated from the one overlying or the one below it. A plus mark (+) indicates that only a portion of the full series or period is included in the section. In some few cases a section shows a thickness of a series either too great or too small, owing to the dip of the strata where it was made, a reduction to true vertical measurement being impracticable in some of the sections. Sections of this type that effect the accuracy of the table have been marked with an asterisk (*), to indicate that the reader should refer to the detailed section. In all localities where the rocks dip steeply, particularly in the Devonian, all sections were reduced to true vertical measurement and so published. An explanation accompanies each section, where published in the text, detailing the conditions under which it was made:

Sandstone, medium-grained, gray to brown, abundant pyrite (sample 2M)	5		
Sandstone, "wheat grain" conglomerate, porous from leaching of calcareous material, stained light-brown from limonite (sample 3M)..	5	Ridgeley Sandstone ..	20+ 93+
Sandstone, medium-grained, porous from leaching of calcareous material, stained light-brown by limonite (sample 4M).....	5		
Sandstone, medium- to coarse-grained, white to brown, porous from leaching (sample 5M).....	5		
Loose fragments, doubtful.....			

Penbrier County and Adjacent Areas.

[illegible]

Summary of Measured Sections

NAME OF SECTION OR NUMBER OF BORING	PENNSYLVANIAN			MISSISSIPPIAN								
	POTTSVILLE			MAUCH CHUNK								
	New River	Pocahontas	Total	Bluestone	Princeton	Hinton	Blackfield	Total	Greenbrier	Macerady	Pocono	
Aeme		90+	90+	663	35	840	1202	2749	297+	30+		
Alderson							293+	293+	610	75+		
Alta							330+	330+	35+			
Alum Bum												
Alvon								521+				
Big Clear Creek Mountain.....	157+	284	441+	?	?	?	155+	155+	25+			
Blaker Mills.....							195+	195+				
Blue Sulphur Springs.....												
Bobs Ridge.....												
*Briery Knob (Pocahontas Co.).....	431+		431+	297	50	?	?	1365	210+			
Burr Valley (Pocahontas Co.).....							505+	505+	65+			
Butler Mountain.....									42+	250	600	
Caldwell	?	?	825+	49+				49+				
Charmco				230+	55	?	?	815+				
Cherry Low Place.....	?	?	315+	395	80	475	935	1885	475	80	205	2
*Cold Knob—Hinkle Well.....												
Duo	277+		277+									
Eckle School.....												
*Goddard Mountain.....	?	?	415+	300	5+			305+				
Green Sulphur Springs.....		120+	120+	435	20	?	?	2220	695	25	600	3
Hawver School—East.....							435+	435+	150+			
Hawver School—West.....							415+	415+				
Hominy Falls (Nicholas Co.).....	1057		1057+	120	140	?	?	1418	393	?	?	2
Horseshoe Bend School.....									70+	180	5+	
Julia Post-Office									551+	3+		
Kieffer					35	300+		335+				
Little Clear Creek.....	?	?	420+	?	?	?		480+				
Little Rocky Run.....	?	?	070+	?	?	?		650+				
Little Sewell Mountain—South End	?	?	369+	?	?	?		608+				
Little Sewell Mountain—West Side	?	?	355+	255	5+			260+				
No. 5 E.....	387+		387+									
No. 6.....	537+	70+	607+									
No. 7.....	480+		480+									
*No. 11.....	557+	271	828+	11+				11+				
*No. 13.....	175+	330	505+	36+				36+				
Patton									285+			
Quinwood	411+		411+									
Renick							260+	260+	405+			
Renick Valley.....							400+	400+	225+			
Richlands—Northwest							55+	55+	95+			
Richlands—Two Miles North.....							540+	540+				
*Roach Run.....				90+	25	520	280+	915+				
Russellville	?	?	733+				85+	85+	151+			
Savannah School.....												
Sims Mountain—North End.....	?	?	427+									
*Sims Station.....	?	?	370+	330	20+			350+				
Spring Creek—North.....										105	60+	
Spring Creek—South.....										125	85+	
Turniphole Mountain.....	?	?	311+									
Unus							100+	100+	150+			

*See detailed section for dip correction.

CHAPTER VI.

STRATIGRAPHY—PENNSYLVANIAN ROCKS.

INTRODUCTION.

The Pennsylvanian System of rocks forms the uppermost grand division of stratified beds in Greenbrier County, being succeeded only by certain terrace gravels and river clays, that may be of Pleistocene age. The Pennsylvanian probably once covered all of the county but any estimate of its original thickness would be conjectural, although it is likely that most of its subdivisions, as known in counties to the north and west, may have been formed in this area and later removed by erosion.

The subdivisions now remaining, and as classified in descending stratigraphic order, are as follows:

	Feet.
Kauawha Group.....	250±
New River Group.....	600 to 950
Pocahontas Group.....	0 to 340
<hr/>	
Apparent maximum.....	1,540

The various groups are composed of sandstone, sandy or fire clay shales, carbonaceous shales, and coals.

The outcrop of rocks of the Pottsville Series is confined to the northwestern fourth of the county. Figure 8 shows the outcrop of the Pottsville rocks, and Map II shows the same

Limestone, light-gray, stylolitic structure, fossiliferous, (quarry, average dip, 23° N. W.).....	25'	} Patton and Sinks Grove	75	823
Limestone, dark-gray, massive.....	50			
Limestone, Hillsdale, blue, gray, massive, nodules of irregular black chert (Coll. 71).....			80	903
Maccrady Series (75'±)			75±	978
Shale, red.....				

MEASURED SECTIONS, FALLING SPRINGS DISTRICT.

Falling Springs is the northernmost district in the county. It includes most of the drainage area of North and South Forks of Cherry River, most of the drainage area of Spring Creek, and the drainage area of several small streams on the east side of Greenbrier River north of the village of Anthony. The surface rocks range from the Kanawha Group of the Pottsville down to the middle Chemung. Sections measured in this district afford the best detailed measurements of the Greenbrier Limestone available in the county.

Little Rocky Run Section.

Falling Springs District; measured with aneroid starting at the top of the high knob (elevation, 4030' L.) north of Little Rocky Run, traversing south to Little Rocky, thence westward to South Fork of Cherry River.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River and Pocahontas Groups (670'±)		
Concealed to top of bench.....	115	115
Sandstone, Welch?.....	20	135
Concealed	10	145
Sandstone, makes cliff.....30'	} Upper Raleigh?.....	200
Concealed		
Sandstone, massive, white, with white quartz pebbles 20		
Concealed	55	200
Sandstone, coarse, conglomerate.....	460	660
	10	670
Mauch Chunk Series (650'±)		
Concealed	120	790
Sandstone, coarse.....	10	800
Concealed	30	830
Shale, red, and concealed.....	185	1015
Sandstone, brown, Princeton?.....	50	1065
Concealed	10	1075
Shale, yellow to brown.....	10	1085
	30	1115

	Feet.	Feet.	
Mauch Chunk Series—Bluestone Group (297')			
Sandstone, reddish-brown, shaly.....	5	436	
Concealed	52	488	
Concealed in flat bench.....	165	653	
Shale, yellow	5	658	
Concealed	70	728	
Mauch Chunk Series—Princeton Group (50')			
Concealed in bench but large conglomerate boulders (Princeton Sandstone).....	50	778	
Mauch Chunk Series—Hinton and Bluefield Groups (1018')			
Shale, red.....	200	978	
Sandstone, Stony Gap , red and brown, cross-bedded, makes bold cliff, shaly at top.....	40	1018	
Shale, red.....	125	1143	
Sandstone, grayish-brown, micaceous.....	25	1168	
Shale, red.....	20	1188	
Shale, partly concealed, but mostly red.....	250	1438	
Sandstone, red.....	5	1443	
Shale, red.....	105	1548	
Sandstone, reddish-brown.....	10	1558	
Shale, red.....	25	1583	
Sandstone, red.....	3	1586	
Shale, red, green, sandy.....	30	1616	
Shale, yellow, olive, to road forks (2715' B.).....	40	1656	
Shale, olive, dark.....	10	1666	
Limestone, Reynolds , very fossiliferous, impure, shale at top.....	20	1686	
Shale, dark, olive, sandy.....35'	} Lillydale	85	1771
Sbale, red.....20			
Sbale, olive, sandy.....30			
Sandstone, Edray , grayish-brown, micaceous.....	25	1796	
Greenbrier Series (210'+)			
Limestone, Alderson , massive, gray, fossiliferous.....	10	1806	
Concealed	90	1896	
Shale, Greenville , dark, carbonaceous, fossiliferous....	40	1936	
Concealed to Hills Creek (2465' B.)....	70	2006	

Butler Mountain Section.

Falling Springs District; beginning at road forks, elevation 2729', on north end of Butler Mountain, traversing southeast along road to $\frac{1}{2}$ mile east of Rapp School; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Bluefield Group (505'+)		
Sandstone, Droop , gray, medium-grained.....	50	50
Shale, red, brown, and concealed.....	215	265
Limestone, Reynolds , shaly.....	45	310
Shale and concealed, mostly reds.....	50	360
Limestone, light-gray, laminated.....	5	365
	90	455

	Feet.	Feet.
Shale, red, yellow, and coneealed, Lillydale.....	40	505
Greenbrier Series—Alderson Member		
Shale, yellow calcereous (top, 2200').....	5	510
Limestone, blue, massive, Archimedes, crinoid stems	5	515
Limestone, shaly, grayish-blue, weathers yellow, fissile, encorals, <i>Athyris</i> , bryozoa, <i>Productus</i> (Coll. 113).....	10	525
Limestone, yellowish-gray, argillaceous, abundant <i>Pentremites</i> , crinoid stems large and small, horn corals, <i>Athyris</i> , <i>Spirifer pellaensis</i> , fenestelloids and other bryozoa (Coll. 112).....	15	540
Limestone, gray-blue, shaly, fissile, abundant <i>Productus</i> , <i>Orbiculoidea</i> , and <i>Ambocoelia</i> (Coll. 111).....	15	555
Limestone, blue, oolitic, crinoid stems, blastoid plates, <i>Pentremites</i> , horn corals.....	5	560
Limestone, yellowish-gray, some red, weathers to a yellow clay, plants, fish plate (Coll. 110).....	10	570
Greenbrier Series—Union Member		
Limestone, bluish-gray, oolitic, stylolitic, massive, pure, (top, 2110' B.).....

In the following section, no division is made between the Hinton and Bluefield Groups of the Mauch Chunk Series. It is probable, however, that the grayish-brown sandstone at 2940' B. is the Stony Gap and the base of the Hinton Group:

Cherry Low Place Section.

Falling Springs District; starting from the top of a small knob $2\frac{1}{4}$ miles north of Leonard and descending the southeast side of the mountain to Panther Camp Creek; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Bluestone Group (230' +)		
Knob capped by fine-grained sandstone (top, 3520' B.)	20	20
Concealed	35	55
Shale, green, brown, sandy, crumbly.....	10	65
Shale, dark, carbonaceous, ostracods, pelecypods (top, 3450' B.).....	5	70
Concealed	50	120
Shale, grayish-brown (top, 3400' B.).....	5	125
Concealed	25	150
Shale, dark, carbonaceous.....	30	180
Concealed	40	220

Greenbrier Series (225'+)

Limestone, olive-green. shaly (top, 2675' B.).....	3'	Alderson	95	495
Shale, olive, weathered.....	8			
Limestone, gray, fossiliferous.....	5			
Shale, gray, weathers yellow.....	59			
Limestone, blue, hard (top 2600' B.).....	10			
Limestone, red, shaly.....	5	Union	130	625
Limestone, gray, weathers white, stylolite	5			
Limestone, gray, oolite (top, 2565' B.).....	10'			
Limestone, light oolite, stylolite.....	10			
Limestone, blue, gray, hard, brittle, sparse black chert.....	110			

Julia P. O. Section.

Falling Springs District; starting 1.5 miles northwest of Julia P. O. on the top of a high knob and continuing southeastward down the highway toward Julia; arrangement in descending stratigraphic order.

		Thickness. Feet.	Total. Feet.
Greenbrier Series (551'+)			
Limestone, and concealed (top, 2565')	171'	Union	200
Limestone, blue, massive (top, 2394' L.).....	29		
Limestone, gray to yellow, shaly (top, 2365' B.).....	5'	Pickaway	130
Limestone, light-gray.....	10		
Limestone, yellow (top, 2350' B.)....	20		
Limestone, dark-gray, weathers yellow, argillaceous.....	10		
Limestone, bluish-gray, hard, massive	30		
Limestone, blue-gray, shaly (top, 2290' B.).....	25		
Limestone, yellow, argillaceous, weathers white.....	2		
Limestone, light-gray, with clay seams, light-blue at base.....	28		
Limestone, red, shaly (top, 2235' B.)	2'	Taggard	25
Limestone, light, weathers yellow....	3		
Limestone, red, shaly.....	5		
Limestone, light-gray and yellow.....	5		
Limestone, yellowish-red.....	3		

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Princeton Conglomerate (55')		
Sandstone, greenish-brown, flaggy (top, 3285' B.).....	5'	Princeton 55 290
Sandstone, brown, medium-grained, massive	20	
Conglomerate, brown, porous, rotten, plants	30	

Mauch Chunk Series—Hinton and Bluefield Groups (530'+)

Coneaealed (top, 3230' B.).....	5	295
Shale, yellow, sandy.....	10	305
Coneaealed	200	505
Limestone blocks, Avis? (3015' B.).....	505
Coneaealed	75	580
Sandstone, grayish-brown (top, 2940' B.).....	15	595
Shale, red, to 2850' B.....	75	670
Coneaealed and red shale to house at Panther Camp Creek (2710' B., 2746' L.).....	140	810

Renicks Valley Section.

Falling Springs District; starting at the junction of Brushy Mountain and Droop Mountain, 400 feet northwest of the Pocahontas-Greenbrier County line; measured descending southeastward along the highway toward Renicks Valley; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Bluefield Group (400'+)		
Sandstone, Droop, white (base, 3035' B.).....	40+	40
Shale, Talcott and Ada, olive-green, yellow, sandy (base, 2980' B.).....	55	95
Concealed	40	135
Shale, yellow, much weathered (2910' B.).....	30	165
Coneaealed	15	180
Shale, green to olive.....	5	185
Shale, red (2860' B.).....	30	215
Coneaealed	10	225
Shale, yellow, weathered.....	5	230
Shale, red.....	5	235
Coneaealed	15	250
Sandstone, Edray or Webster Springs, yellowish- brown, flaggy at top, massive near middle, flaggy and shaly at base (2785' B.).....	40	290
Shale, Lillydale, green to olive-brown, sandy and flaggy at top, fissile with coneretions in lower part, exfoliate type of weathering (2735' B.).....	50	340
Shale, dark.....	45	385

	Feet.	Feet.
Shale, yellowish-brown, fissile.....55'	Lillydale	65 260
Shale, dark-gray, sandy, <i>Orthis</i> , peleypods, (Coll. 91).....10		
Greenbrier Series (405'+)		
Limestone, yellowish-gray, weathers yellow, cup corals, erinoid stems, brachiopods, <i>Archimedes</i> (Coll. 90)10'	Alderson	85 345
Limestone, red, sandy, shaly.....25		
Limestone, yellowish-gray, shaly at top, more solid at base, fossilifer- ous (Coll. 92).....50		
Limestone, blue, massive, oolitic. stylolitic40'	Union	185 530
Limestone, yellow, clayey.....10		
Limestone, yellowish-blue, shaly.....5		
Limestone, blue, massive.....25		
Shale, yellow on weathered exposure..10		
Limestone, blue, hard, abundant erinoid stems, <i>Pterotoechinus</i> , brach- iopods, small stylolitic structure....15		
Concealed at upper end of village.....50		
Limestone, dark, blue, massive.....30	Pickaway	135 665
Limestone, grayish-yellow, shaly.....15'		
Limestone, gray, blue, massive, to post-office85		
Concealed to railroad.....35		

MEASURED SECTIONS, BLUE SULPHUR DISTRICT.

Blue Sulphur District is located in the southwest corner of the county and includes most of the drainage area of Muddy Creek. The outcropping rocks range from the Hinton Group of the Mauch Chunk down to the upper part of the Pocono.

Blue Sulphur Springs Section.

Blue Sulphur District; starting at a point 2 miles south of Blue Sulphur Springs along the Alderson highway; measured with aneroid southward approximately 1 mile, there being a gentle dip to the north-west; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Bluefield Group (195'+)		
Shale, greenish-brown, fissile.....	10	10
Limestone, Reynolds, blue to yellowish-blue, fossil- iferous (top, 1865' B.).....	40	50
	10	60

		Feet.	Feet.
Limestone, light-blue, with streaks of pink and yellow, (base, 2205' B.)	5'		
Limestone, light-gray, partly oolitic	10		
Limestone, weathers yellow	5		
Limestone, grayish-blue, massive, oolitic at top	30	Patton	125
Limestone, light-blue, dense	30		480
Limestone, yellowish-white	10		
Limestone, light-blue, massive, to road forks (base, 2071' L., 2085' B.)	35		
Limestone, light-gray	20'		
Limestone, dark, abundant nodules of black chert	2	Sinks Grove	42
Limestone, light-blue, weathers yellow, pure, white silica geodes in base	20		522
Shale, bluish-green, with red coral colonies	1'		
Limestone, bluish-black, abundant black chert	20	Hillsdale	29
Limestone, yellow	8		551
Maccrady Series (3'+)			
Shale, blue, red, old soil			3
Shale, red			554

Renick Station.

Falling Spring District; beginning on a knoll on Falling Spring Mountain, one mile southwest of Modoc P. O. and descending east to cut in road summit, and thence southeast along highway to Renick village, thence along highway to Renick P. O. and Station; arrangements in descending stratigraphic order.

	Thickness. Feet.	Total Feet.
Mauch Chunk Series—Bluefield Group (260'+)		
Knoll capped by olive-green sandy shale (2650' B.)		
Concealed	35	35
Limestone, blue, hard, exposed	2	37
Concealed	48	85
Shale, red, sandy, summit of road cut, (Coll. 87 at base)	15	100
Limestone, Reynolds, gray, clayey (Coll. 88)	15	115
Shale, yellow, olive, sandy	20	135
Shale, red, sandy	10	145
Sandstone, Webster Springs, grayish-brown, cross-bedded	10	155
Limestone, Glenray, gray, hard, cross-bedded, very fossiliferous, crinoid, blastoid, fenestelloid, Spirifer, Productus, Archimedes, Pterotocrinus, cup coral, coprolite, fish teeth (Coll. 89)	15	170

	Feet.	Feet.
Limestone, Glenray, blue, hard, siliceous, (top, 1800' B.).....	15	90
Shale, yellowish-brown, sandy.....	10	100
Shale, red.....	10	110
Sandstone to shale, brown.....	25	135
Shales, sandy, brown, lamulated, almost sandstone....	60	195

Alum Run Section.

Blue Sulphur District; starting 1.5 miles southwest of Brushy Flat School; measured with aneroid along the highway southward, down Alum Run; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Church Series—Bluefield Group (350' +)		
Sandstone, Droop, grayish-brown, medium-grained, very hard, cemented with silica, iron-stained, streaks of coal and carbonized plants, ripples and cross-bedding, makes cliff (top, 2110' B.).....	75	75
Concealed	270	345
Shale, black, carbonaceous, pelecypods.....	5	350

Greenbrier Series (35' +)

Limestone, yellow, shaly at top; Archimedes, brachiopods, eup corals, etc. (top, 1765' B.).....	5'	Alderson	35	385
Limestone, grayish-blue, shaly weathers yellow; brachiopods, bryozoan, etc.....	30			
Limestone, Union, gray, massive, white, weathers light-blue; Archimedes, Pentremites, cup corals....			

Blaker Mills Section.

Blue Sulphur District; starting at the top of a small hill just north of Blaker Mills; measured with aneroid southeast to Mill Creek; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Bluefield Group (155' +)		
Shale, yellow, sandy at top (1780' B.).....	20	20
Limestone, Glenray, impure, fossiliferous.....	10	30
Shale, yellow, sandy, concretionary.....	100	130
Sandstone, gray, brown to reddish.....	5	135
Concealed (base, 1625' B.).....	20	155
Greenbrier Series (25' +)		
Limestone, Alderson, to creek.....	25	180

The following section, measured by D. B. Reger, is located just outside of Greenbrier County and in the edge of Summers County. It is reprinted from pages 256 to 258 of the Mercer, Monroe, and Summers County⁶ report:

Summers County; Talcott District; starting at the top of Keeney Knob on Keeney Mountain and traversing southeastward to Mt. Zion Church and thence southward down Possum Hollow to its junction with Greenbrier River 1.5 miles west of Alderson; dip northwest about 100 feet per mile; arrangement in descending stratigraphic order; Nos. 1-70, inclusive, were measured with aneroid but the apparent thicknesses of these members were increased approximately $12\frac{1}{2}$ per cent., or a total of 280 feet, to show a true vertical section; Nos. 71 to 79, inclusive, were measured by separate determination or by estimate.

	Thickness. Feet.	Total. Feet.
Pottsville Series—Pocahontas Group (90' +)		
1. Sandstone, Lower Pocahontas, gray, forms top of Keeney Knob (3925' B.).....	22	22
2. Conealed in slope, with yellow sandy soil.....	45	67
3. Sandstone, buff.....	23	90
Mauch Chunk Series—Bluestone Group (663')		
4. Shale, yellow, sandy, with plant fossils (3845' B.)	22	112
5. Sandstone, greenish-gray, flaggy (3840' B.).....	6	118
6. Shale, red.....	73	191
7. Sandstone, brown, shaly, micaceous.....	17	208
8. Shale, red.....	62	270
9. Conealed	56	326
10. Sandstone, massive, coarse, buff, micaceous, cliff rock (3610' B.).....	50	376
11. Shale, red, largely conealed, to highest road fork (2410' B.).....	225	601
12. Conealed	56	657
13. Shale, variegated.....	28	685
14. Sandstone, shaly.....	11	696
15. Conealed, with dark shale, (3275' B.).....	57	753
Mauch Chunk Series—Princeton Conglomerate (35')		
16. Sandstone, Princeton, gray, massive, coarse, pebbly (3245' B.).....	35	788
Mauch Chunk Series—Hinton Group (849')		
17. Conealed	90	878
18. Shale, sandy.....	17	895
19. Shale, red.....	51	946
20. Sandstone, greenish-brown, shaly (3090' B.).....	17	963
21. Shale, red, and variegated.....	80	1043
22. Sandstone, Avis, green, flaggy, cliff rock (2995' B.).....	30	1073
23. Shale, red.....	16	1089
24. Shale, Upper Avis, greenish-yellow, limy, with marine fossils, pelecypods.....	11	1100
25. Limestone, Avis, steel-gray, shaly at middle, with numerous marine fossils, pelecypods, brachiopods, gastropods, crinoids, and bryozoa (2915' B.).....	28	1128
26. Shale, Lower Avis, yellow, calcareous.....	33	1161
27. Shale, red.....	170	1331

	etc.) and bryozoa; (estimated).....	Feet. 60	Feet. 2649
76.	Concealed, horizon of Reynolds Limestone (estimated)	15	2664
77.	Limestone, Glenray, shaly and sandy (estimated)	75	2739
78.	Shale, Lillydale, ("Pencil Cavo"), black and green, fissile (estimated).....	100	2839
Greenbrier Series (30'+)			
79.	Limestone, Alderson, hard, yellowish-blue, with marine fossils, bryozoa (Archimedes) etc., visible above Greenbrier River at mouth of Possum Hollow.....	30	2869

The following section and the combined well record, compiled by D. B. Reger, was published in the report cited in the foot-note:

Green Sulphur Springs Section.

Summers County, Green Sulphur District; starting at the top of Big Swell Mountain 1 mile southeast of Mountain View School and thence extending northwestward to this school and thence northeastward to the mouth of Mill Creek 0.4 mile south of Green Sulphur Springs; gentle northwest dip; measured with aneroid and arranged in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Pottsville Series—Pocahontas Group (120'+)		
Sandstones and shales from top of Big Swell Mountain, not examined but stratigraphic thickness estimated after making deduction for southeastward rise	100	100
Sandstone, buff, massive, pebbly, makes top of ridge near Mountain View School (2860' B.).....	20	120
Mauch Chunk Series—Bluestone Group (435')		
Shale, red and variegated.....	50	170
Sandstone, buff, coarse, micaceous, cliff rock (2795' B.).....	15	185
Shale, red.....	75	260
Sandstone, green.....	30	290
Shale, red.....	65	355
Sandstone, green, flaggy, much weathered, makes sharp ridge (2585' B.).....	40	395
Shale, sandy (2550' B.), with marine fossils, pelecypods	35	430
Shale, sandy.....	55	485
Shale, dark.....	35	520
Fire clay shale, streak, (2460' B.).....	520
Shale, green.....	35	555

	Feet.	Feet.
30. Sandstone, reddish-brown, with massive layers interbedded with shale (2645' B.).....	50	1464
31. Shale, variegated.....	23	1487
32. Sandstone, shaly, reddish-brown (2595' B.).....	34	1521
33. Shale, red, partly concealed.....	39	1560
34. Sandstone, massive, reddish-brown (2550' B.).....	11	1571
35. Shale, red and variegated.....	49	1620
36. Sandstone, <i>Stony Gap</i> , massive at base, cliff rock (2490' B.).....	17	1637

Mauch Chunk Series—Bluefield Group (1202')

37. Shale, calcareous, with restricted fauna; ostracods and annolids (<i>Spirorbis</i>).....	6	1643
38. Sandstone, shaly.....	6	1649
39. Shale, variegated.....	5	1654
40. Sandstone, shaly.....	11	1665
41. Shale, variegated.....	28	1693
42. Sandstone, shaly.....	6	1699
43. Shale, green, sandy, partly concealed.....	11	1710
44. Sandstone, greenish, massive, cliff rock (2415' B.)	11	1721
45. Shale, variegated and lmy.....	37	1758
46. Sandstone, greenish-brown, shaly at top (2372' B.)	11	1769
47. Shale, variegated.....	2	1771
48. Limestone, yellow, shaly (2365' B.).....	6	1777
49. Shale, variegated.....	9	1786
50. Sandstone, brown, shaly.....	2	1788
51. Shale, red.....	4	1792
52. Limestone, yellow, earthy, and brecciated (2350' B.).....	2	1794
53. Shale, red.....	11	1805
54. Sandstone, reddish-brown, shaly (2330' B.).....	11	1816
55. Shale, red and green.....	23	1839
56. Sandstone, shaly.....	6	1845
57. Shale, yellowish-green, calcareous, with marine fossils (2290' B.), numerous pelecypods.....	17	1862
58. Shale, red and variegated.....	39	1901
59. Sandstone, reddish-brown, shaly (2250' B.).....	6	1907
60. Shale, red.....	11	1918
61. Shale, green.....	6	1924
62. Shale, red.....	6	1930
63. Sandstone, reddish-brown, shaly (2225' B.).....	6	1936
64. Shale, red, streaked with green.....	88	2024
65. Shale, sandy.....	40	2064
66. Shale, red and variegated, sandy.....	96	2160
67. Sandstone, shaly (2005' B.), outcrops at Mt. Union Church.....	22	2182
68. Shale, red and variegated, partly concealed.....	165	2347
69. Sandstone, flaggy, cliff rock (1845' B.).....	17	2364
70. Shale, red and variegated, partly concealed.....	135	2499
71. Limestone boulders (1725' B.).....	2499
72. Concealed (estimated).....	50	2549
73. Sandstone, <i>Droop</i> , shaly, cliff rock.....	25	2574
74. Shale, green, sandy.....	15	2589

	Thickness.	Total.
Sandstone, shaly (1605' B.).....	10	1375
Shale, red.....	53	1428
Shale, green, to well mouth (1545' L.).....	7	1435
Section continued by record of James H. Gwinn No. 1 Well (No. 7 on Map II):		
Gravel	20	1455
Red rock.....	215	1670
Slate, white.....	15	1685
Lime, white.....	10	1695
Slate, white.....	20	1715
Red rock.....	20	1735
Slate	30	1765
Red rock.....	20	1785
Slate, white.....	15	1800
Red rock.....	25	1825
Lime, black.....	20	1845
Red rock.....	30	1875
Slate, white.....	20	1895
Red rock.....	40	1935
Lime, black.....	10	1945
Red rock.....	45	1990
Slate, black.....	15	2005
Sand, Maxton (Droop Sandstone?).....	50	2055
Slate, white.....	100	2155
Lime, Reynolds, black.....	15	2170
Slate, white.....	3	2173
Lime, black.....	17	2190
Slate, white.....	10	2200
Little Lime (Glenray Limestone), black.....	65	2265
Pencil Cave (Lillydale Shale).....	75	2340
Greenbrier Series (695')		
Big Lime (Greenbrier).....	695	3035
Maccrady Series (25')		
Slate, white.....	25	3060
Pocono Series (600')		
Sand, (Big Injun, Logan, Burgoon).....	75	3135
Slate and shells.....	325	3460
Sand, (Weir), gray.....	15	3475
Shells, hard lime (gas show at 2100').....	60	3535
Slate	15	3550
Shells, hard lime.....	85	3635
Sand, (Berea), white.....	25	3660
Chemung Series (775'+)		
Shells, flinty.....	65	3725
Slate, black.....	50	3775
Lime and shells, flint (gas shows at 2365' and 2760')..	480	4255
Shale, black (gas show at 2825').....	5	4260
Lime and flint.....	175	4435

"A strong stream of salt sulphur water now flows from the 14-inch easing which protrudes from the mouth of this well there being numerous bubbles

	Feet.	Feet.
Mauch Chunk Series—Princeton Conglomerate (20')		
Sandstone, Princeton, green, massive, with gray streaks, somewhat soft and weathered (2405' B.)....	20	575
Mauch Chunk Series—Hinton and Bluefield Groups (1765')		
Shale, red and variegated.....	25	600
Sandstone, green, massive, with streaks of shale.....	15	615
Shale, dark-green, (2342' B.); contains numerous marine fossils, pelecypods.....	23	638
Limestone, yellowish-green, shaly.....	2	640
Shale, green.....	15	655
Concealed, and red shale.....	30	685
Shale, green, with sandstone.....	20	705
Sandstone, green, somewhat massive.....	5	710
Shale, yellowish-green.....	15	725
Sandstone, greenish-brown, somewhat massive (2245' B.).....	10	735
Shale, red.....	60	795
Fire clay shale, streak.....		795
Sandstone, shaly (2170' B.).....	15	810
Shale, red.....	25	835
Sandstone, reddish-brown.....	5	840
Shale, yellow, sandy.....	10	850
Shale, red.....	15	865
Shale, green, sandy.....	15	880
Sandstone, Avis, shaly.....	5	885
Shale, red.....	10	895
Shale, Upper Avis, yellow, calcareous (2060' B.); contains marine fossils, pelecypods.....	25	920
Limestone, Avis, gray, shaly (2025' B.); contains marine fossils, brachiopods, pelecypods, erinoids, and bryozoa	35	955
Shale, Lower Avis, yellow, limy; contains marine fossils, pelecypods, and brachiopods (including <i>Orthotetes</i>)	10	965
Shale, red.....	45	1010
Sandstone, reddish at top, green and shaly at base (1920' B.).....	50	1060
Shale, red.....	95	1155
Shale, yellow.....	15	1170
Shale, red.....	15	1185
Sandstone, reddish-brown, hard, flaggy, with streaks of red shale (1780' B.).....	15	1200
Shale, red.....	10	1210
Shale, yellowish-green.....	10	1220
Shale, red.....	25	1245
Sandstone, reddish-brown, shaly (1720' B.).....	15	1260
Shale, red.....	14	1274
Limestone, yellow, very shaly (1705' B.).....	1	1275
Shale, red.....	15	1290
Sandstone, very shaly, mixed with red shale.....	20	1310
Shale, red, with a little sandstone.....	15	1325
Shale, green, sandy (1650' B.); contains marine fos-		

	Feet.	Feet.
19. Limestone, thinly laminated, weathers yellow....	5	253
20. Limestone, siliceous (Sample 173).....	9	262
21. Limestone, bluish-gray, weathers yellow, jointed (Sample 172).....	15	277
22. Limestone, siliceous, conchoidal weathering (Sample 171).....	20	297

Horseshoe Bend School Section.

Irish Corner District; starting near the top of a hill $\frac{1}{4}$ mile east of Horseshoe Bend School, 1.5 miles southeast of Roneeverte and measured down the highway westward to a point just west of Horseshoe Bend School; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Greenbrier Series (70'+)		
Limestone, Hillsdale, gray, cherty, fossiliferous.....	20+	20
Limestone, yellow, earthy, shaly.....	10	30
Shale, olive, yellow, crumbly.....	4	34
Clay, earthy, yellow, oolitic, cut with calcite veins	1	35
Shale, purplish-red, with streaks of yellow and olive, sandy shale.....	15	50
Limestone, ocher, brown, shaly, weathers ribbon-like and yellow.....	10	60
Shale, yellow, sandy, laminated.....	10	70
Maerady Series (180')		
Shale, red.....	100	170
Shale, variegated, brown, yellow, purplish.....	10	180
Shale, red.....	50	230
Sandstone, yellowish-brown.....	5	235
Shale, yellow, brown.....	15	250
Pocono Series (5'+)		
Sandstone	5+	255

Patton Section.

Monroe County: Second Creek District; starting just south of Patton and measured along road to Second Creek; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Greenbrier Series (285'+)		
Limestone, dark-blue, massive, tough, clay veins, chert sparse, few fos- sils (top, 2135' B.).....	20'	
Limestone, dark-blue, abundant chert, some chert nodules blend into limestone, calcite veins cut- ting chert nodules.....	50	
Limestone, gray-blue, semi-oolitic may be small fossils.....	15	
Conchoidal	40	
Patton	130	130

MEASURED SECTIONS, IRISH CORNER DISTRICT.

Irish Corner is a small district in the south central part of the county. It occupies the area bounded by the Greenbrier River, Monroe County and a line drawn from the south end of Kates Mountain to the town of Caldwell. The surface rocks include the basal part of the Mauch Chunk Series, the Greenbrier, Maccrady, and Pocono Series, and on the headwaters of Harts Run the Upper Devonian is exposed.

In the following section the sample numbers refer to chemical samples, the results of which are published in Chapter XII:

Acme Limestone Quarry Section.

Irish Corner District; measured at the Acme Limestone Quarry, near Fort Spring; arrangement in descending stratigraphic order.

	Thickness.	Total.
	Feet.	Feet.
Greenbrier Series—Alderson Member (49' +)		
1. Shale, brown, weathers yellow, sandy, calcareous.....	30	30
2. Limestone, blue, hard.....	8	38
3. Limestone, yellow, very fossiliferous.....	4	42
4. Limestone, sandy.....	2.5	44.5
5. Limestone, impure, hard, abundant, Pentremites and Archimedes.....	5	49.5
Greenville Shale Member (8')		
Shale, yellow, calcareous, algal structure.....	8	57.5
Union Member (178')		
7. Limestone, dark-gray, crystalline (Sample 183)	16	73.5
8. Limestone, white, oolitic, stylolitic vertically and horizontally (Sample 182).....	20	93.5
9. Limestone, bluish-gray, crystalline, oolitic in part, very fossiliferous (Sample 181).....	47	140.5
10. Limestone, light-gray, argillaceous (Sample 180)	5	145.5
11. Limestone, bluish-gray, oolitic, fossiliferous (Sample 179).....	18	163.5
12. Limestone, grayish-blue, weathers yellow, tough, one band of nodular chert, large erinoid stems (Sample 178).....	10	173.5
13. Limestone, gray, oolitic, Pentremites, stylolitic vertically and horizontally.....	26'	
14. Limestone, dark-gray, crystalline... 4	4	
15. Limestone, gray, fossiliferous (Sample 176).....	16	219.5
16. Limestone, grayish-blue, very fossiliferous, a few chert nodules (Sample 175).....	16	235.5

		Feet.	Feet.
Shale, Patton, weathered horizon of very irregular sandy chert with red clay (erosion surface?, 5-10')..	10		140
Limestone, light-blue, abundant light-gray chert, some nodules quite large, numerous battered bryozoa and erinoid stems.....	30		
Shale, olive-green and sandy (1925' B.)	10	Sinks Grove	60
Limestone, blue-black, massive, good	5		
Limestone, weathers sandy.....	5		
Limestone, light-blue, weathers white, clay seams.....	5		
Limestone, light-yellow, argillaceous	5		
Limestone, light-gray, massive, black chert (top, 1900' B.).....	5		
Limestone, gray-blue, massive, one band of black nodular chert.....	12		
Shale, yellow, sandy, limy.....	3	Hillsdale	85
Limestone, light-blue, yellowish, sandy, shaly, with chert at base....	15		
Limestone, much weathered, partly concealed, with chert remaining in place, to Second Creek.....	50		
			285

MEASURED SECTIONS, FORT SPRINGS DISTRICT.

Fort Springs is a small triangular-shaped district lying just north of Irish Corner District. The outcropping rocks range from the Bluefield Group of the Mauch Chunk to and including the upper part of the Pocono Series.

Hawver School Section—West.

Fort Springs—Blue Sulphur District; starting $\frac{1}{2}$ mile west of Hawver School on Muddy Creek Mountain; measured with aneroid eastward to the road forks at 2440' L., then with the road down the west side of the mountain to the top of the Alderson Limestone. Measurements shown for that portion above 2440' are greater than true vertical and for that portion below 2440' the measurements are less than true vertical. Arrangement in descending stratigraphic order.

Thickness. Total.
Feet. Feet.

Mauch Chunk Series—Bluefield Group (415'+).

Sandstone, white, massive, Droop (in part) (top, 2640')	25	25
Shale, yellow, olive, sandy.....	90	115
Limestone, impure, shaly, fossiliferous, Reynolds (Coll. 114) (top, 2525').....	15	130
Shale, olive, sandy.....	35	165
Shale, red sandy.....	35	200
Limestone, blue, gray, impure, heavy-bedded, Glenray (top, 2440' L.).....	40	240
Shale, yellow, olive, sandy, and concealed.....	175	415

record of the coal test boring above illustrates some of the variations in lithology found in the Pottsville within short distances:

Duo Section.

Meadow Bluff District; measured with aneroid, starting just above Duo and continuing along the road southwestward down the mountain to the C. & O. Railroad tracks. The measurements are slightly greater than true vertical due to a dip of about 25'. Arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River Group (277')		
Coal blossom, Sewell "A" (3470' B.) (No. 8 on Map II).....
Sandstone, brown, irregular bedding, Lower Guy-andot	10	10
Shale, brown to gray.....	27	37
Shale, black, Hartridge.....	8	45
Coal, Sewell, at old opening (No. 150 on Map II) (base, 3422' L.).....	3.5	48.5
Shale, grayish-brown and concealed.....	25.5	64
Coal, Welch (3395' B.).....	1	65
Sandstone, tough, grayish-white, abundant plants, some standing.....	5	70
Shale, gray to brown, sandy, fissile and concealed.....	27	97
Sandstone, brown, irregular bedding at base, Upper Raleigh	35	132
Shale, gray, fissile, 1" beds, iron-stained.....	25	157
Sandstone, brown, fine-grained, shaly.....	10	167
Concealed, shale talus.....	35	202
Sandstone, gray to brown, massive, medium-grained..	10	212
Shale, variegated.....	10	222
Shale, dark to black, iron-stained, fissile.....	5	227
Shale, brown, sandy, and concealed.....	40	267
Sandstone, gray to pink, medium-grained, massive, to C. & O. railroad track at 3190' B.....	10	277

The following coal test boring gives much information about the rocks in the upper half of the New River Group:

Raine Lumber and Coal Company Coal Test Boring No. 4— No. 7 on Map II.

Meadow Bluff District; one mile east of Duo; elevation, 4015' L.

	Thickness.		Total	
	Ft. In.		Ft. In.	
Pottsville Series—New River Group (489'±)				
Surface	10	0	10	0
Shale, dark, soft.....	20	0	30	0

Sandstone, Lower Guyandot.....	5	6	26	6						
Shale, dark.....	14	6	41	0						
Shale, dark, sandy.....	12	0	53	0						
Shale, dark.....	19	6	72	6						
Slate, black, Hartridge Black Shale.....	2	6	75	0						
Slate, black, and coal.....0' 3"	Sewell (3551')	4	3	79	3					
Coal						0	3			
Coal and slate.....						0	2			
Coal						3	7			
Shale, dark.....	8	4	87	7						
Coal, dirty.....	0	11	88	6						
Fire clay, soft.....	4	0	92	6						
Sandstone, hard, Welch and Upper Raleigh.....	77	6	170	0						
Shale, gray, sandy.....	38	0	208	0						
Shale, dark.....	21	0	229	0						
Slate	2	3	231	3						
Coal	1'	1"	Little Raleigh..	4	1	235	4			
Coal and slate.....	0	4								
Slate	0	3								
Coal and slate	1	6								
Fire clay.....	0	3	Fire clay, with coal.....	0	8	0	6	235	10	
Fire clay, soft.....	22	2								258
Shale, gray, sandy.....	43	0	301	0						
Sandstone, hard.....6' 0"	Lower Raleigh	43	0	301	0					
Shale, gray, sandy.....16						Sandstone	26	0	327	0
Sandstone, hard.....21										
Shale, dark.....	1	0	332	0						
Slate, black.....	4	10	336	10						
Coal	7	2	344	0						
Fire clay and coal.....0' 8"	75	6	419	6						
Fire clay, light, sandy.....	9	0	428	6						
Sandstone, and shale.....	26	6	455	0						
Sandstone, hard, Quinnimont.....	6	0	461	0						
Sandstone, hard, and shale, mixed, Fire Creek	75	0	536	0						
Coal, horizon?.....	0	9	536	9						
Sandstone, hard.....										
Sandstone, and shale mixed, Little Fire Creek										
Coal horizon?.....										
Sandstone, hard, Pineville.....										
Shale, dark.....										
Pottsville Series—Pocahontas Group (70' 3")										
Sandstone, hard.....13' 0"	Flat top and	60	3	597	0					
Shale, dark.....						0	6			
Sandstone and dark						1	6			
shale mixed.....						42	9			
Sandstone, hard.....	Pierpont	Sandstones	60	3	597	0				
Sandstone and coal							1	0		
spars							1	6		
Sandstone, hard.....	4	0	601	0						
Shale, dark, sandy.....	1'	5"	No. 6 Poca-	2	4	603	4			
Coal	0	1								
Sulphur streak.....	0	7								
Coal			(3027')							

No. 5E on Map II.

Meadow Bluff District: three-fourths mile west of Beech Knob; reported elevation, 3832' R.

	Thickness.		Total		
	Ft. In.		Ft. In.		
Pottsville Series—New River Group (387')					
Surface	13	6	13	6	
Sandstone, Lower Nuttall.....	19	0	32	6	
Shale, dark, sandy.....	30	0	62	6	
Fire clay, shaly.....	6	0	68	6	
Shale, dark, sandy.....	16	6	85	0	
Shale, dark, soft.....	25	6	110	6	
Slate, black.....	0	4	110	10	
Coal, dirty, Hughes Ferry.....	1	3	112	1	
Fire clay.....	1	0	113	1	
Sandstone, Middle laeger.....	19	6	132	7	
Shale, dark, sandy.....	15	0	147	7	
Shale, gray.....	15	6	163	1	
Coal, Lower laeger.....	0	1	163	2	
Fire clay.....	3	4	166	6	
Shale, gray, sandy, Lower laeger.....	25	6	192	0	
Sandstone, hard, Harvey Conglomerate.....	58	11	250	11	
Coal	0	1	251	0	
Sandstone	10	2	261	2	
Coal, Castle?.....	0	4	261	6	
Fire clay.....	1	6	263	0	
Sandstone and shale.....18' 0"	} Guyandot	39	0	302	0
Sandstone21 0					
Shale, dark.....	15	9	317	9	
Shale, sandy.....	15	0	332	9	
Shale, gray.....	6	6	339	3	
Shale, dark.....	2	0	341	3	
Coal, Sewell "A".....	1	3	342	6	
Shale, dark, gray.....	36	0	378	6	
Coal, Sewell (elevation reported 3450').....	3	5	381	11	
Fire clay, sandy.....	5	1	387	0	

The following record of a coal test boring furnishes important data concerning the character of the Pottsville rocks, below the Sewell Coal, in the general vicinity of Grassy Knob. In addition it is an important link in the chain of evidence establishing the correlation of the coal beds on Little Clear Creek Mountain. The measurements shown in this record as well as those shown in the record of boring No. 13, immediately following No. 11, must be used with caution. Unfortunately the cores were not always cut at right angles to the bedding-planes of the formations penetrated. Only parts of the cores were found but they showed a variation of 3° to 20° off vertical. The harder sandstone beds caused the greater

				Ft.	In.	Ft.	In.
Sandstone, shale streaks..	26'	10"	Lower Nuttall	86	10	123	0
Shale, dark, sandy.....	4	0					
Sandstone, broken.....	10	0					
Sandstone, shale streaks..	46	0					
Coal	0'	6"	laeger "A"..	11	1	134	1
Fire elay.....	0	6					
Shale, gray, sandy.....	9	0					
Slate	0	10					
Coal	0	3					
Fire elay, soft.....				4	5	138	6
Shale, dark, with sandstone streaks.....				11	6	150	0
Shale, dark.....				5	6	155	6
Sandstone				1	4	156	10
Shale, dark, sandy.....				17	2	174	0
Shale, dark.....				9	6	183	6
Coal, Hughes Ferry (3831').....				0	10	184	4
Fire elay, sandy.....				3	0	187	4
Sandstone, dark shale streaks, Middle laeger..				27	8	215	0
Shale, dark.....				17	0	232	0
Fire elay, soft, Lower laeger Coal horizon (?)				4	6	236	6
Sandstone, Lower laeger.....				23	0	259	6
Shale, dark, sandstone streaks.....				7	6	267	0
Coal				0	3	267	3
Fire elay, dark.....				3	3	270	6
Shale, gray, sandy, with sandstone streaks.....				57	2	327	8
Sandstone				1	1	328	9
Coal, Castle (3685').....				1	3	330	0
Fire elay.....				3	7	333	7
Shale, gray, sandy.....				1	0	334	7
Sandstone	34'	0"	Guyandot ...	37	5	372	0
Sandstone, with shale streaks	3	5					
Shale, dark, soft.....				13	6	385	6
Sandstone				5	3	390	9
Shale, dark, soft.....				14	3	405	0
Sandstone, soft.....				4	0	409	0
Shale, dark, soft.....				14	0	423	0
Slate, black.....				3	6	426	6
Bone	0'	4"	Sewell "A".....	2	0	428	6
Coal	1	8					
Fire elay				4	6	433	0
Shale, dark.....				31	0	464	0
Slate, Hartridge Black Shale.....				15	6	479	6
Coal	0'	3	Sewell (3532')	3	5½	482	11½
Black slate.....	0	0½					
Coal	3	0½					
Coal and slate.....	0	1½					
Fire elay, dark.....				1	0½	484	0
Fire elay, shaly.....				5	0	489	0

			Ft.	In.	Ft.	In.
Shale, dark.....			14	6	352	6
Coal			0	1	352	7
Gray elay shale.....			2	5	355	0
Shale, dark, sandy.....			13	6	368	6
Slate, black.....			1	0	369	6
Fire elay.....			1	6	371	0
Shale, dark, sandy.....			14	0	385	0
Shale, dark.....			1	4	386	4
Coal, bony.....			1	4	387	8
Fire elay, shaly.....			12	0	399	8
Shale, gray.....			8	8	408	4
Slate, with coal spars.....			1	1	409	5
Shale, gray.....			28	7	438	0
Sandstone, hard.....			8	8	446	8
Coal and slate mixed, Fire Creek?.....			0	6	447	2
Shale, dark, sandy.....			18	0	465	2
Bone coal, Little Fire Creek?.....			0	1	465	3
Shale, dark.....			4	0	469	3
Sandstone, hard.....	40'	0"	62	9	532	0
Shale, dark.....	2	0				
Sandstone	7	9				
Shale, dark.....	0	10				
Sandstone	12	2				
Shale, dark.....			2	2	534	2
Slate, black.....			2	0	536	2
Slate, black, bony.....			1	0	537	2
Coal, dirty.....	0'	4"	0	11	538	1
Fire elay.....	0	1				
Slate, soft.....	0	2				
Coal	0	4				
Slate, soft.....			0	2	538	3
Fire elay, shaly.....			1	3	539	6
Shale, gray.....			10	0	549	6
Slate, black, No. 8 Pocahontas Coal horizon....			0	6	550	0
Fire elay.....			0	10	550	10
Shale, gray, sandy.....			6	2	557	0

Pottsville Series—Pocahontas Group (271')

Sandstone and shale....	6'	0"	85	0	642	0
Sandstone, hard.....	79	0				
Coal, No. 6 Pocahontas.....			0	10	642	10
Sandstone with coal spars and shale spots.....			4	8	647	6
Shale, sandy.....			2	6	650	0
Sandstone, Eckman.....			51	0	701	0
Coal and sandstone.....	0'	5"	2	9	703	9
Sandstone, with coal spars	2	4				
Dark elay shale.....			0	10	704	7
Fire elay, soft.....			2	8	707	3
Shale, gray, sandy.....			4	9—	712	0
Sandstone, with coal spars, Upper Pocahontas			30	0	742	0
Shale, sandy.....			2	0	744	0
Shale, dark.....			9	0	753	0
Sandstone, with coal spars, Pocahontas Coal horizon			5	0	758	0

No. 11 on Map II.

In Meadow Bluff District; four and one-half miles east of Duo and one mile east of Job Knob; elevation, 4010' L.

	Thickness.		Total	
	Ft. In.		Ft. In.	
Pottsville Series—New River Group (557' +)				
Surface	16	0	16 0	
Shale, soft, brown.....	14	0	30 0	
Shale, gray.....	12	0	42 0	
Slate, Hartridge	8	2	50 2	
Coal, Sewell (3957').....	3	2	53 4	
Fire clay.....	0	6	53 10	
Shale, light clay, sandy.....	5	2	59 0	
Sandstone, hard, Welch?.....	32	0	91 0	
Coal, Welch?.....	1	6	92 6	
Sandstone and coal.....	0	6	93 0	
Sandstone, hard.....	5	6	98 6	
Shale, dark, soft.....	8	6	107 0	
Coal and slate mixed, Welch?.....	2	8	109 8	
Fire clay.....	2	0	111 8	
Shale, sandy, hard.....14' 4"	} Upper Raleigh	34 4	146 0	
Sandstone, hard.....20 0				
Shale, gray.....	7	9	153 9	
Slate, with coal spars.... 3' 6"	} Little Raleigh "A".....	9 11	163 8	
Slate				5 0
Coal				0 6
Slate				0 5
Coal	0 6	7 4	171 0	
Fire clay.....	11 5			
Shale, dark.....	1	5	183 10	
Slate				
Slate, with coal spars.... 0' 10"	} Little Raleigh	11 2	195 0	
Fire clay, soft.....				5 0
Gray clay shale.....				4 10
Slate, black.....				0 4
Coal	0 2	0 10	195 10	
Fire clay.....				
Shale, gray, sandy.....	33	2	229 0	
Shale, dark.....	7	0	236 0	
Coal and slate.....	0	5	236 5	
Fire clay.....	4	7	241 0	
Sandstone, hard, white, Lower Raleigh.....	23	0	264 0	
Shale, dark.....	10	3	274 3	
Coal	0	3	274 6	
Shale, gray, sandy.....	14	0	288 6	
Shale, dark.....	10	0	298 6	
Fire clay, sandy.....	4	6	303 0	
Shale, dark.....	23	6	326 6	
Slate, black.....	7	0	333 6	
Coal, bony..... 0' 1"	} Beckley	4 6	338 0	
Fire clay and slate.....				3 7
Coal				0 2
Fire clay.....				0 2

	Ft.	In.	Ft.	In.
Shale, sandy and sandstone, Lower Pocahontas Sandstone?.....	17	0	777	0
Slate, black, No. 2 "A" Pocahontas Coal horizon?	0	11	777	11
Fire clay, soft.....	1	7	779	6
Shale, gray, sandy.....	3	0	782	6
Sandstone and shale.....	6	6	789	0
Shale, dark.....	8	0	797	0
Slate, black, No. 2 Pocahontas Coal horizon....	5	0	802	0
Fire clay, sandy.....	6	0	808	0
Sandstone and shale mixed.....	5	6	813	6
Slate, black, No. 1 Pocahontas Coal horizon....	2	6	816	0
Sandstone and shaly clay.....	12	0	828	0
Mauch Chunk Series (11'+)				
Fire clay, hard.....	0	6	828	6
Shale, green.....	10	6	839	0

The following record of a coal test boring confirms the correlation of the coal beds on Little Clear Creek Mountain. As noted in the comment preceding coal test boring No. 11 above, the measurements shown in this record must be used with caution:

Gauley Coal Land Company Coal Test Boring No. 1— No. 13 on Map II.

Meadow Bluff District; six miles north 77° E. of Anjean, on Little Clear Creek Mountain; elevation, 3808' L.

Clear Creek Mountain; elevation, 3508' L.		Thickness.	Total	
		Ft. In.	Ft. In.	
Pottsville Series—New River Group (175'+)				
Surface		15 6	15 6	
Shale, gray.....		0 4	15 10	
Sandstone, hard.....33' 8"	} Quinnimont	40 2	56 0	
Sandstone, with shale spots				1 0
Sandstone, hard.....				5 6
Coal, Fire Creek?.....		0 2	56 2	
Sandstone and coal spars.....		2 6	58 8	
Sandstone		1 4	60 0	
Shale, gray, soft.....		7 4	67 4	
Black slate, soft, coal spars.....		0 10	68 2	
Fire clay, soft.....		1 2	69 4	
Shale, gray, soft, broken.....		2 8	72 0	
Shale, gray, sandy.....		3 0	75 0	
Sandstone, fine-grained. Pineville.....		60 8	135 8	
Shale, dark, sandy.....		8 4	144 0	
Sandstone, hard, coarse.....		12 7	156 7	
Shale, gray.....		3 5	160 0	
		2 6	162 6	

	Ft.	In.	Ft.	In.
Clay shale, with fossils.....	1	0	163	11
Shale, gray, sandy.....	11	1	175	0

Pottsville Series—Pocahontas Group (330' 6")

Sandstone, hard, fine-grained, Flattop.....	33	3	208	3
Shale, dark.....	1	1	209	4
Sandstone	2	10	212	2
Slate, black.....	0	7	212	9
Bone coal, No. 7 Pocahontas.....	0	1	212	10
Clay, shale, gray.....	0	10	213	8
Shale, gray, sandy.....	15	0	228	8
Sandstone11' 10"	23	8	252	4
Sandstone, occasional				
coal spars.....10 6				
Sandstone, with coal	6	0	258	4
spars and shale.....1 4				
Shale, gray.....	2	0	260	4
Shale, soft, Royal?.....	3	6½	263	10½
Bone coal.....0' 2 "				
Coal2 5				
Bone coal.....0 1½				
Fire clay.....0 6				
Coal0 3				
Slate0 0½	1	0½	264	11
Coal0 0½				
Fire clay, dark.....	15	6	280	5
Shale, gray.....	0	6	280	11
Coal	13	1	294	0
Shale, gray, sandy.....	0	3	294	3
Slate, with coal spars.....	10	6	304	9
Clay shale.....	8	6	313	3
Shale, gray.....	15	7	328	10
Shale, gray, sandy.....	0	5	329	3
Slate, gray.....	0	8	329	11
Coal	1	5	331	4
Fire clay, dark.....	2	4	333	8
Shale, dark.....	0	9	334	5
Coal	2	2	336	7
Fire clay.....	12	3	348	10
Shale, gray.....	1	4	350	2
Fire clay.....	3	8	353	10
Shale, gray, sandy.....	30	0	383	10
Sandstone, Upper Pocahontas.....	1	0	384	10
Shale, dark.....	0	1	384	11
Bone coal, No. 3 Pocahontas.....	4	7	389	6
Shale, dark.....	2	8	392	2
Fire clay, soft.....	20	4	412	6
Shale, dark.....	22	0	434	6
Shale, dark, sandy and sandstone, Lower Poca-				
hontas Sandstone?.....	10	6	445	0
Shale, dark.....	2	9	447	9
Sandstone, hard, fine, coal spars.....	1	4	449	1
Shale, dark.....				

	Feet.	Feet.	
Flre clay. streak, (laeger "B" Coal horizon) (2870' B.).....	85	
Sandstone, massive, eoarse, soft, Lower Nuttall	115	200	
Coneealed	40	240	
Spring, Hughes Ferry Coal horizon (2710' B.).....	240	240'
Slate, blaek.....	15	255	
Coneealed and shale.....	25	280	
Sandstone, Harvey.....	40	320	
Coneealed	60	380	
Sandstone, massive, Guyandot, and coneaeled in steep bank.....	143	523	
Shale, dark, Hartridge.....	2	525	
Coal, soft..... 2' 4"	} (5' 1") Sewell (2420' L.).....	530	290'
Slate, bony..... 0 7			
Coal, bony..... 1 6			
Coal, soft, good..... 0 8			
Sandstone and coneaeled to stratigraphie level of Well (S).....	7	537	

Continued by Ganley Coal Land Company (Granville
O'Dell) No. 1 (No. 8 on Map II) Well Reeord:

Conduetor	16	553	
Slate shell (hole full of water at 20').....	64	617	
Coal Blossom, Welch?.....	2	619	
Slate, blaek.....	148	767	
Sand. gray. Lower Raleigh.....	20	787	
Lime, white.....	30	817	
Coal, Beekley?.....	5	822	
Slate, shell, dark (hole full of water at 304').....	68	890	
Coal, Fire Creek?.....	5	895	
Slate, dark.....	7	902	
Lime, gritty (hole full of water at 385').....	55	957	
Slate, dark.....	15	972	
Llme, dark, hard.....	65	1037	
Sand, gray, hard, Plneville (hole full of water at 510').....	20	1057	

Mauch Chunk Series (1418')

Red roek.....	120	1177	
Sand, gray, hard, Princeton (gas at 694', steel- line measure).....	140	1317	
Slate and shells, dark.....	175	1492	
Red roek and lime shells.....	75	1567	
Red roek lime shells.....	125	1692	
Lime shells.....	25	1717	
Lime shells, red roek.....	75	1792	
Llme, gritty, Terry?.....	40	1832	
Sand, very hard.....	45	1877	
Slate, soft.....	15	1892	
Lime, very hard.....	20	1912	
Slate, soft.....	6	1918	
Lime, very hard.....	12	1930	
Slate, soft.....	10	1940	
Llme, broken up.....	15	1955	
Slate, white.....	10	1965	

Shale, dark, sandy.....	14	0	469	0
Shale, dark.....	7	0	476	0
Slate, soft.....	3	6	479	6
Bone coal, No. 2 Pocahontas?.....	0	2	479	8
Shale, dark.....	0	8	480	4
Fire clay, sandy.....	2	8	483	0
Shale, dark, sandy.....	10	4	493	4
Slate, black.....	0	6	493	10
Dark clay shale.....	0	10	494	8
Coal, No. 1 Pocahontas?.....	0	6	495	2
Fire clay.....	2	3	497	5
Shale, gray.....	2	3	499	8
Sandstone	3	10	503	6
Shale, gray.....	2	0	505	6
Mauch Chunk Series (35' 6" +)				
Fire clay and gray and green shale mixed.....	12	0	517	6
Gray and green and red shale mixed.....	6	9	524	3
Fire clay, hard, sandy.....	5	2	529	5
Fire clay and green shale.....	1	4	530	9
Red and gray and green shale mixed.....	8	3	539	0
Red shale.....	2	0	541	0

The following section and well record, with remarks about the same, by D. B. Reger, are taken from the Nicholas County Report, W. Va. Geological Survey, pp. 174 to 177; 1921. It provides much information on the subsurface strata of southeastern Nicholas and western Greenbrier Counties:

"In the following section, arranged in descending order the surface portion was measured with aneroid, starting at the top of the plateau just east of Snow Hill School, and extends northeastward with the strike of the rocks along the public highway to an opening in the Sewell Coal at the foot of the mountain, one-third mile west of Hominy Creek. The lower portion is the record of the Gauley Coal Land Company (Granville O'Dell) No. 1 Oil Test Well (No. 8 on Map II) located just west of Hominy Creek, and 1.4 miles southward from Hominy Falls, and being 0.4 mile northwest of the foot of the measured section. Inasmuch as the Sewell Coal is opened within a few feet of the well and only 7 feet above the level of the top of the hole, no difficulty was experienced in making connection with the stratigraphic measurement described above. The well was drilled by the Wick-Laing Oil and Gas Company, its record having been furnished by Mr. C. M. Boyd, Secretary, of Youngstown, Ohio. It was abandoned as a dry hole, only a small amount of gas having been found in the Princeton Sandstone:"

Hominy Falls Section, Wilderness District, Nicholas County.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River Group (1057')		
Sandstone, massive, from road fork, Upper	75	75

Lime, hard.....	13	2008
Slate	10	2018
Sand, Maxton.....	39	2057
Slate, soft.....	12	2069
Slate and lime, broken up.....	118	2187
Lime, white.....102'	} Hinton (Little Lime)	215 2402
Slate, white.....15		
Lime		
Sand, Webster Springs.....	35	2437
Slate, black.....	5	2442
Sand	13	2455
Pencil Cave.....	20	2475
Greenbrier Series (393')	393	2868
Big Lime.....		
Pocono Series (311')	12	2880
Sand, Keener.....	3	2883
Red rock.....	16	2899
Sand, Big Injun.....	49	2948
Slate	19	2967
Squaw Sand.....	110	3077
Sand shells.....	102	3179
Sand, Berea.....		
Catskill and Chemung Series (1111')	363	3542
Limo shells, slate.....	20	3562
Lime	85	3647
Shells and slate.....	60	3707
Lime	150	3857
Slate and shells.....	125	3982
Lime, gritty.....	308	4290
Lime, gritty, and shells (steel-line measure).....		
Began spudding, October 26, 1915; shut down May		
11, 1916, 6 p. m.		
10" easing, 345'; 8¼" easing, 975'; 6⅝" easing, 1658'.		
Total depth of hole.....		3753

Kieffer Section.

Meadow Bluff District; starting at the edge of Cross Mountain; descending to Beaver Creek; measured with aneroid along road and arranged in descending stratigraphic order.

		Thickness.	Total.	
		Feet.	Feet.	
Mauch Chunk Series—Princeton Group (35')				
Conglomerate, massive,	} Princeton Conglomerate (top, 2875')	35	35	
white				15'
Rotten sand and pebbles.....				10
Coneealed				10
Mauch Chunk Series—Hinton Group (300'+)				
Coneealed, with red and olive shale.....		100	135	
Shale, red.....		75	210	
Sandstone, greenish-yellow, shaly, cross-bedded.....		25	235	
Shale, red.....		80	315	
Shale, red.....		10	325	
Shale, yellow, Avls.....				
Shale, yellow, shaly, fossiliferous (top				

Williamsburg District; starting at the top of Cold Knob and traversing generally southward 2 miles to the S. W. Hinkle well No. 4 on Map II and combined with the log of that well; measurements for that portion of the Bluefield Group above the Hinkle well are slightly greater than true vertical owing to the dip of the rocks; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Pottsville Series (315'+)		
Concealed from top of Cold Knob, not examined in detail.....	315	315
Mauch Chunk Series—Bluestone Group (395')		
Shale, red and concealed (top, 4030' B.).....	15	330
Shale, red, variegated, with thin flaggy, greenish, argillaceous sandstones.....	380	710
Mauch Chunk Series—Princeton Group (80')		
Sandstone, greenish-brown, massive, many quartz pebbles, mostly concealed, but abundant drift, Princeton Conglomerate (top, 3635' B.).....	80	790
Mauch Chunk Series—Hinton Group (475')		
Shale, red, and concealed.....	25	815
Sandstone, greenish-gray, calcareous, thick-bedded, hard	20	835
Sbale, variegated, interbedded with green to red argillaceous sandstones, and concealed.....	110	945
Limestone, argillaceous, fossiliferous, Avis (top 3400' B.).....	15	960
Shale, red, with brown to green sandstones, partly concealed	270	1230
Sandstone, reddish-brown, cross-bedded, medium-hard, Stony Gap (top, 3115' B.).....	35	1265
Mauch Chunk Series—Bluefield Group (935')		
Shale, red, variegated, with argillaceous, green to brown sandstones, and concealed to top of Hinkle Well	545	1810
(Continued with record of Hinkle Well No. 4 on Map II—elevation, 2535' B.):		
Soil	5	1815
Sandstone, Droop.....	125	1940
Sbale, blue.....	70	2010
Shale and sand, Webster Springs.....	45	2055
Sbale, gray	145	2200
Greenbrier Series (475')		
Shale and lime (top, 2145' B.).....	5	2205
Lime	435	2640
Shale, blue, coarse, and some lime.....	35	2675
Macerady Series (80')		
Shale, red $\frac{1}{2}$, gray $\frac{1}{2}$	55	2730
Shale, red $\frac{1}{2}$, gray $\frac{1}{2}$	25	2755
Pocono Series (205')		
Shale, black, and gray limy sand.....	105	2860
Gray lime.....	40	2900
	25	2925

Williamsburg District joins Meadow Bluff District on the northeast and east. It is shaped somewhat like an hour-glass with the narrow part at Grassy and Cold Knobs. The fan-shaped northwest end of the district includes the drainage area of Laurel Creek and Little Laurel Creek extending from Beech Ridge on the south to Sugar Knob on the north. The southern half of the district is centered on the town of Williamsburg and includes most of the drainage area of Sinking and Culverson Creeks. The outcropping rocks range from the New River Group of the Pottsville down to the top of the Pocono.

In the following record the Hinton Group of the Mauch Chunk appears to be too thin, due, no doubt, to a northwest dip:

Roach Run Section.

Williamsburg District; measured along the road on southeast end of Cross Mountain near Roach Run; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Bluestone Group (90' +)		
Sandstone, grayish-brown, weathers white (base, 3125' B.).....	15	15
Shale, brown to yellow, and concealed.....	75	90
Mauch Chunk Series—Princeton Group (25')		
Sandstone, Princeton, gray, massive, conglomerate....	25	115
Mauch Chunk Series—Hinton Group (520')		
Concealed	15	130
Shale, buff, sandy.....	40	170
Shale, red.....	80	250
Sandstone, grayish-brown, flaggy, shaly at base.....	10	260
Shale, red, sandy.....	50	310
Sandstone, yellow to olive, calcareous, shaly.....10'	Avis Limestone (Coll. 132)..... (top, 2830' B.)	25
Limestone, blue, banded, shaly		
Shale, yellow, fissile..... 5		
Shale, red.....	40	375
Sandstone, reddish-brown, massive, cross-bedded.....	35	410
Shale, red and concealed.....	190	600
Sandstone, Stony Gap, reddish-brown at top, more massive and grayish-brown at base.....	35	635
Mauch Chunk Series—Bluefield Group (280' +)		
Shale, red.....	20	655
Sandstone, reddish-brown.....	5	660

Thickness. Total.
Feet. Feet.

Catskill Series (50')

Red rock..... 50 3010

Chemung Series (400'+)

Dark fine sand lime, fairly good sand..... 175 3185
 Sand, gray, coarse..... 45 3230
 Lime and coarse sand..... 5 3235
 Sand, light, hard, gray..... 15 3250
 Lime, light, hard and gray..... 24 3274
 Sand, gray, soft (some water)..... 11 3285
 Sand, gray, hard..... 15 3300
 Lime, shale 35 3335
 Lime, black, hard..... 18 3353

Alta Section.

Williamsburg-Blue Sulphur District line; starting 1½ miles west of Alta and measured southeastward along the Midland Trail; arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Mauch Chunk Series—Bluefield Group, (293'+)		
Sandstone, Droop, brown to grayish-white, massive, cross-bedded, makes cliff at quarry.....	50	50
Shale, yellow, fissile (Coll. 69 at base).....	50	100
Limestone, Reynolds, shaly (Coll. 68).....	3	103
Shale, yellow, sandy, fissile, thin streaks of red.....	30	133
Limestone, gray, hard, cut with calcite veins.....10'	60	193
Limestone, tough, siliceous.....10'		
Limestone, blue, hard, broken.....10'		
Shale, Lillydale, dark, carbonaceous, fissile, mica- ceous, plants and pelecypods at base (Colls. 70, 72)	100	293

Greenbrier Series (610')

Limestone, Alderson, bluish-gray, siliceous, upper part shaly (Coll. 80), lower part more massive (Coll. 79).....	40	333
Shale, Greenville, yellowish-green to dark (Coll. 78)	10	343
Limestone, blue, hard, massive, some oolite, very fossiliferous, upper part; light-gray to white oolite in lower part; stylolitic; abundant marine fossils; (Coll. 77 from upper part); (Coll. 76 from lower part).....120'	195	538
Limestone, bluish-gray, massive..... 75		
Shale, dark to yellow (Coll. 75)..... 10'	135	673
Limestone, gray, massive, loose chert, fragments (Coll. 74 near base)125		
Shale, yellow, sandy, few fossils.....	40	713
Limestone, yellowish-gray, weathers yellow, mud-cracks, (photograph, Plate XXI).....15'	35	748

The following section, measured just across the county line from Sugartree Bench, together with comments about the same by P. H. Price, is taken from pages 111 to 113 of the Pocahontas County Report⁵. It is now believed that approximately 150 feet should be added to the interval between the Sewell Coal and the top of the Mauch Chunk. This is in addition to the amount to be added to the Mauch Chunk as noted in the comment.

"The following section, measured by the writer and arranged in descending stratigraphic order, affords a view of the basal Coal Measures including the Sewell Coal. A complete section of the Mauch Chunk Series was measured by aneroid, using vertical measurements on rising strata, thus shortening its true thickness by approximately 400 feet. An attempt was made to reopen the Sewell Coal here at the prospect of the Preston Clark Heirs, from which considerable coal was mined several years ago. In order to get a true thickness several hours were spent by the writer, Walter Mason, and Lee Clark, one of the heirs, in facing up the coal as indicated below:"

Briery Knob Section.

Pocahontas County, Little Levels District; beginning at high point on Briery Knob and following southeastward along the old coal road to forks near Mt. Lebanon Church and thence northeastward to Hills Creek.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River Group (431')		
Sandstone, (Harvey Conglomerate), grayish-brown, weathering almost white, coarse.....	15	15
Coneaeled in flat bench.....	90	105
Sandstone, Guyandot, white, weathering to white sand, coarse; small, white quartz pebbles.....	55	160
Shale, Hartridge, mostly coneaeled, but 4' of dark carbonaceous shale with plants and Naiadites? visible	35	195
Coal, good, clean....2' 4"	6.4 (No. 11 on Map II)	201.4
Shale1 10		
Coal, good, clean....1 3		
Coal, coneaeled.....1 0		
Coneaeled	24.6	226
Sandstone, Upper Raleigh (Sharon), brown and white, coarse, cross-bedded, with white quartz pebbles	105	331
Coneaeled	95	426
Shale, dark, carbonaceous, Fire Creek Coal horizon?	5	431

	Thickness. Feet.	Total. Feet.
Limestone, light-gray, stylolitic structure, fossiliferous, (quarry, average dip, 23° N. W.).....	25'	823
Limestone, dark-gray, massive.....	50	
Limestone, Hillsdale, blue. gray, massive, nodules of irregular black chert (Coll. 71).....	80	903
Maccrady Series (75'±).....	75±	978
Shale, red.....		

MEASURED SECTIONS, FALLING SPRINGS DISTRICT.

Falling Springs is the northernmost district in the county. It includes most of the drainage area of North and South Forks of Cherry River, most of the drainage area of Spring Creek, and the drainage area of several small streams on the east side of Greenbrier River north of the village of Anthony. The surface rocks range from the Kanawha Group of the Pottsville down to the middle Chemung. Sections measured in this district afford the best detailed measurements of the Greenbrier Limestone available in the county.

Little Rocky Run Section.

Falling Springs District; measured with aneroid starting at the top of the high knob (elevation, 4030' L.) north of Little Rocky Run, traversing south to Little Rocky, thence westward to South Fork of Cherry River.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River and Pocahontas Groups (670'±)		
Concealed to top of bench.....	115	115
Sandstone, Welch?.....	20	135
Concealed	10	145
Sandstone, makes cliff.....	30'	200
Concealed	5	
Sandstone, massive, white, with white quartz pebbles 20		
Concealed	55	
Sandstone, coarse, conglomerate.....	460	660
	10	670
Mauch Chunk Series (650'±)		
Concealed	120	790
Sandstone, coarse.....	10	800
Concealed	30	830
Shale, red, and concealed.....	185	1015
Sandstone, brown, Princeton?.....	50	1065
	10	1075
	10	1085

Two and one-half miles north of Alderson a small fault was mapped that involves the basal members of Mauch Chunk Series. The Lillydale Shale is overturned with an 80-degree dip to the southeast, while a short distance northwest the same shales are seen in a normal position with a 10-degree northwest dip. It is not possible to determine the amount of displacement but it must be small since the shales are rarely more than 100 feet thick.

Two miles farther north and $2\frac{1}{2}$ miles southeast of Blue Sulphur Springs a similar condition was noted but here the upper beds of the Greenbrier Series are exposed at the fault. The Alderson Limestone is slightly overturned with an 85-degree dip to the southeast. Above the Alderson (to the northwest) is a concealed interval of about 20 feet and the next visible bed is a limestone that is probably the Glenray. The latter bed which contains a number of small rectangular blocks of limestone that have been cemented together, is right side up and has a northwest dip of 5 degrees. The Lillydale Shale that would normally occur between these two limestones should have a thickness of about 100 feet, which indicates a displacement of about 80 feet. No entirely satisfactory explanation can be given to account for the rectangular blocks in the limestone but the most plausible theory is that of jointing, plus solution and cementation. Joints that are closely akin to true cleavage joints have been developed in the Alderson Limestone.

Along the Midland Trail (U. S. Route 60) 1.4 miles northwest of Alta an apparent fault was noted that is similar to the two just described. The Glenray Limestone is standing nearly vertical while a short distance northwest the Droop Sandstone is nearly horizontal. No absence of beds could be proved although the interval between the Droop Sandstone and Glenray Limestone is smaller than would be expected.

A small vertical fault with a displacement of five feet was noted $\frac{3}{4}$ mile northwest of Oscar P. O. The lower part of the Alderson Limestone and the Greenville Shale are ex-

described, is a major overthrust, located along the west side of Beaver Lick Mountain. It receives its name from the small settlement of Burr, in Pocahontas County, on the west side of Beaver Lick Mountain, $\frac{1}{2}$ mile north of the Greenbrier County line. The outcrop of the fault-plane is usually concealed so that its exact location and extent (as shown on Map II) is, in some respects, approximated. On the headwaters of Little Creek the fault contact was found and at this point the Red Medina sandstones and sandy shales are thrust up and over the Marcellus black shales. The red sandstones and sandy shales have been so mashed and metamorphosed that it was not possible to distinguish the true bedding-planes and the underlying black shales show numerous enfolded drag folds. The thickness of rocks that normally occur between the Marcellus and the Red Medina is about 1700 feet, which with the 800 feet of Red Medina exposed and an undetermined thickness of the Marcellus, indicates a total throw at this point, of more than 2500 feet. Cross-section A—A' was drawn to illustrate the fault at the point just described and it is reproduced on the margin of Map II.

CHAPTER V.

MEASURED SECTIONS.

INTRODUCTION.

The surface or outcropping rocks of Greenbrier County include the Quaternary, with Recent and Pleistocene deposits, and a considerable portion of the Paleozoic, including the lower portion of the Pennsylvanian, the Mississippian, the Devonian, and the greater part of the Silurian sediments. A classification of these beds, approximating 14,385 feet of rocks, is shown in Figure 7, pages 131-133.

The Quaternary Rocks are represented by clays, gravels, and sand beds, present along the river and creek valleys, and by river-terrace deposits now resting many feet above the present streams. Some of these terraces are undoubtedly of Pleistocene age, although there is no evidence of glacial origin. These two types of formations, which make up the best farming lands along the larger streams, are represented on Map II under Alluvium.

The Kanawha, New River, and Pocahontas Groups of the Pottsville Series of the Pennsylvanian, with an approximate thickness of 1,540 feet of strata, are the youngest of the Paleozoic rocks present, and they undoubtedly once covered Greenbrier County. They are now confined to the western part of the county, their eastern extension having been removed by erosion.

The Mauch Chunk Series of the Mississippian is subdivided into four groups, Bluestone, Princeton, Hinton, and Bluefield, and contains approximately 2,805 feet of sediments, constituting a considerable portion of the surface of Greenbrier County west of the Greenbrier River.

The Greenbrier Series of the Mississippian contains about 750 feet of rocks that are predominantly calcareous. Its best

brier River. The new road cuts along the Greenbrier River (Route 219) afford many good exposures that offer opportunity for study.

The outcrop of the Maerady Series of the Mississippian lies immediately beneath the Greenbrier Series. It is found in a belt west of the Greenbrier River the entire length of the county, from Monroe on the south to Pocahontas on the north. It varies in thickness from 60 feet at the northern end of the county to 250 feet at the southern end as compared to 700 feet or more at its type locality in Smyth County, Virginia.

The Pocono Series comprises the basal members of the Mississippian¹ in Greenbrier County and is seen to its best advantage along the Greenbrier River. This series decreases in thickness from approximately 600 feet at its best development in this area, to some 205 feet, in the Hinkle Well near Trout P. O.

The Devonian outcrops in Pocahontas County are confined to the area east of the Greenbrier River, with the exception of the Catskill Series which outcrops along the river and occasionally west of it. The entire assemblage has a thickness of approximately 6,390 feet as compared to 11,000 feet in northeastern West Virginia. The Chemung Series retains a good development throughout the county and may be seen in its entirety along the State road east of Caldwell. Apparently all of the remaining series are retained in this area.

The Silurian rocks comprise the oldest sediments exposed in the county and are limited to the region east of the Greenbrier River along Beaver Lick Mountain. Their maximum thickness is approximately 2,050 feet.

In the area west of the Greenbrier River the gently dipping beds permit the measurement of numerous vertical sections, and the study in detail of the character of the surface rocks, while east of this area where the rocks are steeply dipping, additional sections have been obtained along streams and road cuts, where it was possible to determine, approximately, the vertical thicknesses by trigonometric computation. All of these sections appear in the following pages.

Fayette County, Quinnimont District; measured with aneroid from the road summit, 0.7 mile northwest of Turniphole Mountain, southward along the hill road to the top of the Mauch Chunk Red Shales.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River and Pocahontas Groups		
(311')		
Sandstone, grayish-white, Pineville (?).....	45	45
Concealed	5	50
Sandstone, shaly.....	35	85
Concealed	5	90
Shale, black, Royal, Lingula fossil shells abundant.....	9	99
Coal, soft, No. 6 Pocahontas (No. 411 on Map II).....	4	103
Shale, gray and dark.....	10	113
Coal, blossom, heavy, No. 6 Pocahontas, lower bench (No. 411A on Map II).....	2	115
Concealed and shale, sandy.....	39.5	154.5
Coal, slaty, (6"), No. 4 Pocahontas.....	0.5	155
Sandstone, coarse, broken, Upper Pocahontas.....	13	168
Shale	1.5	169.5
Coal, soft, (5"), No. 3 Pocahontas.....	0.5	170
Shale, sandy	5	175
Shale, flaggy, and sandy.....	29	204
Coal, soft 0' 8" } No. 2 Pocahontas.....	1	205
Shale, gray 0 1 } (10")		
Coal, soft 0 1 }		
Sandstone, shaly at bottom.....	15	220
Concealed and sandstone.....	55	275
Fire clay shale.....	5	280
Concealed and sandstone to red shale, top of Mauch Chunk Series.....	31	311

Sims Station Section.

Meadow Bluff District; starting along road ascending Sims Mountain one mile south of Sims Station, measured with aneroid. Rewritten in descending stratigraphic order. The measurements are somewhat greater than true vertical owing to a dip of about 125 feet.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River and Pocahontas Groups (370'+)		
Coal, reported, Fire Creek? (3255' B.).....
Concealed	5	5
Sandstone, medium-grained, gray to brown.....	10	15
Concealed	28	43
Coal, supplied from other side of hill (No. 372 on Map II) Little Fire Creek?.....	2	45
Concealed	20	65
Sandstone, brown.....	5	70
Concealed, with sandy shale.....	25	95

collections were made and reference is often shown in parenthesis by number, referring to the particular zone described. These collections have all been examined by the late Dr. John L. Tilton and/or Prof. Dana Wells, and the results of their examinations are published as Chapter XIV, Notes on Paleontology.

Additional fossil collections were made by Dr. David White, David B. Reger, and Paul H. Price with particular emphasis on the fossil flora, but the results of these collections will not be available for this report.

MEASURED SECTIONS, MEADOW BLUFF DISTRICT.

Meadow Bluff District, the largest district, occupies a vast area in the extreme western part of Greenbrier County. It is bounded on the west by Fayette County and on the northwest by Nicholas County. The district line, along the northeast, follows the crest of Beech Ridge to Grassy Knob, thence southwest along Old Field Mountain, Buffalo Mountain, and Meadow Mountain to Clintonville. From this point the district line turns more to the west passing through Smoot and reaches the Greenbrier-Fayette County line 3.2 miles southeast of the town of Springdale (Fayette County). Its surface rocks range from the Kanawha Group of the Pottsville down to the base of the Hinton Group of the Mauch Chunk. All of the commercial coal mines operating in the county are located in this district.

The following section, prepared by Ray V. Hennen², was measured along the eastern boundary of Fayette County and shows the development of the Pocahontas Group of the Pottsville in eastern Fayette and southwestern Greenbrier Counties:

²Hennen, Ray V., Fayette Report, W. Va. Geol. Survey, p. 219; 1919.

	Feet.	Feet.
Shale, "fire clay," plant fossils abundant.....	2.2	63
Concealed	12	75
Sandstone, gray to pink, weathers brown, irregularly bedded	30	105
Shale, chocolate colored, with coal streaks, Little Fire Creek? (2940' B.) (No. 373 on Map II).....	3	108
Sandstone, much weathered, limonite veins.....	20	128
Shale, almost a sandstone, much weathered.....	35	163
Sandstone, massive, fine-grained.....	7	170
Concealed and sandstone, mostly concealed.....	35	205
Coal, No. 6 Pocahontas? (2840' B.) (No. 415 on Map II)	0.2	205.2
Concealed, sandy.....	59.8	265
Coal, soft, impure, No. 4 Pocahontas? (2780' B.) (No. 470 on Map II).....	1	266
Shale, "fire clay," abundant plant fossils.....	2	268
Sandstone, weathered, brown, loosely cemented, Upper Pocahontas.....	55	323
Concealed, sandy.....	77	400
Shale, black, No. 1 Pocahontas Coal horizon? (No. 500 on Map II).....	2	402
Estimated interval to top of Mauch Chunk Series.....	25	427

Goddard Mountain Section—West Side.

Meadow Bluff District; starting at a point near the top of Goddard Mountain and measured with aneroid down the trail on the west side of the mountain to Boggs Creek. The measurements are greater than true vertical owing to a dip of about 70 feet. Arrangement in descending stratigraphic order.

		Thickness.	Total.
		Feet.	Feet.
Pottsville Series—New River and Pocahontas Groups (415'±)			
Sandstone, cap rock (base, 3180' B.).....		25	25
Concealed		95	120
Coal blossom, (3085' B.).....		0	120
Sandstone, massive, cross-bedded.....		25	145
Sandstone, brown, cross-bedded.....		25	170
Concealed		139	309
Shale		3.5	312.5
Coal	0' 2 "	} No. 3 Pocahontas (base, 2890' B.)..... 2.5 (Mine No. 486 on Map II)	315
Shale	0 1		
Coal	0 1		
Shale	0 5		
Coal	0 1½		
Shale	0 2		
Coal, clean, good.....	1 5½		
Shale floor.....		0	315
Concealed		100	415
Mauch Chunk Series—Bluestone Group (300')			
Concealed, (top, estimated, 2790').....		300	715

	Fect.	Feet.
Shale, sandy, gray to brown.....	5	120
Sandstone, medium-grained, gray to brown, irregular bedding, Flattop?.....	30	150
Concealed	22	172
Coal, at road forks, reported at prospect No. 414 on Map II 0' 8" to 1' 0" thick, No. 6 Pocahontas? (base, 3082' L.).....	1	173
Shale, "fire clay".....	1	174
Concealed and sandstone.....	59	233
Shale, weathers light-gray, many plant fossils.....	2	235
Coal, trace	0.0	235
Concealed	4.7	239.7
Coal, No. 4 Pocahontas? (base 3015' B.) (No. 469 on Map II).....	0.3	240
Shale, "fire clay".....	2	242
Shale, sandy, partly concealed.....	10	252
Sandstone, shaly.....	37	289
Shale, sandy.....	5.7	294.7
Coal, No. 3 Pocahontas? (2960' B.) (No. 484 on Map II).....	0.3	295
Shale, "fire clay".....	1	296
Concealed, with shale, yellowish-brown, sandy.....	53	349
Shale, black, No. 2 Pocahontas? (2905' B.).....	1	350
Shale and concealed, yellowish-brown.....	20	370
Mauch Chunk Series—Bluestone Group (330')		
Shale, red, definite, in road in front of church (top, 2890' B.).....	5	375
Shale, red, variegated, and concealed.....	85	460
Sandstone	35	495
Shale, red, and concealed.....	90	585
Concealed.....	115	700
Mauch Chunk Series—Princeton Conglomerate (20' +)		
Sandstone medium-grained	5'	} Princeton Conglomerate 20 720 (top, 2560' B.)
Sandstone, coarse grained, with some pebbles	15 +	
Concealed to Sims Station.....		

Sims Mountain Section—North End.

Meadow Bluff District; starting on the north end of Sims Mountain, 1 mile east-southeast of Rainelle, and measured with aneroid along the road descending the mountain. The measurements are greater than true vertical owing to a dip of about 120 feet. Arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River and Pocahontas Groups (427' +)		
Sandstone, pink, broken, limonite veins.....	30'	} Quinnimont 60 60
Sandstone, white, thin.....	30	

	Feet.	Feet.
Sandstone, fine- to medium-grained, no pebbles seen, Princeton?	7	777
Shale, brown, sandy, and concealed to B. M. 2467' at road forks.....	200	977

Little Sewell Mountain Section—West Side.

Meadow Bluff District; measured with aneroid along the road down the west side of Little Sewell Mountain. The measurements above the Mauch Chunk are somewhat greater than true vertical owing to a dip of about 80 feet as shown by the contours on Map II. Arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River and Pocahontas Groups (355' +)		
Concealed from road forks.....	15	15
Sandstone, gray, medium-grained, zone of carbonized plants 20' from base, thin-bedded at top, more mas- sive at base, but irregular bedding throughout.....	35	50
Concealed, sandy	95.2	145.2
Coal, soft, No. 6 Pocahontas, (2885' B.) (No. 418 on Map II).....	0.8	146
Shale, dark-gray, many fossil plants.....	3	149
Sandstone, white, micaceous.....	4	153
Shale, fissile, iron-stained.....	5	158
Shale, dark-gray, slightly calcareous, fossiliferous.....	3	161
Concealed	54.8	215.8
Coal, No. 4 Pocahontas (2810' B.) (No. 472 on Map II)	0.2	216
Concealed	31	247
Sandstone	2	249
Shale, sandy, many plant fossils.....	2	251
Coal, soft, good, No. 3 Pocahontas, (supplied from opening below road at No. 488 on Map II) (2780' B.)	2	253
Concealed	10	263
Sandstone, thin-bedded.....	15	278
Concealed	49.5	327.5
Coal (2705' B.).....	0.5	328
Shale, chocolate-colored, many fossil rootlets.....	2	330
Sandstone, thin-bedded at top, massive at base.....	25	355
Mauch Chunk Series—Bluestone Group (255')		
Concealed	100	455
Sandstone, much weathered, reddish-brown.....	5	460
Concealed and red shale.....	150	610
Mauch Chunk Series—Princeton Conglomerate (5' +)		
Sandstone, medium-grained, Princeton Conglomerate (2415' B.).....	5	615

The following section, prepared by Ray V. Hennen³, starts at the top of a hill one-half mile west of Russellville, Nuttall District, Fayette County, and extends eastward, with aneroid

Meadow Bluff District; measured with aneroid from the top of the point on the south end of Little Sewell Mountain, traversing southwestward to the county road, thence south to the road forks at 2987' L., thence southeastward to the B. M. 2467'.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River and Pocahontas Groups (369' +)		
Sandstone, (cliff) makes top.....	50	50
Shale, dark, lenticular, 3' to.....	4	54
Coal, Little Fire Creek? (No. 375 on Map II). (3390' B.) fallen shut, thickness reported as.....	2.5	56.5
Shale, fire clay.....	0.5	57
Sandstone, brown, coarse, and coneaeled.....	94	151
Coal, bony..... 0' 3 "	} No. 7 Pocahontas (No. 383 on Map II).... (3290' B.)	3.4 154.4
Coal, block..... 0 3		
Pyrite 0 0¼		
Coal, columnar, laminated with fusain (mineral charecoal).. 0 9		
Coal, hard..... 0 5		
Coal, laminated with fusain (mineral charecoal) and pyrite 1 4		
Coal, bony..... 0 4		
Shale	17.6	172
Coal, No. 6 Pocahontas, (No. 421 on Map II) (3270' B.)	3	175
Coneaeled	80	255
Coal, No. 3 Pocahontas, (No. 491 on Map II) (3190' B.)	1	256
Coneaeled	13	269
Sandstone	35	304
Coal (3140' B.).....	0.8	304.8
Fire clay.....	1.2	306
Shale, brown, sandy.....	8	314
Coal (3130' B.).....	0.3	314.3
Shale, sandy.....	9.7	324
Shale, chocolate-colored.....	1	325
Shale, sandy.....	14	339
Sandstone, shaly.....	8	347
Coneaeled and shale, brown, sandy.....	22	369
Mauch Chunk Series (608' +)		
Shale, red and variegated (top, 3075' B.).....	5	374
Sandstone, shaly, brown to green.....	5	379
Coneaeled	20	399
Sandstone, shaly, green.....	10	409
Shale, red and variegated.....	20	429
Sandstone, shaly, green.....	10	439
Coal and black shale (3005' B.).....	1	440
Shale, red, variegated, and coneaeled.....	100	540
Sandstone, medium- to fine-grained, makes cliff.....	25	565
Shale, sandy, brown.....	20	585
Sandstone, brown, shaly at top.....	25	610
Shale and shaly sandstone.....	80	690
	75	765

measurement, to Meadow Bluff river at Russellville and is continued with the record of the Mrs. E. T. Martin Coal Test Boring—No. 1 on Map II located in Meadow Bluff District, Greenbrier County, just opposite the town. The record of the coal test was kindly furnished the Survey by Samuel Stephenson, of Charleston, West Virginia. In line with recent studies a few minor changes in correlation have been made:

Russellville Section.

	Thickness.		Total.				
	Ft. In.		Ft. In.				
Pottsville Series—New River and Pocahontas Groups (733'±)							
Concealed in gentle slope with small grayish-white boulders from summit of hill.....	65	0	65	0			
Concealed in bench.....	10	0	75	0			
Sandstone, grayish-white.....	20	0	95	0			
Concealed in bench.....	15	0	110	0			
Sandstone, current-bedded, grayish-white, Guyandot	30	0	140	0			
Concealed, mostly sandstone.....	25	0	165	0			
Shale, buff, sandy.....	20	0	185	0			
Coal, Sewell "B", and concealed.....	10	0	195	0			
Concealed	5	0	200	0			
Sandstone, current-bedded, Lower Guyandot.....	25	0	225	0			
Concealed	3	6	228	6			
Coal, Sewell (2045' B.).....	1	6	230	0			
Concealed, steep slope, mostly sandstone.....	45	0	275	0			
Concealed, gentle slope.....	25	0	300	0			
Concealed, steep slope.....	15	0	315	0			
Sandstone, grayish-white, making cliff, Upper Raleigh	45	0	360	0			
Concealed to top of coal test boring.....	5	0	365	0			
(Continued with log of Mrs. E. T. Martin Coal Test Boring—No. 1 on Map II, Elevation top of hole, 1930' B.)							
Surface	10	0	375	0			
Sandstone	10	0	385	0			
Slate, gray.....	57	0	442	0			
Bone	0	4	442	4			
Sandstone, hard, Lower Raleigh.....	26	0	468	4			
Slate, gray.....	65	3	533	7			
Shale, dark, sandy.....	17	10	551	5			
Slate, gray.....	27	5	578	10			
Sandstone and shale..... 4' 7"	} Pineville	49	2	628	0		
Sandstone						40	7
Sandstone, pebbly.....						2	0
Sandstone						2	0
Fire clay.....	3	6	631	6			
Sand, shale.....	10	8	642	2			
Slate, gray.....	12	5	654	7			

Coal, Little Raleigh (2925' B.) (No. 231A on Map II)	1	349
Shale, sandy.....	4	353
Sandstone, gray to brown.....	8	361
Concealed	7	368
Shale, sandy, gray to brown and concealed.....	10	378
Concealed	20	398
Sandstone, brown, shaly.....	10	408
Concealed	5	413
Sandstone, brown, shaly.....	5	418
Shale, brown, sandy.....	10	428
Concealed	35	463
Shale, gray to brown.....	20	483
Sandstone	5	488
Concealed	8	496
Sandstone	22	518
Shale, sandy, gray to brown, and concealed.....	17	535
Fire clay (2735' B.).....	3	538
Shale, sandy.....	10	548
Concealed	27	575
Coal, No. 7 Pocahontas, (2695' B.) (No. 385 on Map II) fallen shnt, visible 2' to.....	3	578
Fire clay.....	2	580
Shale, sandy, variegated, and concealed.....	27	607
Coal, banded bright and dull 1' 8" } No. 6 Pocahontas (2565' B.).....	3	610
Coal, soft, partly columnar 1 4 } (No. 430 on Map II)	15	625
Shale, sandy and concealed.....	5	630
Sandstone	35	665
Concealed	10	675
Sandstone	10	685
Concealed	5	690
Sandstone, brown, fine-grained.....	5	695
Concealed		
Coal 0' 1" } No. 3 Pocahontas (top, 2580' B.).....	2	697
Fire clay 1 9 } (No. 492 on Map II)		
Coal 0 2 }	35	732
Concealed	4	736
Sandstone, shaly, fine-grained.....	1	737
Sandstone, many carbonized plants.....	5	742
Sandstone, medium- to coarse-grained, massive.....	5	747
Sandstone, shaly.....	17	764
Concealed		
Shale, black, No. 1 Pocahontas Coal horizon? (2510' B.) (No. 502 on Map II).....	1	765
Concealed	30	795
Sandstone	30	825

Mauch Chunk Series—Bluestone Group (49' +)

Shale, red, found in J. E. Dorsey water well above road	5	830
Concealed to road fork at Charmco (BM 2401').....	44	874

	Ft.	In.	Ft.	In.
Fire clay.....	1	5	658	10
Sandstone	11	2	670	0
Coal	1	2	671	2
Fire clay.....	2	0	673	2
Sandstone	3	1	676	3
Shale, dark, sandy.....	6	6	682	9
Slate, gray.....	6	7	689	4
Sandstone	4	0	693	4
Shale, dark, sandy.....	5	9	699	1
Fire clay.....	1	10	700	11
Shale, sandy.....	15	4	716	3
Sandstone	9	4	725	7
Slate, black.....	1	1	726	8
Coal, No. 6 Pocahontas.....	2	5	729	1
Fire clay.....	2	0	731	1
Sandstone, to bottom of hole.....	1	11	733	0

Charmco Section.

Meadow Bluff District; starting on a high knob on Laurel Creek Mountain 0.3 mile southwest of Orient Hill Church measured with aneroid and hand-level to the road forks on the divide, then south along the highway to Charmco. Measurements are less than true vertical due to a northwest dip of about 150 feet as shown by the green contours on Map II. Arrangement in descending stratigraphic order.

	Thickness.	Total.
	Feet.	Feet.
Pottsville Series—New River and Pocahontas Groups (825'±)		
Interval from top of knob to the Joe Neff mine on Snowden Crane property.....	200	200
Shale, dark, Hartridge.....	1.9	201.9
Coal, hard, laminated and blocky..... 0' 11½"	} Sewell 6.1 208 (3065' B.) (No. 51 on Map II)	
Shale, with coal streaks	0 4	
Coal, columnar, soft..	1 1	
Mineral charecoal.....	0 0¼	
Coal, laminated, light and dull.....	0 11½	
Coal, laminated, soft	0 4	
Coal, hard.....	0 2	
Shale, reported.....	1 0	
Coal, reported.....	1 3	
Concealed to top of bench.....	45	253
Sandstone, brown to gray, cross-bedded, medium-grained, Upper Raleigh.....	55	308
Shale, sandy and concealed.....	15	323
Coal, Little Raleigh "A" (2950' B.) (No. 231B on Map II).....	0.3	323.3
Fire clay	2.7	326
Concealed	12	338

Big Clear Creek Mountain Section.

Meadow Bluff District; measured with aneroid along the public road descending the east side of the south end of Big Clear Creek Mountain, starting at a point 1.95 miles north of Rupert.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River Group (157' +)		
Concealed from top of knob.....	70 +	70
Coal, Fire Creek, (3273' L.) (No. 318 on Map II).....	2.1	72.1
Concealed	29.9	102
Sandstone	7	109
Coal	0.5	} Little Fire Creek (3230' B.)..... (No. 375A on Map II)
Shale	2.5	
Sandstone conglomerate... ..	2.5	
Coal	0.5	
Sandstone, irregular-bedded, Pineville.....	37	152
Concealed	5	157
Coal blossom, No. 8 Pocahontas (3188' B.) (No. 378A on Map II).....		157

Pottsville Series—Pocahontas Group (284')		
Concealed	35	192
Sandstone, grayish-brown, micaceous.....	12	204
Bone, supplied from mine above road.....	1.3	205.3
Coal, bright.....20" } No. 7 Pocahontas (3135' L.)	3.7	209
Coal, dull.....24 } (Mine No. 400 on Map II)		
Concealed and shale.....	36	245
Shale, black.....	3.5	248.5
Coal and bone, No. 6 Pocahontas (3093' L.) (No. 437 on Map II).....	2.5	251
Concealed and shale.....	14.8	265.8
Coal, No. 5 Pocahontas (3078' B.) (No. 468 on Map II)	0.2	266
Shale, sandy.....	18	284
Coal, No. 4 Pocahontas (3058' B.) (No. 477 on Map II)	2	286
Shale, fire clay.....	1.5	287.5
Concealed and shale.....	22.5	310
Coal, No. 3 Pocahontas (3033' B.) (No. 494 on Map II)	1	311
Shale, fire clay.....	1	312
Concealed and shale.....	114	426
Sandstone, massive, fine-grained, micaceous, zone of carbonized plants, near base.....	10	436
Concealed	5	441

Mauch Chunk Series (521' +)		
Shale, red (top, 2913' B.).....	5	446
Shale, variegated.....	5	451
Concealed	10	461
Sandstone, fine-grained, green, shaly at base.....	10	471
Coal, very impure.....	0.8	471.8
Shale, fire clay.....	1.2	473
Shale, green.....	8	481
	2	483

Quinwood Section.

Meadow Bluff District; starting at the road forks at the western edge of Quinwood and measured with aneroid ascending the mountain westward along the road. The intervals are somewhat less than true vertical measurements due to a northwest dip of about 75 feet as shown by the contours on Map II. Arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River Group (411'+)		
Sandstone, medium-grained, brown, irregular bedding, caps knob, Upper Nuttall (top, 3432' B.).....	20	20
Shale, sandy and conealed.....	25	45
Coal blossom, laeger "B" (No. 1 on Map II).....		45
Conealed	25	70
Conealed in bench.....	20	90
Sandstone, gray, medium-grained, irregular bedding, Lower Nuttall.....	30	120
Shale, gray, sandy, Upper laeger.....	20	140
Conealed	20	160
Coal, slaty, Hughes Ferry (No. 2 on Map II) (top, 3272' B.).....	2	162
Shale, sandy, gray.....	9	171
Conealed	5	176
Sandstone, irregular bedding, Middle laeger.....	19	195
Conealed	4	199
Coal, impure, Lower laeger (No. 4 on Map II) (top, 3233' B.).....	3	202
Sandstone, brown to gray, shaly, Lower laeger.....	28	230
Shale, gray to brown, fissile.....	5	235
Shale, black.....	1	236
Conealed	3	239
Sandstone, fine- to medium-grained, gray to brown, Harvey Conglomerate.....	20	259
Conealed	10	269
Coal	0.5	269.5
Shale, "fire clay".....	1.5	271
Shale, gray to brown, sandy, Sandy Huff.....	52	323
Coal, Castle, (No. 5 on Map II) (top, 3108' B.).....	1	324
Shale, gray, "fire clay".....	3	327
Sandstone, fine-grained, gray to brown, thin-bedded, Guyandot	7	334
Conealed	40	374
Shale, black, Skelt.....	6.5	380.5
Coal, Sewell "B" (No. 6 on Map II).....	0.5	381
Shale, sandy, gray to brown.....	28	409
Coal, at road forks, Sewell "A" (base, 3021' B.).....	2	411
Interval to Sewell Coal estimated....	30	441

In the following section the interval between the No. 8 Pocahontas Coal and the Little Fire Creek Coal is about 30 feet less than true vertical measurement due to a compass

Little Clear Creek Section.

Meadow Bluff District; measured with aneroid, starting at the point where the fire trail leaves the top of Little Clear Creek Mountain and continuing southwestward with the trail to Little Clear Creek. The intervals above the No. 6 Pocahontas Coal are too great, there being a dip of about 75 feet as shown by the contours on Map II. The intervals below the coal represent nearly true vertical measurements. Arrangement in descending stratigraphic order.

	Thickness. Feet.	Total. Feet.
Pottsville Series—New River and Pocahontas Groups (420'+)		
Sandstone medium-grained, gray, irregular bedding, (top, 3400' B.) Pineville.....	50	50
Concealed	70	120
Sandstone, fine-grained.....	5	125
Concealed	50	175
Coal blossom, No. 6 Pocahontas? (3225' B.) (No. 461 on Map II).....		175
Concealed	5	180
Sandstone, brown, fine-grained.....	5	185
Concealed	75	260
Sandstone	8	268
Concealed	6	274
Sandstone, coarse-grained, gray to brown.....	2	276
Coal, No. 3 Pocahontas (top, 3125' B.) (Prospect No. 488 on Map II).....	4+	280
Shale, "fire clay," numerous fossil rootlets.....	2—	282
Shale, sandy, numerous fossil rootlets.....	4	286
Sandstone, fine-grained, massive at top, thin-bedded at base.....	19	305
Concealed	70	375
Sandstone, argillaceous.....	5	380
Concealed.....	5	385
Black slate, traces, with "fire clay".....		385
Concealed	35	420
Mauch Chunk Series (480'+)		
Concealed to Little Clear Creek.....	480	900

The following record of a boring 1 mile south of Duo is included in this Chapter because of its prime stratigraphic importance:

Raine Lumber and Coal Company Coal Test Boring No. 5— No. 6 on Map II

Meadow Bluff District; on Shellcamp Ridge, one mile south of Duo; elevation, 3630' L.

	Thickness. Ft. In.	Total Ft. In.
Pottsville Series—New River Group (537'+)		
Surface	12 6	12 6

Knob, crosses Beech Ridge one mile northwest of Clearco, passes just west of Duo and follows the west side of Big Clear Creek to disappear on the south end of Pollock Mountain, about one mile north of Anjean.

The Kovan Syncline is a very shallow fold with the elevation of the key bed along its axis rarely 100 feet lower than it is along the axis of the Webster Springs Anticline. Along the axis of the syncline at the county line the Sewell Coal has an elevation of a little less than 3100 feet, gradually rises to an elevation of about 3710 feet $1\frac{1}{2}$ miles south of Mann Knob. From this point southwestward along the axis the elevation declines to a low point just west of Duo where the Sewell Coal is about 3420 feet. From Duo to the south end of Pollock Mountain, where the syncline disappears, the elevation of the Sewell Coal rises about 40 feet.

The outcropping rocks along the syncline are mostly the New River Group of the Pottsville Series with the Mauch Chunk at the surface along the valleys of North and South Forks of Cherry River, and on Big and Little Laurel Creeks.

Boggs Knob Anticline.—The Boggs Knob Anticline of Hennen³ received its name from a knob of the same name in western Greenbrier County. It has been traced west and south from that point to its southern termination in Summers County, three miles northwest of Hinton. It is a very shallow fold with a reversal of less than 100 feet. The close similarity of this fold to the Webster Springs Anticline, described above, led to an attempt to prove that both were part of the same anticline. All field evidence refutes such an idea and as shown on Map II the Boggs Knob Anticline disappears near the southern end of Goddard Mountain.

The surface rocks along the two miles of the anticline in Greenbrier County belong to the New River and Pocahontas Groups of the Pottsville Series and to the Bluestone and Princeton Groups of the Mauch Chunk Series.

point along the crest of the fold the elevation of the coal again rises and at the high point on north Pollock Mountain the elevation of the Sewell Coal is slightly over 3550 feet. South from this point the fold pitches at the rate of about 50 feet to the mile. The antiline and the Kovan Syncline come together and disappear about one mile north of Anjean. As indicated there is a dome with a closure of approximately 100 feet between Anjean and the headwaters of Sam Creek.

Correspondence with Mr. W. W. Coleman, Chief Engineer of the Leckie Smokeless Coal Company, indicates that the elevations for mine openings 92, 93, 94, 95, and 96, as used in making the structure map are each 17.57 feet too high. The net result of this error is to shift the closed 3500-contour northeast until it passes between mines 92 and 93 instead of between 93 and 94. The 3450-contour should be moved east with a rather sharp bend, passing between mines 94 and 95 and back between 95 and 96. The other contours are not materially affected.

The Webster Springs Antiline has a length, in Greenbrier County, of 18 miles and throughout its length it is asymmetrical, the dip being greater on the west side than on the east. The surface rocks along the crest of the antiline in Greenbrier County are mostly the New River Group of the Pottsville Series with the Bluestone, Princeton, and Hinton Groups of the Mauch Chunk Series coming to the surface along the North Fork of Cherry, South Fork of Cherry, Little Laurel Creek, and Big Laurel Creek.

Kovan Syncline.—The Kovan Syncline of Reger² roughly parallels the Webster Springs Antiline and has been traced from its northern end, near Hodam, Webster County, to the Greenbrier County line $\frac{1}{2}$ mile east of the common corner of Webster, Nicholas, and Greenbrier Counties. The axis of the syncline crosses the North Fork of Cherry River between Coats Run and Little Lick Run, turns a little more south to the mouth of Beech Lick Run on the South Fork of Cherry River, follows along the river to the mouth of Mill Run, turns more to the

Springdale Syncline.—The Springdale Syncline of Alderson¹ roughly parallels the Boggs Knob Anticline and the surface rocks belong in the Pocahontas Group of the Pottsville Series and in the Bluestone Group of the Mauch Chunk Series. With a length of less than two miles in the county the syncline dies out just south of Goddard Mountain. The exact location of the axis of the syncline is difficult to find but its probable location is shown on Map II, being about two miles southeast of the crest of the Boggs Knob Anticline.

Alderson Anticline.—The Alderson Anticline of Reger² has been traced from Summers County, across the western corner of Monroe County, to the city of Alderson at the Greenbrier County line. Extending through Alderson, from which the fold derives its name, the anticline has been traced to its northern end near Muddy Creek Church. Throughout the four miles of the fold in Greenbrier County the surface rocks are the limestones of the Greenbrier Series.

Creamery Syncline.—The Creamery Syncline of Reger², roughly parallels the Alderson Anticline. Starting in Summers County, one mile southeast of Bangers Springs it extends northeastward into Monroe County, passes just east of the village of Creamery, from which it is named, and reaches the Greenbrier County line about $\frac{1}{2}$ mile east of Alderson. From the county line it extends northeastward to Blaker Mills and disappears about $1\frac{1}{2}$ miles north of that village.

The surface rocks along the axis of the syncline in Greenbrier County belong to the basal part of the Mauch Chunk Series and the upper part of the Greenbrier Series.

Williamsburg (Mount Pleasant) Anticline.—The Mount Pleasant Anticline of Reger² has been described as a weak fold starting $1\frac{1}{2}$ miles northeast of Wolf Creek Post-Office, extending northward and passing just east of Mt. Pleasant School, reaching the Greenbrier County line $1\frac{1}{2}$ miles east of Alderson,

¹Op. cit., p. 95.

²Reger, David B., Mercer, Monroe, and Summers Counties, W. Va. Geol. Surv. p. 149, 1926.

its total length in Monroe County being 4 miles. The Williamsburg Anticline of this report, although apparently connecting with the Mount Pleasant Anticline, has been renamed because of its much greater magnitude in Greenbrier County.

The northern end of the fold is about one mile east of Trout Post-Office, and from that point the axis has been traced in a general southwest direction, passing $\frac{1}{2}$ mile west of Sunlight and is located about 0.7 mile east of the town of Williamsburg, from which the fold was named. Continuing southwestward the axis is on the crest of Brushy Ridge, passing through Alta and Brushy Ridge School. Near Asbury there is an offset along the axis to the east; the crest line, as shown by the dashed, red line on Map II, crosses a low saddle in a southeast direction for a distance of about one mile. Resuming its southwest course the fold passes through the south end of Muddy Creek Mountain, the axis passing midway between Hawver School and Fearnster School and reaching the Greenbrier River one mile east of Alderson. The total length in Greenbrier as described is 23 miles.

The fold is unusual in that it is quite severe yet very narrow and that the dip is more rapid on the east side than on the west. At Alta the crest of the anticline is structurally more than 1000 feet higher than the area 0.7 mile to the east and an equal amount above the area 1.4 miles to the west; indicating both the sharpness of the reversal and the steeper east limb. The rocks along the crest of the anticline between Brushy Ridge School and about one mile southeast of Williamsburg are nearly horizontal, with the fold pitching to the north and south from these points. The northern end of the anticline plunges more rapidly than the southern end.

The surface rocks along the crest of the anticline are all Mississippian in age, belonging to the lower part of the Manch Chunk Series, the Greenbrier Series, the Maecrady Series, and the Pocono Series. From the northern end, east of Trout Post-Office, to a point 1.2 miles northeast of Williamsburg, the entire thickness of the Greenbrier Series is at the surface. From this point to a point 0.3 mile northwest of Asbury the out-

Continuing southwest the axis makes a gentle curve around the city limits of Roneeverte, crosses the Greenbrier River slightly less than one mile east of Rockland and reaches the Monroe line midway between Hokes Mill and Nickells Mill. From this point the fold has been traced into Monroe County, passing $\frac{1}{3}$ mile west of Sinks Grove, from which the fold was named, to a point 1.1 miles east of Lillydale where it disappears.

The surface rocks along the axis of the fold are almost entirely of the Maerady Series. At a few points the basal beds of the Greenbrier Series may remain on the crest and where streams cut across the antiline the upper members of the Pocono Series are exposed.

Caldwell (Patton) Syncline.—The Patton Syncline of Reger¹⁰ has been described in Monroe County as a weak structural feature starting $1\frac{1}{2}$ miles south of Sinks Grove, extending northeastward for six miles to the Monroe-Greenbrier County line $1\frac{1}{2}$ miles east of Patton. Because of its much greater extent and severity in Greenbrier County the fold has been renamed the Caldwell Syncline, from the town of the same name through which it passes and where it is a prominent structural feature.

From its northern end $1\frac{1}{2}$ miles north of Anthony, the axis of the syncline extends southwestward, in the general direction of the Greenbrier River, passes $\frac{1}{2}$ mile west of Anthony, through Camp Longemount, passes just west of Harpers, through Camp Alleghany to Caldwell. From Caldwell the axis of the fold continues southwest through Holliday School, crosses U. S. Route 219 $\frac{1}{4}$ mile west of Organ Cave, and reaches Second Creek and the Monroe County line one mile east of Patton.

The surface rocks along the syncline north of Caldwell are wholly of the Pocono Series except for a few small areas of Maerady and Greenbrier rocks at and near Caldwell. Southwest of Caldwell the surface rocks along the axis are entirely of the Greenbrier Series except for a very small area of Maerady that is at the surface two miles southeast of Roneeverte.

around Asbury the basal members of the Greenbrier Limestone are at the surface and about $\frac{1}{2}$ mile south of Asbury the Maerady and Pocono again appear. Continuing south along the axis the entire thickness of the Greenbrier Series dips below the surface and on the south end of Muddy Creek Mountain, rocks of the Bluefield Group of the Mauch Chunk Series form the crest of the antiline. The upper part of the Greenbrier Series is again exposed in the Greenbrier River gorge.

Muddy Creek Mountain Syncline.—Muddy Creek Mountain Syncline is a broad structure with the west limb much steeper than the east limb. In many places the exact position of the axis of the fold is very difficult to find but its general location is clearly defined.

As shown on Map II the fold has been traced from its northern end, 1 mile north of Mt. Vernon School, extending in a general southwest direction to Frazier, just south of the Greenbrier River. Describing the fold in more detail: the axis passes $\frac{1}{4}$ mile east of Unus, follows Burus Run for a short distance, crosses U. S. Route 60 about $1\frac{1}{2}$ miles west of Richlands, passes near Persinger School and follows the west side of Muddy Creek Mountain to Fry School, reaching the Greenbrier River just west of Frazier. It is possible that this fold is the northward continuation of the Laurel Creek Syncline of Reger⁸.

The surface geology along the axis of Muddy Creek Mountain Syncline is mainly that of the Bluefield Group of the Mauch Chunk Series but around Unus, on Spice and Burns Runs, on the headwaters of Milligan Creek and along the Greenbrier River there are outcrops of the Greenbrier Limestone.

Sinks Grove Antiline.—The Sinks Grove Antiline of Reger⁹ is a prominent fold crossing most of Monroe and Greenbrier Counties. Having its northern end at Gardner, the axis of the antiline extends in a southwest direction through the villages of Henning and Vago, passes $1\frac{1}{2}$ miles east of Maxwelton and passes just east of Lewisburg through Wagner Hill.






PLATE VI.—Outcrop of Pickaway Limestone and shale in Monroe County on the east or right limb of a symmetrical fold. Note the vertical fracture cleavage developed in the shale. Note also that the joints in the overlying limestone are normal to the bedding. This outcrop apparently proves that the formation of the joints in the limestone, described as "Pickaway Limestone" in Chapter VII, was independent of and occurred prior to the major folding of the Appalachians.



PLATE VIII.—Drag folding in interbedded limestones and shales of the Rondout Group, 0.5 mile west of Alvy



and except for the rocks at the Bobs Ridge the surface rocks along the axis belong in the Oriskany Series. At White Sulphur Springs the surface rocks along the crest belong to the Mareellus Series and southeast the fold continues to pitch with Upper Devonian rocks along the crest of the structure, the rocks at the county line belonging to the Chemung Series.

For details of the structure of this antiform the reader is referred to the cross-sections on the margin of Map II (in Atlas) and to the discussion of faults at the end of this Chapter.

Stony River Syncline.—The Stony River Syncline of Darton and Taff¹⁶ originates along the North Branch of the Potomac River in Mineral County and has been traced southwestward across Grant, Tucker, Randolph, and Pendleton Counties, passing into Highland County, Virginia, two miles east of the common corner of Pendleton, Pocahontas, and Highland. Remaining in Virginia for nine miles the axis of the syncline enters Pocahontas County where the Staunton and Parkersburg Pike crosses the State line, 2.2 miles east of Top of Allegheny. From this locality it continues southwestward and follows, in general, the State line to Laurel Creek, where the main axis is found 1 mile west of Rimel. The fold enters Greenbrier County at Middle Mountain and coincides with this mountain to its southern end one mile northwest of Neola. The axis of the main basin crosses Anthony Creek $\frac{1}{4}$ mile east of Bound School and turning about due south the fold loses its identity on Whitmans Draft four miles south of Alvon.

In Greenbrier County this structural basin is a broad gentle syncline, much complicated by crumpling of the relatively incompetent shales and sandstones of the Upper Devonian. The surface rocks along its axis belong exclusively to the Chemung and Portage Series.

Neola Anticline.—The Neola Anticline, not previously named or described, roughly parallels the Stony River Syncline. Originating $4\frac{1}{2}$ miles south of Alvon the axis extends

cline deepens to the southwest and in general the rise is more rapid on the east side than on the west.

Maple Grove Anticline.—The Maple Grove Anticline of Reger¹¹ named from Maple Grove School, Greenbrier County, is a poorly defined structure with a total length of 10½ miles, six miles being in Monroe County and 4½ miles in Greenbrier County. Starting 1½ miles west of Pickaway it parallels the Caldwell (Pattou) Syncline, entering Greenbrier County ¾ mile southwest of Maple Grove School. Passing ½ mile east of Organ Cave the fold merges into a terrace 1 mile northeast of Forestdale School.

Northeast of the county line the surface rocks along the crest belong to the Greenhrier, Maccrady, and Pocouo Series, appearing in the order named.

Hurricane Ridge Syncline.—The Hurricane Ridge Syncline of Reger¹², described by him as originating in southwest Virginia, has been traced across Mercer and Monroe Counties to the Greenbrier County line ½ mile east of Maple Grove School. The fold has a length of only 3½ miles in Greenbrier County, merging into a terrace 1½ miles northeast of Forestdale School. Northeast of the county line the surface rocks along the axis of the syncline belong in the Greenhrier, Maccrady, and Pocouo Series, appearing in the order named.

Browns Mountain Anticline.—The Browns Mountain Anticline of Darton¹³, described in Pocahontas County in more detail by Price¹⁴, is the same as the Harts Run Anticline of Reger¹⁵. As noted in the Pocahontas County report cited above, the structure is that of an anticlinorium, overturned to the west and it is now known to be faulted along the central west side.

¹¹Op. cit., p. 153.

¹²Op. cit., pp. 146-9.

¹³Darton, N. H., Monterey Folio, No. 61, U. S. Geol. Sur.; p. 6, 1898.

¹⁴Price, Paul H., Pocahontas County, W. Va. Geol. Sur.; pp. 80-1, 1929.

Draft School. From this locality the axis extends northward to a point one mile east of Bound School where it again resumes its northeast course. From this point the main axis follows the western side of Anthony Creek, passing through the western edge of the town of Neola and leaving Greenbrier County 1.3 miles northeast of Trainer. It is probable that the Neola Anticline connects with the unnamed anticline at Rimel in Pocahontas County.

The exact location of the axis of this anticline, like that of the syncline to the west, is difficult to determine due to the crumpling of the rocks. It is not unusual to find six or more reversals of dip in a distance of half a mile across the strike of the rocks. Dips of 80 degrees are common and locally the beds may be overturned. The rocks along the main axis belong to the Portage Series.

Meadow Creek Syncline.—The Meadow Creek Syncline, not previously named or described, is a well-defined basin in eastern Greenbrier County. The axis nearly coincides with Meadow Creek, from which it was named, and with Laurel Run. Paralleling the State line the total length of the syncline is probably not much greater than the 15 miles present in Greenbrier County.

The surface rocks along the axis belong to the Pocono Series.

Kates Mountain Syncline.—The northern end of the Kates Mountain Syncline of Reger¹⁷ is about $\frac{1}{2}$ mile east of Pleasant Valley School. The axis extends in a southwest direction passing along the length of Kates Mountain and leaves Greenbrier at the southern end of Kates Mountain. The syncline has been traced nine miles into Monroe County, terminating $1\frac{1}{2}$ miles northeast of Red Mill in that county. Its length in Greenbrier County is eight miles.

The surface rocks along the axis are confined to the Chemung Series with the basal beds of the Pocono Series being retained on Kates Mountain.

south of the Pocahontas County line and extending across Beaver Lick Mountain to the North Fork of Anthony Creek. The section was drawn to illustrate the Burr Fault. Here the red sandstones and sandy shales of the Red Medina are lying on the overturned Marellus black shales. The fault's projection below the surface is hypothetical but is believed to be as shown in the cross-section.

Cross-Section B—B'.—Cross-Section B—B' begins on Cold Knob on Cold Knob Mountain, extends along Chestnut Ridge, through Falling Springs (Renick P. O.), through the Anthony Creek gorge at Alvon and ends at the State line $\frac{1}{2}$ mile north of Smith Knob. The surface rocks along the section range from the Pottsville Series down to the Clinton Series. The total length of the section is 22 miles.

Cross-Section C—C'.—This is a short section extending from Greenbrier Mountain through the southern tip of Coles Mountain, through Bobs Ridge and ending on Sulphur Lick Run. The surface rocks are entirely Devonian with all of the series represented. The section was drawn at this point to illustrate the complex antiform.

Cross-Section—D—D'.—This 22-mile long section crosses about two-thirds of the county. Starting at Clintonville it extends southeastward, passes just south of Alta, through Lewisburg and Caldwell, and ends at the State line two miles northeast of the common corner of Greenbrier, Monroe, and Alleghany Counties. The surface rocks include the Bluefield Group of the Mauch Chunk Series, the Greenbrier, Maccrady, and Pocahontas Series of the Mississippian, and the Chemung and Portage Series of the Devonian.

UNCONFORMITIES.

All of the regional unconformities noted in Greenbrier County belong to the type known as disconformity, i.e., the beds above and below the surface of erosion are approximately parallel. As a result they are of minor importance from a structural standpoint and only a brief discussion of each un-

Glace Anticline.—The Glace Anticline or Rege^r has a length of five miles in Monroe County, starting $\frac{1}{2}$ mile northeast of Elk Knob in that county and entering Greenbrier County one-half mile northward from Glace. Northeastward the axis passes one mile west of Upper Tuckahoe School, crosses Dry Creek 0.6 mile northwest of Lower Tuckahoe School, and terminates 0.6 mile east of Pleasant Valley School. The length of the fold in Greenbrier County is $8\frac{1}{2}$ miles and the surface rocks along its axis belong to the Chemung Series.

Tuckahoe Syncline.—The Tuckahoe Syncline, not previously named or described, is a small but sharp down warp with a total length of $9\frac{1}{2}$ miles. Starting on Brushy Mountain the axis of the syncline has been traced southwestward, crossing U. S. Route 60 about $1\frac{1}{2}$ miles east of Pleasant Valley School and about two miles west of the Virginia State line. From this point the axis crosses O'Neill Knob, passes 0.2 mile west of the village of Tuckahoe, from which it receives its name, and follows the west side of Dry Creek to Upper Tuckahoe School. From this locality the axis continues southwestward and terminates on Grindstone Ridge near the Monroe County line. The surface rocks along the axis belong in the Chemung Series with a small area of Pocono rocks on O'Neill Knob.

CROSS-SECTIONS.

In central and eastern Greenbrier County the rocks are often standing at steep dips and in some cases are slightly overturned or otherwise so disturbed that structure contouring is not possible. In this area the contours are replaced by dip and strike symbols and in addition four cross-sections have been prepared to show in graphic manner the position of the various beds. All of these cross-sections have been made on a vertical and horizontal scale of 1:62,500, or 5208 feet to the inch, which is the same scale as the topographic map. Each of them extends approximately at right angles to the strike of the rocks and are so spaced as to illustrate the most interesting features.

Cross-Section A—A'.—Cross-Section A—A' is 2.6 miles long beginning on the headwaters of Little Creek, one mile

the unconformities the reader is referred to the discussions of the contacts, given in the Chapters on Stratigraphy of the various series. (See Index for page references).

The uppermost important time break in the geologic column is at the contact of the Pottsville Series of the Pennsylvanian with the underlying Mauch Chunk Series of the Mississippian. The contact is that of an overlap of transgression, with younger and younger beds of the Pottsville resting on the Mauch Chunk. As is the case with all unconformities of this type, the lapse of time between the deposition of the underlying and the overlying beds varies in the direction of the overlap and in this case the interval becomes greater in a north and northwest direction.

The next lower regional unconformity is at the contact of the Greenbrier Series with the underlying Maerady Series. The contact between the massive limestone and the Maerady red shales is usually sharp but occasionally a thin calcareous shale is present, giving to the contact a blended appearance. The apparent absence of beds representing the Warsaw and Spergen Formations of the Mississippi Valley suggests the time value of the unconformity.¹⁹

Another unconformity is found at the contact of the Pocono Series with the Chemung Series.¹⁹

The contact between the Helderberg Series of the Devonian and the Bossardville Series of the Silurian has been reported as unconformable, in reports on near-by areas. In Greenbrier County there is insufficient evidence to determine the exact relationship of the two beds but the relationship is tentatively considered to be that of a disconformity.

FAULTS.

Only one major fault was noted in the county, that being along the west side of Beaver Lick Mountain near the Pocahontas County line. Several small faults were noted but only four of these are worthy of mention.

¹⁹The full time value of the unconformities at the base of the Greenbrier Series and at the base of the Pocono Series can not be determined until paleontologists agree upon the time range of the various fossils

sharp-edged, the rock is usually called, a breccia, not a conglomerate. The cementing material of either conglomerate or breccia is usually calcium carbonate or ferric oxide.

Sandstone is composed essentially of grains of quartz sand. Most sandstones contain smaller quantities of several other minerals such as magnetite (magnetic iron ore) and mica. Sandstone is described as coarse, medium, or fine grained, according to the prevailing size of the sand grains of which it is composed. The varying colors of sandstones are due to the cementing materials and to minor constituents, since pure quartz sand is white or transparent.

Shales are composed of compacted, finely divided sediment, and usually contain a high proportion of clay. Unlike sandstones and conglomerates, they do not require the presence of cementing material. They are the softest of ordinary sedimentary rocks, and disintegrate more rapidly through weathering than any of the others. Some shales are popularly known as "slate," especially in the coal mining districts. True slate, though formed from shale, is quite different and results from more intense pressure and heat.

Limestone consists essentially of calcium carbonate. In addition, however, all limestones contain varying, though frequently small, proportions of other minerals. They are harder than shales and, when well compacted, are among the toughest and strongest of sedimentary rocks. As calcium carbonate is somewhat soluble in water, especially if the latter contains a trace of any acid, limestone is removed directly by running water, without previous weathering. This process of removal of limestone by **solution**, when carried on by underground water, results in the production of the caves and sinks that are so common in thick limestones. A limestone-like rock, which contains, besides calcium carbonate, a considerable percentage of magnesium carbonate, is called a **dolomite**. With a smaller percentage of magnesium carbonate, it is called a dolomitic limestone.

Coal is a term applied to vegetable matter with varying

portions of animal matter which through geological processes has become so changed by loss of volatile matter that it is more or less compact and dark in color. It burns with comparative slowness and decomposes slightly in the atmosphere. It has a variable chemical composition and is not homogeneous. It grades into peat and differs from that substance in composition chiefly in the smaller percentage of water, oxygen, and volatile hydrocarbons.

A few descriptive terms that will be used frequently in the volume will be defined here:

Arenaceous, from Latin *arena*—sand; meaning sandy, or composed largely of sand.

Argillaceous, from Latin *argilla*—white clay; meaning composed largely of clay.

Calcareous from Latin *calx*—limestone; meaning composed largely of calcium carbonate.

Sedimentary rocks, though often occurring as described above, are probably found more often of intermediate composition. Thus a rock may be formed of a mixture of the finely divided particles of which shale is composed, with calcium carbonate. If the latter appear to predominate, the rock is called an **argillaceous limestone**. In the same way, a rock composed of a mixture of sand and calcium carbonate is an **arenaceous limestone** if the main constituent is calcium carbonate; but if it is composed mainly of sand grains it is called a **calcareous sandstone**. So too, a rock made up of shale particles and sand grains is an **arenaceous shale**, or an **argillaceous sandstone**, depending on which constituent predominates.

Derivation of Sediments and Implied Environment.—As stated above, all of the outcropping rocks of Greenbrier County are of sedimentary origin. They consist of sandstone, shale, and limestone of great variety in composition and appearance. These materials were originally gravel, sand, and mud, derived from the decomposition of older rocks, chemical precipitates, and the remains of plants and animals that lived in the seas or swamps while the strata were being deposited.

The rocks reveal the unwritten history of the sedimentation from the early Silurian to middle Pottsville time. From

can be interpreted as to the conditions under which they deposited. For example, rocks marked by ripples, cross-bedded by currents, or cracked by drying on mud-flats, indicate shallow water, while certain fossils indicate marine water and others indicate fresh or brackish water.

Not only can the condition of sedimentation be determined but also the character of the adjacent land. The sand and pebbles of coarse sandstone and conglomerate show that the adjoining land may have been high and the stream gradient steep. Red beds are generally indicative of continental deposits in an arid climate. Limestones are indicative of clear water and if shallow water is also indicated the adjacent land must be low and the streams too sluggish to carry off the coarser sediments.

If we could reproduce the physical environment found at the beginning of the deposition of our sedimentary rocks, which is roughly estimated at 500,000,000 years ago, we would find that the area now occupied by West Virginia was covered with a sea which extended from the Gulf of Mexico on the south to Newfoundland on the north. To the east was a rugged and mountainous continent composed of crystalline (igneous and metamorphic) rocks. This continent roughly paralleled what is now the Atlantic coast. It was from this region that the greater part of the sedimentary rocks now found in West Virginia was derived. The area occupied by this sea was a zone of weakness and was, on the whole, a subsiding basin, in part due to the weight of the accumulating sediments, up to the close of the deposition of the youngest sediments found in the State. During this time minor oscillations caused the withdrawal of marine waters, at times more or less completely. On the whole, however, the area was one of subsidence so that during its history sediments several miles in thickness were accumulated. Generally speaking the water was comparatively shallow and not comparable to our present ocean depths.

The oldest rocks exposed in Greenbrier County are of the Red Medina Series. These rocks outcrop along the west side of Beaver Lick Mountain from the Pocahontas County line

iferous and, as emphasized by their red color, indicate deposition under subaerial conditions. The overlying White Medina (Clineh, Tuscarora) is a dense quartzite, in Greenbrier County, but from its appearance in other counties of the State it is believed to have been deposited in marine waters. The Clinton is poorly exposed in this county but thin limestones in the upper part and scattered fossils indicate that it is at least partly of marine origin. The remaining Silurian beds,—Niagara, Rondout, and Bossardville,—reveal a vast assemblage of marine forms.

It is apparent that Silurian time was one of enervating seas and that during this period the cycle of erosion of the ancient land mass to the east was nearly completed.

The Lower Devonian, next above the Silurian, is abundantly fossiliferous and the environment was quite similar to that prevailing in upper Silurian. The limestones and cherts indicate clear water while the sandstones that occur in the upper part are well sorted and usually quite pure. These sandstones were derived from the east and indicate that the ancient continent was slowly being uplifted.

The Middle Devonian in Greenbrier County is largely black shale. The origin of black shale is still the subject of much debate. However in this county fossils show that marine conditions prevailed for at least a part of Middle Devonian time.

From the bottom to the top of the Upper Devonian the sediments become more and more coarse and the sandstones become more and more massive. The older part (Portage) is only sparingly fossiliferous with both marine and plant fossils. The Chemung coming above the Portage is abundantly fossiliferous with a large assemblage of marine forms. At the top of the Upper Devonian (Catskill) is a succession of red shales with enclosed conglomerates that sometimes reveal plant fossils. These red shales are continental deposits and do not extend over the entire county.

As indicated above, at the start of Devonian time the ancient land to the east was low and the sea clear. Uplift of

predominance of shale. That the eastern land mass continued to rise during Devonian time is shown by the material composing each succeeding group of rocks. From the beginning to the end of this period there is a more or less gradual change from limestone to coarse sandstones and shales, from wholly marine beds to interbedded marine and non-marine beds with red non-marine beds at the top. The direction of the source of the sediments throughout all of Devonian time appears to have been to the northeast of Greenbrier County.

Overlying the Catskill is about 600 feet of sandstones and sandy shales of Mississippian age that are partly of marine and partly of non-marine origin. These beds correlate with the Pocono Series and appear to be the equivalent of the Price Formation of Virginia. The red Macerady shales and thin sandstones are next above and as both the Pocono and Macerady thicken to the southeast the source of the material composing them is assumed to lie in the same direction. Thin lenticular coals in the Pocono indicate a moderate climate.

The source of the detrital material in the Greenbrier limestones is not known but the abundance of marine shells and corals speak eloquently of quiet marine environment and moderate temperatures. Likewise the exact source of the elastic material in the overlying Mauch Chunk Series has not been worked out. However, it is safe to say that the ultimate source of most of the material was the land mass to the east. The Mauch Chunk is composed of red shales and sandstones and some marine limestones with the marine beds occurring less often near the top. Thin coal seams scattered through the middle of this series indicate a generally mild temperature.

The Pottsville Series rests unconformably on the Mauch Chunk with the change quite abrupt from red shales to dark sandy shales and sandstones. Only the lower and middle groups of the Pottsville remain in Greenbrier County, the upper group and all younger rocks of the Paleozoic having been removed by erosion. No distinctly marine fossils have been found in the Pottsville of this area and numerous coal beds testify to a subaerial environment with abundant plant

- Butts, Charles.—Geologic Map of the Appalachian Valley of Virginia with Explanatory Text, Bull. 42, Va. Geol. Sur.; 1933.
- Reeves, Frank.—Manganese Deposits of Eastern West Virginia, Ser. 1, Bull. 6, W. Va. Geol. Sur.; 1935.
- Price, Paul H., McCue, J. B., and Hoskins, Homer A.—Springs of West Virginia, Vol. VI, W. Va. Geol. Sur.; 1936.
- Tucker, R. C.—Deep Well Records, Vol. VII, W. Va. Geol. Sur.; 1936.
- Price, Paul H., Tucker, R. C. and Haught, O. L.—Geology and Natural Resources of West Virginia, Vol. X, W. Va. Geol. Sur.; 1937.

Nomenclature and Correlation.—In Greenbrier County, the problem of proper nomenclature, along with accurate correlations, involves a selection from equivalent titles that have been given the same formations in different regions. In the present instance this discrimination must be made from the published columns and generally accepted terms in the respective localities of the surveys that have been made in adjoining areas, and in part the local area. These are principally the State Surveys of New York, Pennsylvania, and subsequent eastern States; those of Virginia and other southern Appalachian States; those of the general Mississippi Valley; the U. S. Geological Survey; and more especially the column of the West Virginia Geological Survey. Fortunately, general revision is unnecessary, but early deductions must be affirmed, while many of the local names must be considered as such, so that original titles of which there is no longer a doubt may be properly applied.

In this report as in all the West Virginia Geological Survey reports an attempt is made to recognize and follow the earliest nomenclature of authentic publications that have had general circulation and are of sufficient detail to follow.

In the Pennsylvanian Period the amplified Pottsville nomenclature of southern West Virginia, as used in numerous reports of the West Virginia Geological Survey, is employed.

In the Mississippian Period it is necessary to choose between the distinct nomenclatures of the East and the West. In this Period four major series are easily recognized. Particularly is this true in southern West Virginia, where, until the last few years, detailed study with subdivisions of the

rapidly to the north and northwest, due south to the... members from the base and thinning between coal seams. Historical geologists, noting the rapid thickening of Pottsville rocks to the south have ignored much evidence to the contrary and postulated a source for the material in that direction. It is the junior author's belief that the bulk of the material making up these rocks came from some point to the northeast of Greenbrier County and that the rate of subsidence of the filling basin controlled the thickness of the formations. Erosion of the Mauch Chunk shales from the central part of the State may have contributed some detritus, especially in the lower part of the series. This conclusion is based on a study of the unconformity and on the size and distribution of the pebbles, sand, etc., across the northwest part of the county.

Bibliography.—The following is a brief list giving only the principal publications of a geological, historical, or economic nature that have direct bearing on Greenbrier County:

- White, I. C.—Coal Report, Vol. II, W. Va. Geol. Sur.; 1903.
White, I. C.—Petroleum and Natural Gas, Vol. I(a), W. Va. Geol. Sur.; 1904.
Grimsley, G. P.—Clays, Limestones, Cements, Vol. III, W. Va. Geol. Sur.; 1905.
White, I. C.—Supplementary Coal Report, Vol. II(A), W. Va. Geol. Sur.; 1908.
Grimsley, G. P.—Iron Ores, Salt, Sandstones, Vol. IV, W. Va. Geol. Sur.; 1909.
Brooks, A. B.—Forestry and Wood Industries, Vol. V, W. Va. Geol. Sur.; 1911.
Millspaugh, C. F., and White, David.—The Living Flora of West Virginia, Part I, (by Millspaugh); The Fossil Flora of West Virginia, Part II, (by White); Vol. V(A), W. Va. Geol. Sur.; 1913.
Hennen, Ray V.—Fayette County, W. Va. Geol. Sur.; 1919.
Reger, David B.—Webster County, W. Va. Geol. Sur.; 1920.
Reger, David B.—Nicholas County, W. Va. Geol. Sur.; 1921.
Stose, G. W., and Miser, H. D.—Manganese Deposits of Western Virginia, Bull. No. XXIII, Va. Geol. Sur.; 1922.
Wright, Frank J.—The Physiography of the Upper James River Basin in Virginia, Bull. No. XI, Va. Geol. Sur.; 1925.
Reger, David B.—Mereer, Monroe, and Summers Counties, W. Va. Geol. Sur.; 1926.
Price, Paul H.—Poeahontas County, W. Va. Geol. Sur.; 1929.

Summers County Report, however, Reger's careful and detailed study established the relationship of these rocks to those of the East and the West. Because of the close proximity and similarity of conditions the same nomenclature is herein retained so far as applicable. It is true that even in this short distance considerable thinning has occurred but the same major groups have carried through the entire county.

In the Devonian and Silurian it has been the policy of the West Virginia Geological Survey to retain the New York nomenclature where possible. Many of the important subdivisions, although somewhat attenuated, are easily recognized across the State.

In Chapters VI to IX, inclusive, where the various subdivisions are discussed in detail, the nomenclature of several organizations and authors is included, together with that adopted in this report, which should serve to harmonize conflicting names.

Classification of Outcropping Rocks.—Figure 7 is a general columnar section of the outcropping rocks of Greenbrier County, indicating the maximum and minimum thicknesses of all subdivisions of sufficient importance to be mapped geologically, followed by a brief description of their most salient features. Further descriptions and subdivisions are included under the discussions of each series in Chapters VI to IX, inclusive.

GENERAL COLUMNAR SECTION OF ROCKS EXPOSED IN GREENBRIER COUNTY

VERTICAL SCALE: 1 INCH=1000 FEET











ERA	PERIOD OR SYSTEM	SERIES	MAP SYM.	SECTION	THICK. FEET	TOTAL FEET	DESCRIPTION
QUAT.	RECENT		QAL		?	?	Unconsolidated clays and gravel. (River wash)
	PLEISTOCENE		QAL		?	?	Unconsolidated clays and gravel. (River terraces)
PALEOZOIC	UPPER CARBONIFEROUS PENNSYLVANIAN	POTTSVILLE					
		KANAWHA GROUP (PART)	Ca		250±	250	Massive gray sandstones; gray sandy and dark carbonaceous shales; coals; fresh or brackish water fauna; plant fossils.
		NEW RIVER GROUP	Gnr		600-950	1200	Massive gray sandstones; gray sandy and dark carbonaceous shales; minable coals; fresh or brackish water fauna; erratic boulders in Sewell Coal; plant fossils.
		POGAHONTAS GROUP	Cr		0-340	1540	Massive gray sandstones; gray sandy and dark carbonaceous shales; minable coals; fresh or brackish water fauna; plant fossils.
	LOWER CARBONIFEROUS MISSISSIPPIAN	MAUCH CHUK					
		BLUE-STONE GROUP	Cbl		80-675	2215	Red, green and variegated shales; green, gray and brown massive and flaggy sandstones; thin streaks of coal; marine fauna and plant fossils.
		PRINCETON CONGLOM.	Cpr		20-80	2295	Massive gray and brown sandstone with variegated pebbles; poorly sorted; plant fossils.
		HINTON GROUP	Chn		500-850	3145	Red, green and variegated sandy shales; thin limestones; red and brown sandstones; massive sandstone at base (Stony Gap); marine and plant fossils.
		BLUE-FIELD GROUP	Gbf		1000-1200	4345	Brown, red, green, variegated argillaceous, calcareous and fissile shales; massive and flaggy sandstones; limestones in lower part; coal streaks; abundant marine fauna; some plant fossils.
		GREENBRIER	Cgr		475-750	5095	Dark gray, massive limestone; thin streaks of calcareous shale; white oolite 100± feet from top; streaks of red shale or limestone below

FIGURE 7

GENERAL COLUMNAR SECTION (CONCLUDED)

ERA	PERIOD OR SYSTEM	SERIES	MAP SYM.	SECTION	THICK. FEET	TOTAL FEET	DESCRIPTION
PALEOZOIC	UPPER DEVONIAN (CONTINUED)	PORTAGE	Dp		2000±	11345	Gray, green, sandy and argillaceous shale interbedded with grayish green and brown flaggy sandstone; sparse marine fauna; land plants.
		GENESEE	Dg		50-100	11445	Brown to black slaty shale; marine fauna.
		(HAMILTON) MARCELLUS	Dm		500±	11945	Black carbonaceous, fissile slickensided shale; thin limestone in lower portion; septaria; marine fauna.
	MID. DEV.	ORISKANY	Do		80-90	12035	Upper part, light to dark calcareous chert; gray to brown ferruginous sandstone; marine fossils.
		HELDERBERG	Dhl		300±	12335	Massive, blue, tough, cobbly limestone; chert nodules; marine fauna.
	LOWER DEVON.	BOSSARDVILLE	Sbo		250±	12585	Blue to gray, massive to platy ls. calcite streaks; marine fauna.
		RONDOUT	Srd		200±	12785	Flaggy, brittle limestone; sparse marine fauna.
		NIAGARA	Sna		100±	12885	Massive, dark gray limestone; peodes; marine fauna.
		CLINTON	Scl		600±	13485	Massive, gray, quartzitic sandstone at top; thin limestones in upper part; variegated shale near middle; massive gray and red (iron ore) ss. near base; marine fossils.
		WHITE MEDINA	Swm		100±	13585	Massive, white, very hard quartzite, Scolithus and sucoids.
		RED MEDINA	Srm		80±	14385	Deep red shale alternating with red and reddish-brown sandstone; no fossils found.

FIGURE 7
GENERAL COLUMNAR SECTION
(CONTINUED)

ERA	PERIOD OR SYSTEM	SERIES	MAP SYM	SECTION	THICK. FEET	TOTAL FEET	DESCRIPTION
LOWER CARB. MISSISSIPPIAN (CONTINUED)		MACGRADY	Cmcc		60-250	5345	Deep red, purple shale; occasional sandstone.
		POCONO	Cpo		200-600	5945	Brown or red cross-bedded sandstone; conglomerate at top and base; sandy shale; lenticular beds of semi-anthracite coal; numerous marine fossils and land plant fossils.
		CATSKILL	Dck		0-400	6345	Mostly red shale and sandstone; some brown shale; marine fauna and plant fossils.
UPPER DEVONIAN		CHEMUNC	Dch		2000-3000	9345	Gray, green, brown, massive and flaggy sandstone, interbedded with gray, green arenaceous and argillaceous shale; sandstone conglomerate at top (Hendricks); abundant marine fossils; some land plant fossils.
PALEOZOIC (CONTINUED)							
PALEOZOIC							

CHAPTER IV.

Structural Geology.

INTRODUCTION.

In order to appreciate the structural geology of Greenbrier County it is necessary to analyze it in its general position and relationship with the surrounding areas. It must be kept in mind that the county has received its proportionate share of the disturbances that have affected the Appalachian area in general. By its structure is meant the position in which the strata are now found;—their position or deviation from the horizontal, the approximate position in which they were originally deposited.

Preceding discussion has shown us that the sediments were deposited on the floor of a shallow sea, the bottom of which slowly sank to permit the accumulation of thousands of feet of muds, sands, and limes. However, all those buried for any considerable depth had been compacted into their consolidated equivalents, shales, sandstones, and limestones.

These rocks were then subjected to tremendous earth stresses. These stresses were coming from the east and south-east and were of mountain-making proportions. The geologic time was during the latter part of the Permo-Carboniferous Period. The forces were of sufficient magnitude to move the ancient crystalline mountains, on the east, bodily westward so as to squeeze these sediments which had been deposited in the sedimentary trough into many elongated folds. This tangential or compressive stress tended not only to fold or buckle the rocks but mash and telescope them in such a way that they were thickened en masse and raised from beneath the sea. *This episode in geological history is known as the Appalachian*

the approximate interval to the Sewell Coal is known. In this way the position of the key horizon (Sewell Coal) can fairly accurately be determined, whether it is below drainage or whether it has been removed from the tops of the hills.

The detailed work necessary to prepare the structure map included several hundred observations on the key horizon and other known stratigraphic horizons. Elevations were obtained either by aneroid barometer, checked on the nearest Government spirit-level determination as recorded on the topographic maps, or from spirit-level determinations furnished by engineering departments of several operating companies.

In Greenbrier County there is considerable variation in the intervals between the different stratigraphic horizons due to the thickening or thinning of the intervening measures. For this reason it must not be assumed that the structure on other horizons conforms exactly to that of the key horizon (Sewell Coal). In order to better determine the position of other beds, a table of intervals was prepared from numerous detailed stratigraphic cross-sections and measurements of intervals from place to place. The principal results of these data are condensed in the following table which shows the intervals above and below the Sewell Coal. These tables were used in determining the contours on the key horizon in localities where direct observations could not be made:

height, as the exposed sediments were immediately attacked by weathering agencies which would have reduced them to sea-level instead of a fairly even-crested plain during the course of the vast lapse of time that followed, had not the entire area again been subjected to earth stresses of mountain-making proportions. This time, however, the stresses operated vertically rather than horizontally, as had the previous example, and are responsible for the greater part of our present elevation. It is true that the entire area has since been subjected to one more rejuvenation, but of less magnitude than either of the preceding movements. The present topography is the result of the interaction of these forces with the atmosphere or weathering agents.

METHODS OF GEOLOGIC WORK AND REPRESENTATION OF STRUCTURE.

The method of determining the structure, or position of the rocks in Greenbrier County was not the same in all parts of the county. In the western part of the county where the rocks have been only slightly disturbed and where the strata are still practically horizontal, there are some well-defined beds, where it is possible to measure thicknesses and determine dips over fairly wide areas, by means of aneroid barometer levels, with considerable accuracy.

In this region a structure map has been made, showing the position of the base of the Sewell Coal of the New River Group of the Pottsville Series in the region where this coal occurs. This area includes the Meadow Creek and Big Clear Creek commercial fields where many elevations are available. That portion of the Cherry River drainage in Greenbrier County, including the North and South Forks, is practically uninhabited. Second-growth timber is in part about large enough to cut again. Travel is with difficulty and must be made on foot. Under these conditions and with very little prospecting, information on the coal is only slight. However, other key horizons from which the approximate interval to the Sewell Coal is known have been used to show the base

side, where the rocks have suffered greater deformation, different methods of stratigraphic work are necessary. In a large part of this area the rocks have been severely deformed, leaving them tilted, vertical, occasionally overturned, and sometimes faulted. In such areas the aneroid and level are of minor importance, but the combination clinometer and pocket transit takes their place. With this instrument numerous dip and strike readings were taken, most of which are shown on Map II (in Atlas). By using the accurate topographic maps many cross-sections across the dip were made, and accurate contact lines of the different series were mapped. Four cross-sections have been plotted to a scale of 1:62,500 both horizontally and vertically and appear on the upper right corner of Map II. In other localities, where conditions were favorable, horizontal measurements were made across the dips to secure data for compilation of thickness by trigonometric formulae, and the resulting sections, along with those vertically measured in the western half of the county, appear in Chapter V under the heading of "Measured Sections."

DETAILED STRUCTURE.

ANTICLINES AND SYNCLINES.

Webster Springs Anticline.—The Webster Springs Anticline of Reger¹ has been traced from northern Webster County, across the eastern edge of Nicholas County to the Greenbrier County line about $3\frac{1}{2}$ miles northeast of Richwood. Along the crest of the fold at the county line the Sewell Coal has an elevation of about 3175 feet. Along the crest southwestward there is a gradual rise of almost 100 feet to the mile and at the high point along the fold, one mile south of Mann Knob, the Sewell Coal has an elevation of about 3725 feet. From this point the fold trends a little more to the west passing just south of Beech Knob and pitches at the rate of about 30 feet to the mile. From this point the axis of the anticline gradually bends more and more toward the south, and near the headwaters of Sam Creek there is a structural saddle with the ele-

Intervals Above and Below Sewell Coal, Greenbrier County.

	Anjeau	Big Clear Creek Mountain	Boggs Knob (Sims Station)	Charmo	Cross Mountain	Duo	Grassy Knob	Manning Knob	Namo Chapel (Jetsville)	N. F. Cherry River at County Line	Quinwood	Russellville	S. F. Cherry River
Nuttall Sandstone (top).....	510	450	490
s Ferry Coal.....	290	250	280	270
Coal.....	110	100	150	90	85	120	120
Coal.....	0	0	0	0	0	0	0	0	0	0	0	0
Raleigh Coal.....	160	150	140	140	170	155	140	140	140	120	130	140
y Coal.....	265	270	240	250	280	250	270	250	215	190	250	250
reek Coal.....	325	330	325	320	365	340	350	325	260	235	305	340
Pocahontas Coal.....	430	425	450	435	450	450	450	420	420
Pocahontas Coal.....	525	530	500	500	540	525	540	490	500
Pocahontas Coal.....	635	625	600	600	650	615	640	?
of Pottsville.....	660	710	660	750	750	650	720	550	400	300
ton Sandstone (top).....	835	1050	940	1075	1165	900	550	440

Cherry River at Fenwick, W. Va.

Location.—Chain gage at highway bridge at Fenwick, Nicholas County, 1,000 feet below mouth of Laurel Creek.

Drainage area.—150 square miles.

Records available.—September, 1929, to September, 1932.

Extremes.—Maximum gage height during year, 14.58 feet July 4 (discharge not determined); minimum discharge, 0.1 second-foot Sept. 13 (gage height, 2.60 feet).
1920-32: Maximum gage height, that of July 4, 1932 (discharge not determined); minimum discharge, 0.1 second-foot Sept. 22, 1930, Sept. 13, 1932 (gage height, 2.60 feet).

Remarks.—Records good.

Discharge, in second-feet, 1931-32.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	175	115	455	1,700	885	236	1,790	2,390	41	230	21	5.2
2.....	123	163	590	1,340	815	214	1,250	1,620	37	214	20	3.0
3.....	97	152	455	885	1,340	204	780	885	51	197	61	47
4.....	77	109	430	080	4,140	1,160	710	620	50	8,300	300	16
5.....	61	138	380	780	2,290	1,120	430	430	43	5,240	93	12
6.....	58	121	300	1,700	1,100	1,080	530	360	39	3,630	97	9.2
7.....	53	105	280	1,000	710	885	605	300	33	1,700	60	7.6
8.....	50	97	260	1,340	710	650	455	230	26	1,120	50	5.5
9.....	43	90	380	1,250	710	060	340	228	36	530	37	7.0
10.....	39	81	1,000	1,080	620	815	300	222	21	260	34	12
11.....	43	72	960	815	505	560	300	430	19	500	55	3.2
12.....	47	71	1,990	505	1,080	300	815	815	23	340	77	1.4
13.....	43	74	3,040	455	885	197	590	1,160	75	225	47	.1
14.....	48	70	2,390	340	710	142	480	920	138	152	28	2.2
15.....	55	72	1,430	280	530	105	430	050	90	197	21	.8
16.....	63	66	780	230	360	140	340	505	67	187	17	2.0
17.....	86	69	020	204	430	4,270	300	405	72	138	15	1.4
18.....	91	03	455	214	745	3,270	232	300	54	88	16	1.7
19.....	82	64	430	104	530	1,340	204	255	43	64	101	1.2
20.....	00	63	430	106	430	1,120	181	100	35	51	41	2.2
21.....	60	65	430	158	340	1,000	158	194	27	37	34	1.8
22.....	54	54	455	158	380	1,700	140	480	32	46	24	2.4
23.....	48	57	780	269	340	1,120	125	242	25	63	16	0.0
24.....	43	51	650	380	256	815	142	178	20	71	13	11
25.....	54	57	560	360	239	650	300	142	14	46	11	8.8
26.....	48	51	480	300	218	455	530	133	16	30	9.2	5.5
27.....	43	84	430	620	214	1,080	480	109	16	31	7.3	7.3
28.....	57	280	430	560	190	9,560	455	00	2,000	31	5.8	7.0
29.....	72	300	505	1,430	222	3,880	430	74	1,040	47	7.3	4.9
30.....	84	320	480	4,540	1,610	620	71	380	40	7.6	11
31.....	95	455	3,270	1,610	60	28	7.3

Month	Maximum	Minimum	Mean	Per square mile	Run-off in inches
October	175	39	66.2	0.441	0.51
November	320	51	106	.707	.79
December	3,040	260	733	4.89	5.64
January	4,540	158	910	6.07	7.00
February	4,140	190	758	5.05	5.45
March	9,560	105	1,370	0.13	10.53
April	1,700	125	480	3.20	3.57
May	2,390	00	471	3.14	3.02
June	2,090	14	155	1.03	1.15
July	8,300	28	775	5.17	5.96
August	200	50	120

Cherry River at Fenwick, W. Va.

Location.—Chain gage at highway bridge at Fenwick, Nicholas County, 1,000 feet below mouth of Laurel Creek. Zero of gage is 2,088.94 feet above mean sea level.

Drainage area.—150 square miles.

Records available.—September, 1929, to September, 1934.

Extremes.—Maximum discharge recorded during year ending Sept. 30, 1933, 5,520 second-feet Jan. 21 (gage height, 9.84 feet); minimum, 7.2 second-feet Oct. 5 (gage height, 2.97 feet).

Maximum discharge recorded during year ending Sept. 30, 1934, 5,800 second-feet Mar. 5 (gage height, 10.04 feet); minimum, 3.1 second-feet Oct. 7 (gage height, 2.76 feet).

1929-34: Maximum gage height recorded, 14.58 feet July 4, 1932 (discharge not determined); minimum discharge, 0.1 second-foot Sept. 22, 1930, Sept. 13, 1932, (gage height, 2.60 feet).

Remarks.—Records fair. Discharge estimated Oct. 10, 14, 15, Nov. 29 to Dec. 4, Dec. 16, 1932, Jan. 27-29, Mar. 1, 2, Apr. 30 to May 20, Aug. 15, Sept. 27, 1933.

Discharge, in second-feet, 1932-33.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	11	682	110	780	425	500	748		175	358	266	60
2.....	0.2	590	110	590	715	400	815		126	274	285	45
3.....	8.9	880	100	450	650	293	885	283	111	1,200	748	48
4.....	8.0	293	100	380	590	259	815		95	502	4,010	07
5.....	8.9	248	93	335	425	181	650		89	282	1,420	210
6.....	204	194	86	248	282	197	748		63	184	650	69
7.....	115	161	79	197	335	335	1,510		60	118	380	68
8.....	33	156	77	207	2,450	850	995	867	40	81	250	46
9.....	22	1,330	60	358	1,420	650	748		99	68	178	28
10.....	24	1,780	72	530	682	502	560		650	76	148	46
11.....	26	885	76	450	502	380	530		204	150	620	26
12.....	17	650	241	502	314	380	1,780		143	314	560	23
13.....	18	380	314	402	314	380	995	814	146	143	475	16
14.....	15	278	282	335	958	2,750	715		91	109	475	36
15.....	15	241	234	285	1,870	8,750	530		64	278	220	37
16.....	38	234	150	210	1,330	1,090	475		76	425	133	115
17.....	1,240	293	150	220	1,070	958	590		66	270	120	79
18.....	715	259	175	263	850	1,330	502	1,290	54	158	97	28
19.....	402	1,960	175	450	850	2,550	590		44	111	89	38
20.....	380	1,200	140	380	1,960	2,250	475		35	88	63	46
21.....	266	748	136	4,400	1,600	2,250	402	224	26	77	61	36
22.....	181	450	187	2,950	995	1,330	358	230	22	60	53	36
23.....	131	314	358	1,510	715	885	282	200	18	44	45	28
24.....	101	285	475	885	620	650	244	178	16	38	314	17
25.....	81	244	682	850	682	530	293	200	52	36	161	15
26.....	89	220	502	1,030	1,420	425	358	224	167	111	77	12
27.....	748	184	530	700	815	335	335	200	230	1,030	45	12
28.....	402	148	2,250	530	682	402	285	210	650	1,240	60	14
29.....	314	125	1,150	450		450	259	178	1,240	1,780	153	15
30.....	213	120	885	380		450	220	230	850	502	122	12
31.....	227		780	335		502		217		475	84	

Month	Maximum	Minimum	Mean	Per square mile	Run-off in inches
October	1,240	8.0	196	1.31	1.51
November	1,960	120	501	3.34	3.73
December	2,250	69	347	2.31	2.66
January	4,400	197	697	4.65	5.36
February	2,450	282	911	6.07	6.32
March	3,750	181	929	6.19	7.14
April	1,780	220	623	4.15	4.63
May		178	599	3.99	4.60
June	1,240	16	190	1.27	1.42
July	1,780	36	341	2.27	2.62
August	4,010	45	399	2.66	3.07

Cherry River at Fenwick, W. Va.
(Continued)
Discharge, in second-feet, 1933-34.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	9.5	8.9	425	885	172	270	502	234	77	14	32	17
2.....	17	7.5	335	1,240	213	1,420	425	204	54	20	129	15
3.....	35	16	314	715	181	4,680	314	181	45	14	682	24
4.....	18	19	314	560	158	4,400	293	161	30	8.9	358	10
5.....	14	28	402	500	150	4,960	278	143	35	10	175	15
6.....	13	130	815	748	109	2,550	227	131	68	8.3	103	12
7.....	3.3	77	995	1,110	71	1,600	255	120	172	7.8	75	12
8.....	0.2	79	780	958	113	2,150	248	165	91	9.8	68	10
9.....	6.2	50	590	748	103	1,780	220	95	64	13	42	7.6
10.....	5.5	45	314	530	99	1,030	197	130	58	9.5	32	7.0
11.....	12	46	234	380	77	1,510	200	590	77	8.3	66	7.0
12.....	8.6	64	158	335	84	1,110	259	293	99	14	60	5.3
13.....	7.0	71	204	285	103	715	335	248	55	12	74	9.8
14.....	5.3	81	224	244	95	502	380	335	40	14	52	15
15.....	8.3	131	207	227	122	203	450	682	32	15	35	12
16.....	8.0	120	234	181	138	358	550	650	24	8.3	682	402
17.....	181	164	380	133	140	335	650	550	20	5.3	780	335
18.....	07	380	650	101	146	380	850	314	10	0.0	278	130
19.....	36	335	1,110	143	158	650	815	194	278	6.0	143	86
20.....	42	207	3,750	93	148	995	780	194	107	5.3	105	58
21.....	14	184	2,550	101	143	780	650	158	81	4.5	88	38
22.....	31	285	2,150	118	122	715	530	138	45	5.8	55	40
23.....	20	293	2,150	138	170	530	330	133	42	4.9	63	31
24.....	31	220	1,600	156	129	425	358	122	49	6.8	89	26
25.....	25	178	358	178	133	402	285	111	31	8.6	101	48
26.....	11	158	358	175	172	560	244	95	22	6.2	84	45
27.....	12	197	293	190	230	1,900	293	84	18	5.3	74	30
28.....	20	285	241	194	282	2,250	335	74	16	402	46	41
29.....	18	282	153	213	1,330	293	66	12	95	31	131
30.....	20	314	93	200	850	259	58	14	72	35	1,690
31.....	17	207	190	682	57	48	22

Month	Maximum	Minimum	Mean	Per square mile	Run-off in inches
October	181	3.3	24.5	0.163	0.19
November	380	7.5	149	.993	1.11
December	3,750	93	732	4.88	5.63
January	1,240	93	389	2.59	2.90
February	282	71	141	.940	.08
March	4,960	270	1,360	9.07	10.46
April	850	197	396	2.64	2.94
May	682	57	215	1.43	1.55
June	278	12	58.9	.393	.44
July	402	4.5	28.0	.187	.22
August	780	22	150	1.00	1.15
September	1,690	5.3	111	.740	.83
The year	4,960	3.3	310	2.11	23.59

Cherry River at Fenwick, W. Va.

Location.—Chain gage, lat. 38°13'45", long. 80°35', at highway bridge at Fenwick, Nicholas County, 1,000 feet below mouth of Laurel Creek. Zero of gage is 2,088.04 feet above mean sea level.

Drainage area.—150 square miles.

Records available.—September, 1929, to September, 1935.

Extremes.—Maximum discharge observed during year, 4,740 second-feet Mar. 12 (gage height, 8.95 feet); minimum, 5 second-feet Oct. 5 (gage height, 2.07 feet).

1929-35: Maximum observed gage height, 14.58 feet July 4, 1932 (discharge not determined); minimum discharge, 0.1 second-foot Sept. 22, 1930. Sept. 13,

Rating tables, water year 1934-35 (gage height, in feet, and discharge in second-feet)							
Table for Oct. 1 to Mar. 11				Table for Mar. 12 to Sept. 30			
3.3	23	4.5	293	2.9	3.0	4.5	326
3.4	32	5.0	530	3.0	5.5	5.0	570
3.5	45	5.5	850	3.1	10	5.5	880
3.6	58	6.0	1,240	3.2	16	6.0	1,260
3.7	74	7.0	2,210	3.4	34	7.0	2,230
3.8	91	8.0	3,380	3.6	63	8.0	3,380
4.0	133	9.0	4,740	3.8	102	9.0	4,740
4.2	187			4.0	153		

Discharge, in second-feet, water year October, 1934, to September, 1935

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	425	336	1,700	402	185	425	2,900	142	348	100	240	18
2.....	278	422	1,110	475	153	402	1,820	209	302	290	174	12
3.....	190	203	815	380	136	358	1,140	302	320	320	100	10
4.....	136	203	682	314	105	314	845	348	1,440	174	110	01
5.....	111	380	502	278	79	314	655	313	1,060	515	142	022
6.....	402	475	402	227	86	358	625	465	655	055	92	508
7.....	402	500	335	204	64	314	778	3,220	410	1,020	2,310	322
8.....	282	450	274	217	99	203	055	1,720	322	1,020	2,450	225
9.....	190	335	217	450	133	260	685	880	415	1,440	950	168
10.....	143	241	187	620	530	314	508	715	300	778	542	156
11.....	122	190	187	502	530	2,030	655	490	218	309	465	87
12.....	95	170	238	358	475	4,040	745	302	183	268	326	67
13.....	74	153	230	282	402	2,230	1,220	490	183	180	199	47
14.....	04	146	111	259	475	1,180	1,060	810	140	140	369	44
15.....	61	111	01	227	743	988	880	1,180	159	232	225	43
16.....	53	105	95	417	748	1,020	655	1,060	153	102	148	33
17.....	50	103	81	2,000	560	745	570	988	187	04	100	30
18.....	42	101	80	1,700	402	625	542	715	369	77	74	21
19.....	42	105	30	335	335	515	542	515	348	63	68	21
20.....	37	89	80	2,000	270	745	440	405	348	52	55	22
21.....	36	80	289	2,430	187	685	360	773	204	70	57	20
22.....	31	90	217	2,540	230	830	326	655	490	76	44	18
23.....	31	2,130	175	2,100	314	2,500	279	685	515	72	38	14
24.....	35	1,600	173	1,800	748	2,900	232	685	415	00	28	11
25.....	30	748	207	995	650	2,070	205	745	286	860	22	10
26.....	54	530	958	590	1,070	1,815	171	625	106	490	10	7
27.....	49	380	815	425	885	1,100	142	440	156	392	61	9
28.....	58	335	682	358	560	845	148	343	119	246	37	7
29.....	70	1,310	560	314	715	137	294	137	162	22	7
30.....	45	1,110	502	244	593	153	655	124	98	21	5
31.....	38	425	215	1,960	515	63	20

Month	Second-foot-days	Maximum	Minimum	Mean	Per square mile	Run-off in inches
October	3,090	425	30	119	0.793	0.01
November	13,386	2,180	86	446	2.97	3.31
December	12,526	1,700	81	404	2.69	3.10
Calendar year 1934.....	110,934.2	4,060	4.5	320	2.13	28.98
January	24,208	2,540	204	781	5.21	6.01
February	11,150	1,070	64	309	2.66	2.77
March	34,204	4,040	266	1,103	7.35	8.47
April	20,172	2,900	13.7	672	4.48	5.00
May	21,939	3,220	142	708	4.72	5.44
June	10,554	1,440	119	352	2.35	2.62
July	10,704	1,620	52	345	2.80	2.05
August	9,520	2,450	16	307	2.05	2.36
September	2,724	622	5	90.8	.005	.68

resulting from differences in the composition and structure of the rocks.

Another type of sink-hole, quite common in Greenbrier County, is due to the collapse of the roof of underlying caverns. Sinks due to this cause are quite irregular in shape and are often elongated. They are usually steep walled and are often quite large. It is the exception, rather than the rule, for the cavern roofs to collapse suddenly and usually the settling is so gradual that it would hardly be noticed by a resident of the region.

The average sink-hole in Greenbrier County owes its origin to a combination of the two main causes discussed above. In general they have been excavated above the water-table, drain downward through openings in their floors and are therefore usually dry. The outlets of some, however, are clogged by clay, humus, and other insoluble matter washed into them, allowing the development of small lakes whose levels are above the water-table and independent of it. In some sinks the water leaks away slowly; in others the insoluble stopper is suddenly broken through and the lake disappears with a rush.

Caverns.—Caverns of many sizes and shapes occur in the limestones of Greenbrier County. In so far as they have been explored most of the caverns are small, many of them hardly extend beyond the twilight zone. From the vast area in which no surface streams are present it is apparent that many of these small caverns must interconnect. However, these connecting passageways may be small and difficult to traverse. Some of the caves are smooth walled, showing only the effects of solution, while others are sparingly ornamented with calcite deposited from solution.

The process of precipitation by subsurface waters is clearly evident in the deposits of calcite in the form of drip-stone. Vadose (ground) water charged with calcium carbonate percolates downward from the surface of the ground to the roof of the cavern, where, clinging to the ceiling, it forms drops. While at rest it evaporates a little, loses some

Solution in Carbonate Rocks.—Pure water dissolves mineral matter but water containing oxygen, carbon dioxide, and acids is a vastly more efficient solvent. The rain water that reaches the rocks is not pure. In falling through the atmosphere it acquires oxygen and carbon dioxide, and in percolating through the crust of vegetation and the underlying soil in humid regions it absorbs more carbon dioxide as well as various organic acids formed by the decomposition of plant matter.

Limestone is soluble in water charged with carbon dioxide and therefore in humid regions, where rainfall is plentiful and evaporation relatively slight, it is vigorously attacked by subsurface water with striking results. Great holes are formed in the surface, caverns are hellowed out below ground, and surface streams are undermined and led away through subterranean channels. That subsurface water is responsible for this work is shown by the fact that the water of springs and wells in regions of limestone and dolomite is "hard"; that is, it contains much calcium carbonate in solution.

Sinks.—In compact, well-stratified limestones, such as those in Greenbrier County, the easiest descent for vadose water is through vertical joints and along bedding-planes. Those avenues most favorably situated with respect to supply from above and free circulation below are readily enlarged by solution as the descending water passes through them. Enlargement is most effective at the surface, where movement of the water is most rapid and where the water is freshly charged with carbon dioxide from the atmosphere and from decaying vegetation, and decreases rapidly downward. In consequence the point of intersection of two joints near the surface becomes a funnel-shaped depression. As the depression widens, the overlying mat of insoluble mantle and vegetation collapses into it, and a sink is formed. Sinks of this (funnel) type range in size from small openings only a few inches in di-

Then, as more water is added from above, it is forced to drop, falling on the floor below, it evaporates still further, leaving another minute deposit. As the drops slowly but endlessly succeed each other, long "icicles" of calcite (stalaetites) grow downward from the roof, while broader accumulations (stalagmites) grow upward from the floor. If the process goes on long enough each pair coalesces and forms a column. Drip-stone assumes many fantastic shapes, curious to the cavern visitor, but all are formed in this simple way.

In past times caverns often served as refuges for primitive man and as dens for animals that are now extinct. Because of this the bones of men and animals, stone implements, and other objects have accumulated in the caves and have often been sealed up beneath deposits of calcium carbonate slowly accumulating on their floors. Relics of this kind, especially in certain parts of Europe, have revealed much concerning the life and culture of the times before the beginning of written history.

The following item taken from "The Pleistocene of North America and its Vertebrated Animals from the States East of the Mississippi River and from the Canadian Provinces East of Longitude 95°", by Oliver P. Hay, Carnegie Institution of Washington, Washington, D. C., pp. 34-35, 1923, records the finding of several bones of a prehistoric sloth in a cave in Greenbrier County:

"In a cave situated somewhere in this county were found the bones described in 1799 by President Thomas Jefferson (Trans. Amer. Philos. Soc., Vol. IV, pp. 246-260) under the name *Megalonyx*. Colonel John Stewart became interested and saved some of the bones from being carried away by curious inhabitants of the region.

"The bones, a distal end of a femur, a complete radius, a complete ulna, three claws, and some other foot-bones were secured and presented to the American Philosophical Society of Philadelphia, from which they passed into the possession of the Academy of Natural Sciences, where they are still preserved. Some of these were described by Dr. Caspar Wistar (Trans. Amer. Philos. Soc., Vol. IV, 1799, p. 526, plates I, II).

"Inasmuch as this species may have existed during a large part of the Pleistocene and certainly after the passing of the Wisconsin epoch, and inasmuch as no other species were found associated with

own satisfaction that the bones were found in what is now known as Organ Cave in southern Greenbrier County. (See Maps I and II in Atlas).

Present Fauna in the Caves.—An interesting account of the life to be found in caves of the State is to be found in the Proceedings of the West Virginia Academy of Science, West Virginia University Bulletin, series 34, No. 15, pp. 39 to 53, 1934. In this paper Professor A. M. Reese, of the University Department of Biology, gives a detailed account of his visit to 43 caves. The following descriptions of the caves of Greenbrier County are taken from the paper just cited:

"Organ Cave, visited April 26, 1932.

"This, as has been said, is one of the few commercial caves of the State and is easily located by watching for the advertising signs along Route 24, (U. S. Route 219), in the lower side of the county near the Monroe County line. It is situated about one-half mile east of Route 24, (U. S. Route 219). The entrance is large and is at the base of a high, rocky cliff. The cave is partially lit by electricity. Some interesting formations are to be seen. In this cave, also a number of wooden troughs, for collection of saltpeter, said to have been used during the war between the States. A considerable pond of water is here but at the time of our visit it was very cloudy and no animals could be found in it. No insects were seen. Several bats were collected, but were misplaced and so are not named here.

"The West Virginia Biological Expedition on July 30, 1931, found adults and larvae of the salamander, *Desmognathus fuscus fuscus*. This party also found *Rana clamitans* in the cave and *R. sylvatica* at the cave entrance, both probably accidental visitors."

On June 21, 1929, the senior author was shown through the cave by the manager, Mr. S. M. Sively. The cave, which is electrically lighted is in the Hillsdale member of the Greenbrier Limestone. Water that was colored for testing was found to emerge on Second Creek. In addition to the many interesting formations of dripstone, one of the main attractions to the visitor is the presence of 37 saltpeter hoppers used in making gunpowder by the Confederates in the Civil War. Of interest to the geologist is the fact that in this cave was found the bones of the Pleistocene *Xenarthra Megalonyx Jeffersonii*, named and described by President Jefferson in 1799.

"A dirt road leads northwest from Route 24, (U. S. Route 219), 3.5 miles north of Frankford; if this road be followed for one mile it will lead to the home of Mr. J. Rapp; the cave is about 200 yards behind and below the house, opening by a fairly large hole into the side of a steep hill. The rooms are fairly large but do not extend very far, perhaps 200 yards. The floor is very rough with fallen rocks; it was damp but no actual stream was present at time of visit. Considerable numbers of stalactites and other formations were present. No rats were found but three bats were collected, two Georgian, *Pipistrellus subflavus subflavus*, and one brown, *Myotis lucifugus*. The only other animals found were a few crickets, *H. subferraneus* Scudder."

"Arbuckle's Cave, visited October 1, 1932.

"This small cave is located one-half mile east of Route 24, (U. S. Route 219), in the rear of the brick residence of Dr. Arbuckle, at Maxwellton. Its fairly large opening is about 50 feet behind a small farm-house and leads into a roomy passage, with a smooth floor, that extends about 100 yards into a hill. The only actual water in the cave is a rocky tank, about 3 x 8 feet in size. The animals found were one Georgian bat, *Pipistrellus subflavus subflavus*; three salamanders, *Plethodon wehrlei*; many crickets, *H. subterraneus* Scudder; a few flies, *Amoebaleria defessa* O. S.; a few blind beetles, *Pseudanophthalmus grandis* Valentine; and one milliped, *Pseudotremia caverarum* Cope.

"McClung's Cave, visited October 1, 1932.

"This interesting cave is easily reached by following the road that leads from the east side of Route 24, (U. S. Route 219), at Maxwellton. About 2½ miles northeast of Maxwellton this road leads directly to the residence of Mr. McClung; the entrance to the cave is about 50 feet from the house. The cave which extends in a westerly direction passes almost directly beneath the house; its chambers are very roomy for a hundred feet or more, then contract to a high, narrow cleft with many fallen rocks. A considerable number of stalactites may be seen. Even in the extremely dry season, when the cave was visited, a small stream flowed towards the west, away from the entrance. The walking was difficult, and time allowed the cave to be followed for only 200 to 300 yards; but in this short distance the following animals were found: many crickets, *Hadenoeus subterraneus* Scudder; some diptera, *Amoebaleria defessa* O. S.; several blind beetles, *Pseudanophthalmus grandis* Valentine; several small gastropods, *Helicodiscus parallelus* Say; myriapods, four unidentified specimens; earthworms, *Helodrilus ealiginosus trapezoides* Duges; several salamanders, *Eurycea lucifuga* and *Desmognathus fuscescens*; no bats or rats were seen."

"Saltpeter Cave No. 1, visited April 26, 1932.

"This is one of the numerous caves known as saltpeter caves; it is located near the Tennant homestead at Blaker's Mill, between the Fort Spring road and the road from Alderson to Blue Sulphur Springs. The cave has a fairly large entrance but is not very extensive. Two or three bats, *Pipistrellus subflavus subflavus*; several crickets, *H.*

Sanpete. Cave No. 1, 1932.
"Located in the same hillside and about 100 yards from the preceding cave. It may be that the two caves are one. The cave was entered by climbing down a tree ladder into a large sink hole. One bat was seen, and some cave crickets, *H. subterraneus* Seudder, were found."

Bunger's Cave No. 1, visited April 26, 1932.

"This cave is about 1½ miles south of Route 60 (Midland Trail) about 8 miles west of Lewisburg. The road to this cave leaves Route 60 just east of the schoolhouse, close to the south side of Route 60. The entrance is large, rough and steep, with a stream about 50 yards from the opening. The cave can be followed only a short distance beyond the twilight zone. The only life found were three bats, two *Myotis lucifugus lucifugus*, and one *Pipistrellus subflavus subflavus*."

"Bunger's Cave No. 2, visited April 26, 1932.

"Located about one-half mile from the preceding cave at the side of a broad meadow. A wide, steep entrance ends, after about 50 feet, in a clear stream about 10 feet wide and 1 foot deep. This stream was waded for about 200 yards but no life of any sort was seen."

"Higginbotham's Cave No. 1, visited June 24, 1932.

"This cave is located about one mile northwest of Frankford on the farm of Mr. O. D. Higginbotham, in the side of a hill. The main passages of the cave extend in opposite directions from the fairly large entrance and are high enough for erect walking in most places, so that it is an easy cave to explore. A slow-moving stream flows towards the south. Numerous stalactites are present. No bats or rats were seen. The animals found were: one salamander, *Plethodon cinereus* (dark phase); numerous crayfish, probably *Cambarus bartonii ecarinirostris* Hay; numerous cave crickets, *H. subterraneus* Seudder; numerous diptera, *Amoebaleria defessa* O. S.; and eight or ten blind beetles, *Pseudanophthalmus grandis* Valentine."

"Higginbotham Cave No. 2, visited June 24, 1932.

"This is a small cave, situated about one-half mile southwest of the preceding cave; its entrance is a sort of small shuk-hole. No running water was present at the time it was visited. The only animals seen were many cave crickets, *H. subterraneus* Seudder, and two or three blind beetles, *Pseudanophthalmus grandis* Valentine."

"Coffman's Cave, visited June 24, 1933.

"This cave lies about one mile southwest of the preceding cave and 100 yards from the Coffman residence. The entrance is large and lies at the base of a rocky cliff. A good stream flows in the cave, in a direction away from the entrance, which has to be waded at places; lack of time and the appearance of a large pond stopped further investigation of this cave. The animals collected were: four salamanders, *Gyrinophilus porphyriticus*, (two larvae and two adults); a few crayfish, *Cambarus bartonii ecarinirostris* Hay; and many cave crickets, *H. subterraneus* Seudder. Numerous traps, set by Dr. Val-

Lizard Cave.

"This cave is $1\frac{1}{2}$ miles west of Alderson, on the road from Alderson to Hinton. It was not visited by the writer, but a specimen of *Eurycea lucifuga* was secured from the cave through Mr. R. H. Fletcher."

"Mud Cave.

"This is also on the road from Alderson to Blue Sulphur Springs, about $2\frac{1}{2}$ miles from the preceding cave. It was not visited by the writer. A specimen of *Eurycea lucifuga* from the cave was received from Mr. Richard H. Fletcher."

"Muddy Creek Cave.

"This cave was not visited by the writer. It lies about one mile north of Alderson on the road to Blue Sulphur Springs. A specimen of *Eurycea lucifuga* from the cave was secured through Mr. Richard H. Fletcher."

Subsurface Drainage.—No tests were made by the Survey to determine the outlets of the various streams that sink into the limestone but the structural position of the rocks and field data suggest the following:

Stream	Probable Point of Emergence of Streams
Sinking Creek.....	Piercys Mill
Milligan Creek.....	0.6 mile N. W. of Fort Spring
Culverson Creek.....	Tributary to Spring Creek (?)
Roaring Creek.....	0.7 mile N. of Sunlight
Buekeye Creek.....	Tributary to Spring Creek (?)

From a structural standpoint it is possible that Culverson Creek and Buekeye Creek flow southwest on their subsurface course and emerge either on Mill Creek or near Fort Spring.

the original substances. Water permeating the pores of rocks dissolves and removes any soluble substance originally present, as well as those formed by the chemical action of oxygen or carbon dioxide. Thus, the grains of substances neither subject to chemical change nor appreciably soluble in water are separated from one another in so finely divided a state that running water can easily carry them away. Sand, for example, is formed in this way from granites and from sandstones. The sand grains originally present in these rocks are simply left separated one from another through the removal of the other materials that, with the sand grains, compose such rocks.

Effects of Changes of Temperature.—Changes of temperature, especially in the Temperate Zones, are very active in breaking rocks to pieces, thus exposing fresh surfaces to the action of air and water. All substances change in volume with changes of temperature, and the change is nearly invariably expansion with rise in temperature. Since each of the several minerals of which rocks are composed has its own rate of change of volume with temperature, the result of considerable temperature change in a mass of rock is generally a weakening of the adhesion of unlike minerals to one another.

Another powerful disintegrating agent is the freezing of water which has been absorbed into the pores of the rock. As is well known, when water freezes the volume change is a decided **expansion**. Just as water freezing in pipes bursts them, so freezing in crevices of rocks pushes the pieces farther apart, while the freezing in the very small pores within the rock tends to break down the entire mass into a pile of mineral fragments. In this area we do not see piles of minerals so produced because abundant rainfall carries away the products of disintegration as fast as they are produced.

The Processes of Erosion and Deposition Never Cease.—The processes of the removal of material from the higher portions of the earth's surface and its deposition in the lower

PART II.

Geology.

CHAPTER III.

GEOLOGIC PROCESSES: EROSION AND DEPOSITION.

Hills and Valleys are Temporary Features.—When we look at the hills and valleys of our State, and think of the fact that the first inhabitants of this region, probably several thousand years ago, saw the same hills and valleys practically as they are to-day, it is hard to realize that they are, after all, quite temporary features—that there was a time in the earth's history before they existed, and that in the future they must surely vanish. Yet, whenever we see a stream flowing turbid with suspended matter after a rain, we have before us the process through which the valleys were made, leaving the hills as temporary remnants of the formerly continuous beds of rocks. And by this one process the hills too will, in time, be worn away and the materials of which they are composed carried seaward, finally to rest, in the case of material from most of our State, in the growing delta at the mouth of the Mississippi.

Weathering is a process of physical and chemical change which goes on whenever rocks are exposed to air, moisture, and changes of temperature. The active agents contained in air—oxygen and carbon dioxide—attack certain compounds and change them to other compounds.

throughout the millions of years of geological history. There is no area of the earth's surface that remains quite unaffected by these processes for any considerable length of time. How is it then, that the higher parts of the earth have not, long ago, been worn away entirely? Since the oceanic basins are larger than the land areas, to have this cycle go on to completion would mean that the earth would be entirely covered with water. This would certainly have happened long ago if the outer zone of the earth, (which we call the "crust" of the earth, because it was once thought that all of the earth within this zone was liquid), were stationary. Just as surely as these weathering processes with the aid of running water are trying to remove the irregularities of the surface of the earth internal processes or forces are tending to prevent it. We know that vast masses of this outer zone of the earth have moved upward even as far as several miles, while other masses have sunk downward. This fact is not so immediately evident as is that of the erosion processes just stated. Remains of sea animals, shells, corals, teeth and spines of marine fishes are found in many beds of rock now thousands of feet above sea-level. As a matter of fact all of these have been found in the rocks of Greenbrier County. The Greenbrier Limestone which is so conspicuous along the Greenbrier Valley contains literally millions of beautifully preserved marine shells and corals, while in the western part of the county fossil fish teeth are found a few feet above the Sewell Coal.

The processes of sinking and of elevation have actually been observed in many parts of the world. For hundreds of years parts of Denmark and Sweden have been slowly rising. On the other hand a part of the coast of Greenland has been sinking at the rate of several feet a century since the first settlement of that coast by Europeans.

Not only are portions of the outer zone of the earth elevated or depressed, but they are often times deformed into large or small arches, such as may be seen in the county (Alvon) and particularly in Peudleton and other counties of the Eastern Penhandle. Sometimes the stress on the beds of

opposite the fracture have slid past one another. Since we will frequently have occasion to speak of the features just mentioned these terms will be defined here:

Anticline.—A fold that is arched upward or convex upward. The oldest beds are in the middle.

Syncline.—A fold that is arched downward or convex downward. The youngest rocks are in the middle.

Fault.—A fracture or break along which there has been movement. The masses on opposite sides have moved past one another.

It can be seen that while a land area remains, as a whole, higher than the surrounding districts, not only will no new deposits (except volcanic) be laid down upon it, but the deposits already present will be continuously worn away. Now, the area of the State has, for a very long time, remained at least as high as any neighboring region. For this reason no very young rocks are found in Greenbrier County, or even in West Virginia, and many of the older rocks have been removed in places.

Classification of Rocks.—The rocks of the earth's crust fall into three main groups:—igneous, metamorphic, and sedimentary. **Igneous** rocks are those that have solidified from a molten magma. **Metamorphic** rocks are those that have been subjected to such intense heat and pressure that their physical and chemical properties have been changed. **Sedimentary** rocks are made up of the transported products of decomposition of older rocks or of organic material.

It is important to remember that all of the outcropping rocks in Greenbrier County are sedimentary rocks.

How Sediments Change to Stone.—As sediment is deposited, whether under water or on land, the lower beds become subject to an ever increasing pressure, due to the weight of the sediments that are constantly being laid down upon these lower beds. This slowly forces the particles of which the lower beds are composed closer together, besides flattening all particles of softer material. As the depth to which the lower beds are buried increases with deposition of new sediment, the temperature to which they are subjected also

is deposited, the temperature of the basal beds will be about 175 to 200° Fahrenheit. The pressure under the same thickness of sediments of average density will be in the neighborhood of 10,000 pounds per square inch. It must be remembered that beds of sediment are subjected to such pressures and temperatures, not for periods of time as we are well able to comprehend, but for periods of hundreds of thousands and millions of years. Under these conditions beds of soft clay and silt are changed into compact shales.

However, pressure and moderate heat alone are entirely ineffective in changing beds of reasonably pure quartz sand to solid sandstone. This takes place only through the deposition of some kind of cementing material,—usually from circulating water,—among the sand grains. The more important of these cementing materials are **calcium carbonate**, **ferric oxide**, and **silica**. Calcium carbonate (CaCO_3), is the chief constituent in ordinary limestone and is soluble in slightly acid water. Ferric oxide (Fe_2O_3) is more familiar to us as iron rust ($\text{Fe}_2\text{O}_3 \cdot n\text{H}_2\text{O}$) and when found in large quantities the minerals, limonite ($\text{Fe}_2\text{O}_3 \cdot n\text{H}_2\text{O}$) and hematite (Fe_2O_3) are valuable iron ores. Silica (SiO_2) is simply the material (quartz) of the grains themselves. Although practically insoluble in cold water it is soluble in hot water which already has certain substances in solution.

Limestone may be deposited as a mass of shell fragments, as a fine-grained lime mud, or as a mixture of these components. In either case it is readily and rapidly consolidated through formation of crystals of calcite, and through the effect of high pressure.

The Sedimentary Rocks.—All rocks thus formed through compaction and cementation of sediments under conditions of moderate temperatures and comparatively moderate pressures are called **sedimentary rocks**. The main classes into which they are divided are as follows:

Conglomerates are sedimentary rocks composed largely of pebbles and boulders, that is, of fragments larger than coarse

Meadow River at Nallen, W. Va.

Location.—Chain gage at highway bridge at Nallen, Fayette County.

Drainage area.—297 square miles.

Records available.—July, 1908, to September, 1916; November, 1928, to September, 1931.

Extremes.—Maximum discharge during year, 3,970 second-feet Apr. 4 (gage height, 10.70 feet); no flow Oct. 1-23, Oct. 27 to Nov. 5.

1908-1910, 1928-1931: Maximum discharge, about 7,300 second-feet Feb. 3,

1915 (gage height 13.25 feet); practically no flow at times in 1930.

Remarks.—Records good.

Daily and monthly discharge, in second-feet, 1930-31.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	0	0	4.6	69	308	237	1,350	375	340	35	179	190
2.....	0	0	4.6	40	237	530	1,460	324	340	29	141	150
3.....	0	0	6.4	45	179	610	1,200	278	278	50	2,940	190
4.....	0	0	25	40	150	510	3,610	250	212	88	1,980	200
5.....	0	0	26	56	108	410	3,520	212	308	70	1,840	159
6.....	0	.1	23	1,300	100	324	2,700	190	392	83	1,460	116
7.....	0	.1	54	1,100	124	293	1,910	179	375	132	1,010	94
8.....	0	.1	141	740	132	392	1,580	1,060	450	124	570	66
9.....	0	.1	100	530	324	550	1,910	1,350	470	169	375	54
10.....	0	.2	61	375	1,250	400	1,840	1,100	430	159	200	48
11.....	0	.2	45	212	965	350	2,140	875	324	179	212	44
12.....	0	.2	40	190	650	320	1,910	650	237	159	200	73
13.....	0	.2	35	159	510	300	1,350	740	190	73	530	37
14.....	0	.2	40	124	510	570	1,010	1,100	179	49	410	33
15.....	0	.2	37	108	650	1,770	785	1,150	490	34	308	33
16.....	0	.2	30	88	610	1,910	570	875	1,200	23	212	70
17.....	0	.3	24	88	650	965	430	695	610	33	132	150
18.....	0	.4	23	100	920	1,150	358	570	570	37	108	169
19.....	0	.4	21	141	1,200	1,010	308	610	324	32	88	108
20.....	0	.3	17	324	1,060	965	250	1,000	224	66	250	132
21.....	0	.3	15	358	830	830	212	1,640	179	62	1,520	610
22.....	0	.3	16	308	650	830	212	1,770	141	48	1,400	430
23.....	0	.3	18	224	400	785	570	3,180	124	78	2,380	278
24.....	.1	.4	13	141	392	905	740	2,300	179	250	1,640	324
25.....	.1	.4	12	100	358	1,150	740	1,520	132	430	1,150	308
26.....	.1	.5	12	150	308	1,100	695	1,150	100	308	785	470
27.....	0	.6	24	293	250	1,010	740	875	79	190	530	1,520
28.....	0	.7	94	650	224	1,150	695	650	63	150	375	1,400
29.....	0	.9	159	610	2,700	570	430	54	88	308	830
30.....	0	1.4	104	490	2,300	470	358	47	72	278	610
31.....	0	83	392	1,580	358	60	237

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
October	0.1	0	0.01	0.000034	0.00004
November	1.4	0	.30	.0010	.001
December	159	4.6	42.2	.142	.16
January	1,300	40	308	1.04	1.20
February	1,250	100	505	1.70	1.77
March	2,700	237	902	3.04	3.50
April	3,610	212	1,100	4.01	4.47
May	3,180	179	899	3.03	3.49
June	1,200	47	301	1.01	1.13
July	430	23	108	.364	.42
August	2,940	88	760	2.58	2.97
September	1,520	32	297	1.00	1.12

Meadow River at Nallen, W. Va.

Location.—Chain gage at highway bridge at Nallen, Fayette County.

Drainage area.—297 square miles.

Records available.—July, 1908, to September, 1916; November, 1928 to September, 1932.

Extremes.—Maximum discharge during year, 7,840 second-feet June 28 (gage height, 14.08 feet); minimum discharge, 0.4 second-foot Sept. 18, 19; minimum gage height, 2.58 feet Sept. 10.

1908-16, 1928-32: Maximum discharge, that of June 28, 1932; minimum discharge, less than 0.1 second-foot at times in 1930.

Remarks.—Records good. Discharge interpolated June 5.

Discharge, in second-feet, 1931-32.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	324	100	920	570	2,140	392	2,140	2,780	107	1,400	56	0.2
2.....	212	102	1,100	1,520	1,770	358	1,640	2,780	93	830	03	7.0
3.....	179	124	470	1,400	1,580	324	1,300	2,060	84	450	110	4.9
4.....	150	124	358	1,350	4,240	785	965	1,150	62	358	107	3.1
5.....	124	116	324	1,010	4,780	1,580	740	875	70	1,010	88	1.5
6.....	105	104	293	920	3,340	1,640	490	650	79	1,910	60	4.0
7.....	91	03	264	1,400	1,700	1,580	470	530	132	1,640	53	7.2
8.....	80	87	224	1,400	1,200	1,580	392	392	107	1,000	07	8.8
9.....	69	79	570	1,460	1,010	1,300	324	308	63	785	43	5.8
10.....	66	74	470	1,250	875	1,060	324	293	35	695	30	4.6
11.....	58	07	650	1,150	785	1,100	375	830	41	610	32	4.9
12.....	50	62	1,100	920	1,060	695	570	1,520	70	450	33	8.0
13.....	52	58	1,770	095	1,580	470	740	1,770	116	308	32	4.0
14.....	52	62	2,540	530	1,250	450	695	1,350	159	278	63	3.7
15.....	68	01	2,540	410	1,010	340	530	695	212	250	47	3.1
16.....	83	52	1,700	375	785	324	450	570	302	324	33	1.4
17.....	78	48	1,150	324	830	1,640	358	470	264	570	22	.6
18.....	68	52	785	308	920	3,430	324	392	141	392	31	.4
19.....	54	48	570	293	920	2,380	278	340	116	237	570	.4
20.....	47	48	510	293	830	1,400	250	293	124	141	278	1.0
21.....	43	40	610	278	650	1,200	224	237	278	124	169	1.0
22.....	41	44	830	204	610	1,060	293	785	392	141	100	.6
23.....	35	42	1,350	264	610	1,200	358	740	264	212	62	1.1
24.....	33	39	1,300	278	570	1,150	375	650	159	237	46	4.6
25.....	37	34	1,250	308	570	965	965	490	108	179	36	8.8
26.....	33	33	1,000	308	490	740	1,580	324	75	132	29	7.2
27.....	32	43	920	490	410	010	1,350	237	83	100	16	9.0
28.....	37	72	785	650	450	3,700	1,010	212	5,240	83	12	11
29.....	50	101	650	695	430	3,610	785	159	5,640	76	14	11
30.....	73	200	530	3,340	2,860	650	132	2,860	69	10	10
31.....	100	490	2,940	2,800	116	63	10

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
October	324	32	81.7	0.275	0.32
November	200	33	73.8	.248	.28
December	2,540	224	906	3.05	3.52
January	3,340	264	886	2.98	3.44
February	4,780	410	1,290	4.34	4.68
March	3,700	324	1,360	4.58	5.28
April	2,140	224	698	2.35	2.62
May	2,780	116	778	2.62	3.02
June	5,040	35	586	1.97	2.20
July	1,910	63	488	1.04	1.80
August	570	10	75.6	.255	.29
September

Meadow River at Nallen, W. Va.

Location.—Chain gage at highway bridge at Nallen, Fayette County, a quarter of a mile below Youngs Creek.

Drainage area.—297 square miles.

Records available.—July, 1908, to September, 1916; November, 1928, to September, 1934.

Extremes.—Maximum discharge recorded during year, 8,740 second-feet Mar. 5 (gage height, 15.64 feet); minimum, 3.2 second-feet July 25 (gage height, 2.57 feet).
1908-16, 1928-34: Maximum discharge recorded, that of Mar. 5, 1934; practically no flow at times in 1930.

Remarks.—Records good.

Discharge, in second-feet, 1933-34.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	33	24	108	010	212	264	890	212	53	11	30	14
2.....	30	23	101	1,250	237	293	685	100	51	9.8	34	12
3.....	24	21	104	1,100	212	3,260	472	159	85	0.3	48	9.8
4.....	17	19	108	875	190	5,740	403	150	73	7.8	264	9.0
5.....	15	17	110	695	169	8,540	354	141	67	6.7	179	8.6
6.....	14	86	050	695	169	6,840	323	132	56	6.0	00	8.1
7.....	13	79	965	920	169	4,870	293	124	46	4.7	64	7.6
8.....	12	84	785	1,400	150	3,340	308	116	100	4.0	48	7.1
9.....	12	83	740	1,200	141	3,100	203	124	68	6.4	34	6.4
10.....	10	70	650	065	132	2,140	293	132	60	7.8	28	5.8
11.....	8.8	72	570	740	132	1,460	278	141	49	9.0	25	5.1
12.....	13	62	490	530	141	990	250	150	44	9.8	22	7.4
13.....	13	73	264	430	141	840	237	159	42	12	19	8.3
14.....	12	86	212	410	159	665	200	150	35	15	16	8.6
15.....	11	93	200	358	141	625	224	200	27	14	15	9.8
16.....	16	107	237	308	141	386	338	370	21	12	46	11
17.....	53	116	410	278	169	508	386	403	17	9.8	88	33
18.....	101	116	570	224	190	490	437	354	10	8.1	108	73
19.....	95	116	610	212	224	472	750	278	20	0.7	82	48
20.....	79	124	3,430	200	250	545	890	212	42	0.0	48	31
21.....	53	150	2,380	179	224	508	090	169	04	5.3	36	24
22.....	40	179	1,400	160	190	454	940	150	40	4.8	27	19
23.....	30	212	1,100	169	179	420	890	124	32	4.5	24	16
24.....	33	224	740	169	160	370	750	116	26	3.9	22	14
25.....	33	224	570	159	179	338	585	108	22	4.3	27	12
26.....	29	212	470	150	224	370	545	101	20	0.2	27	12
27.....	26	190	410	159	250	1,340	437	88	17	12	24	14
28.....	22	169	324	169	278	2,060	278	73	11	53	22	15
29.....	19	150	308	124	1,910	250	68	10	64	20	19
30.....	19	110	264	58	1,520	237	61	9.8	58	18	124
31.....	22	264	212	1,190	55	36	16

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
October	101	8.8	29.5	0.099	0.11
November	224	17	110	.370	.41
December	3,430	101	631	2.12	2.44
January	1,400	58	487	1.64	1.80
February	278	132	184	.620	.65
March	8,540	264	1,802	6.07	7.00
April	990	200	470	1.58	1.76
May	403	55	162	.545	.63
June	100	9.8	40.8	.137	.15
July	64	3.9	13.8	.046	.05
August	264	15	50.0	.168	.19

Meadow River at Nallen, W. Va.

Location.—Chain gage at highway bridge at Nallen, Fayette County.

Drainage area.—297 square miles.

Records available.—July, 1908, to September, 1916; November, 1928 to September, 1933.

Discharge.—Maximum during the year, 3,970 second-feet Mar. 20 (gage height, 10.07 feet); minimum, 6.0 second-feet Oct. 3 (gage height, 2.90 feet).

1908-16, 1928-33: Maximum, about 7,840 second-feet June 23, 1932 (gage height, 14.68 feet); practically no flow at times in 1930.

Remarks.—Records good. Discharge estimated Apr. 3-10.

Discharge, in second-feet, 1932-33.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	8.8	278	141	1,150	920	740	490	108	400	160	510	95
2.....	7.2	785	141	1,150	1,250	570	610	159	340	132	430	74
3.....	6.4	610	132	875	065	470	1,000	450	204	308	740	07
4.....	6.8	470	124	740	020	375	1,240	785	212	570	1,350	81
5.....	11	358	116	610	875	308	880	1,350	160	324	1,200	109
6.....	15	204	108	490	785	264	700	1,010	124	200	1,000	109
7.....	20	212	108	410	785	264	640	1,640	110	141	830	124
8.....	26	170	105	340	2,700	570	780	1,350	150	102	308	94
9.....	20	324	97	510	3,100	920	1,120	1,400	250	80	237	74
10.....	34	1,980	93	875	1,640	020	920	1,910	308	63	190	61
11.....	31	1,400	94	875	1,150	830	610	2,000	224	159	237	49
12.....	21	1,200	116	830	875	920	605	1,770	100	124	212	44
13.....	14	875	293	785	740	740	020	1,520	141	132	169	42
14.....	12	570	430	570	740	1,010	1,150	1,300	93	86	141	49
15.....	9.2	302	400	510	1,100	3,340	965	1,250	74	01	110	224
16.....	11	278	005	450	1,700	2,540	920	1,520	62	64	110	570
17.....	94	278	695	340	1,400	1,770	830	1,640	56	40	150	358
18.....	605	264	740	324	1,250	1,640	740	1,350	49	39	100	250
19.....	010	204	050	392	1,060	3,340	650	1,200	41	33	160	141
20.....	740	1,400	510	1,100	2,380	3,610	610	1,100	30	28	124	124
21.....	650	1,200	358	2,380	2,860	3,700	530	020	24	22	108	116
22.....	375	1,055	250	3,610	2,060	3,020	450	650	21	22	88	102
23.....	250	785	340	2,860	1,400	1,980	375	530	18	45	81	78
24.....	124	570	570	1,770	1,010	1,520	324	470	13	32	86	64
25.....	94	302	740	1,300	830	1,150	308	302	21	35	87	54
26.....	70	308	375	1,520	920	695	264	410	42	110	83	43
27.....	64	250	875	1,700	1,000	570	237	510	278	050	64	36
28.....	212	212	1,640	1,640	875	570	212	490	278	1,300	53	43
29.....	278	200	2,220	1,350	530	200	470	237	1,770	64	46
30.....	237	100	1,010	1,060	510	100	510	200	1,300	159	43
31.....	190	1,460	920	470	570	965	150

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
October	740	0.4	100	0.538	0.62
November	1,980	179	585	1.07	2.20
December	2,220	93	552	1.86	2.14
January	3,010	324	1,080	3.64	4.20
February	3,100	740	1,330	4.48	4.06
March	3,700	264	1,290	4.34	5.00
April	1,240	190	652	2.20	2.46
May	2,000	108	1,020	3.43	3.05
June	490	13	150	.505	.56
July	1,770	22	294	.989	1.14
August	1,350	53	306	1.03	1.10
.....

Location.—Chain gage, lat. 35 55 30 N., long. 83 55 30 W.

Fayette County.

Drainage area.—287 square miles (revised).

Records available.—July, 1908, to September, 1916; November, 1928, to September, 1935.

Extremes.—Maximum discharge observed during year, about 5,940 second-feet Apr. 1 (gage height, 12.84 feet); minimum, 16 second-feet Sept. 30 (gage height, 2.99 feet.)

1908-16, 1928-35: Maximum discharge observed, about 8,740 second-feet Mar. 5, 1934, (gage height, 15.64 feet); practically no flow at times in 1930.

Remarks.—Records good.

Rating tables, water year 1934-35 (gage height, in feet, and discharge, in second-feet)

Table for Oct. 1 to Mar. 12

3.1	10	5.0	338
3.2	24	6.0	706
3.4	39	7.0	1,190
3.6	58	8.0	1,770
3.8	81	9.0	2,540
4.0	108	11.0	4,240
4.5	200		

Table for Mar. 13 to Sept. 30

3.0	16	5.0	805
3.2	20	6.0	780
3.4	46	7.0	1,225
3.6	67	8.0	1,770
3.8	91	9.0	2,540
4.0	120	11.0	4,240
4.5	216	13.0	6,140

Discharge, in second-feet, water year October, 1934, to September, 1935

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	370	58	2,060	420	490	940	5,549	104	547	330	128	26
2.....	212	169	1,010	585	437	750	5,140	184	402	330	128	26
3.....	132	169	1,400	545	370	585	3,020	208	314	314	120	38
4.....	97	141	1,090	545	323	472	1,840	610	912	298	87	55
5.....	70	200	840	403	278	403	1,420	610	1,770	736	71	267
6.....	74	420	665	354	250	472	1,270	780	1,140	824	59	956
7.....	293	585	508	293	200	472	1,420	3,700	736	868	384	780
8.....	264	437	403	264	190	437	1,470	4,240	505	1,090	2,700	547
9.....	200	204	308	386	212	380	1,420	2,140	484	1,040	1,840	384
10.....	141	224	264	1,040	940	323	1,270	1,320	443	912	1,140	228
11.....	100	160	237	990	1,190	1,040	1,090	912	330	526	736	173
12.....	79	150	200	840	990	4,060	1,040	604	240	282	384	128
13.....	62	132	190	665	795	3,700	1,180	568	267	173	228	103
14.....	50	116	169	400	706	2,620	1,379	662	205	128	956	90
15.....	43	195	150	386	1,240	1,640	1,270	912	194	173	824	73
16.....	36	94	124	323	1,240	1,270	1,140	1,000	228	205	402	63
17.....	33	82	124	1,840	1,140	1,090	868	1,140	104	128	254	62
18.....	30	79	132	2,220	940	868	780	1,000	194	95	173	42
19.....	26	76	159	1,709	706	736	736	956	184	66	128	40
20.....	24	71	338	1,340	545	1,220	652	736	194	57	99	35
21.....	24	67	472	1,520	437	1,370	568	868	154	69	92	36
22.....	23	65	472	2,460	386	1,320	484	1,040	206	101	87	32
23.....	22	169	386	4,060	403	2,460	402	1,000	205	145	70	29
24.....	21	990	308	3,610	625	4,330	348	1,140	205	145	96	28
25.....	21	705	278	2,140	625	3,790	282	1,840	136	240	57	26
26.....	24	508	338	1,290	640	3,790	254	1,520	104	610	46	23
27.....	24	403	1,140	840	1,290	2,620	210	1,140	89	547	42	20
28.....	23	278	000	790	1,090	1,640	205	824	79	422	49	17
29.....	22	625	795	730		1,420	104	610	72	267	37	17
30.....	24	1,460	665	680		1,189	194	694	72	164	35	16
31.....	23		499	625		3,920		736		128	32	

Month	Second-foot-days	Maximum	Minimum	Mean	Per square mile	Run-off in inches
October	2,587	370	21	83.5	0.291	0.34
November	9,011	1,469	65	300	1.05	1.17
December	17,605	2,069	124	568	1.98	2.22
Calendar year 1934.....	128,231.3	8,549	3.9	351	1.22	16.11
January	34,374	4,060	264	1,109	3.86	4.4
February	18,876	1,299	190	674	2.35	2.4
March	50,514	4,330	323	1,629	5.68	6.5
April	37,083	5,540	194	1,236	4.31	4.8
May	34,148	4,240	184	1,102	3.84	4.4
June	10,894	1,770	72	360	1.25	1.4
July	11,413	1,090	57	368	1.28	1.4
August	11,454	2,700	32	369	1.20	1.4
September	4,350	956	16	146	.506	.5

Location.—Gage, at highway bridge at Nallen, Fayette County.
 Drainage area.—287 square miles (revised).
 Records available.—July, 1908, to September, 1916; November, 1928, to September, 1935.
 Extremes.—Maximum discharge observed during year, about 5,940 second-feet Apr. 1 (gage height, 12.84 feet); minimum, 16 second-feet Sept. 30 (gage height, 2.99 feet.)
 1908-16, 1928-35: Maximum discharge observed, about 8,740 second-feet Mar. 5, 1934, (gage height, 15.64 feet); practically no flow at times in 1930.
 Remarks.—Records good.

Rating tables, water year 1934-35 (gage height, in feet, and discharge, in second-feet)
 Table for Oct. 1 to Mar. 12 Table for Mar. 13 to Sept. 30

3.1	10	5.0	338	3.0	16	5.0	365
3.2	24	6.0	706	3.2	29	6.0	780
3.4	39	7.0	1,100	3.4	46	7.0	1,225
3.6	58	8.0	1,770	3.6	67	8.0	1,770
3.8	81	9.0	2,540	3.8	91	9.0	2,540
4.0	108	11.0	4,240	4.0	120	11.0	4,240
4.5	200			4.5	216	13.0	6,140

Discharge, in second-feet, water year October, 1934, to September, 1935

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	370	58	2,060	420	490	940	5,540	104	547	330	128	26
2.....	212	169	1,910	585	437	750	5,140	184	402	339	128	26
3.....	132	169	1,400	545	370	585	3,020	298	314	314	120	38
4.....	07	141	1,090	545	823	472	1,840	610	912	298	87	55
5.....	70	200	840	403	278	403	1,420	610	1,770	736	71	207
6.....	74	420	665	354	250	472	1,270	780	1,140	824	59	056
7.....	293	585	508	293	200	472	1,420	3,700	736	808	384	780
8.....	264	437	403	264	190	437	1,479	4,240	505	1,990	2,700	547
9.....	200	264	308	380	212	386	1,420	2,140	484	1,040	1,840	384
10.....	141	224	264	1,040	940	323	1,270	1,320	443	912	1,140	228
11.....	100	169	237	990	1,190	1,040	1,090	912	330	526	736	173
12.....	79	150	290	840	900	4,060	1,040	694	240	282	384	128
13.....	62	132	190	665	795	3,790	1,180	568	267	173	228	103
14.....	50	110	169	490	705	2,620	1,370	652	205	128	956	90
15.....	43	105	159	386	1,240	1,640	1,270	912	194	173	824	73
16.....	36	94	124	323	1,240	1,270	1,140	1,000	228	206	402	63
17.....	33	82	124	1,840	1,140	1,090	868	1,140	104	128	254	52
18.....	30	79	132	2,220	040	868	780	1,090	194	95	173	42
19.....	26	76	159	1,700	705	736	736	956	184	66	128	40
20.....	24	71	338	1,340	545	1,220	652	736	194	57	99	35
21.....	24	67	472	1,520	437	1,370	568	868	154	69	92	36
22.....	23	65	472	2,460	386	1,320	484	1,040	205	101	87	32
23.....	22	169	386	4,060	403	2,460	402	1,000	205	145	79	29
24.....	21	990	308	3,610	625	4,330	348	1,140	205	145	06	28
25.....	21	705	278	2,149	625	3,790	282	1,840	130	240	57	26
26.....	24	508	338	1,290	840	3,790	254	1,520	104	610	46	23
27.....	24	403	1,140	840	1,290	2,620	216	1,140	89	547	42	20
28.....	23	278	990	790	1,090	1,040	205	824	79	422	40	17
29.....	22	625	795	730	1,420	194	610	72	267	37	17
30.....	24	1,460	665	680	1,180	194	694	72	164	35	16
31.....	23	490	625	3,020	736	128	32

Month	Second-foot-days	Maximum	Minimum	Mean	Per square mile	Run-off in inches
October	2,587	370	21	83.5	0.201	0.34
November	9,011	1,460	65	390	1.05	1.17
December	17,605	2,960	124	568	1.98	2.28
Calendar year 1934.....	128,231.3	8,540	3.9	351	1.22	16.18
January	34,374	4,000	264	1,100	3.86	4.45
February	18,876	1,290	190	674	2.35	2.45
March	50,514	4,330	323	1,629	5.68	6.55
April	37,088	5,540	194	1,236	4.31	4.81
May	34,148	4,240	184	1,102	3.84	4.43
June	10,804	1,779	72	369	1.25	1.40
July	11,413	1,090	57	368	1.28	1.48
August	11,454	2,700	32	369	1.29	1.49
September	4,350	956	16	145	5.05	5.0

Cherry River. The tributaries of Cherry River, Laurel Creek, Little Laurel Creek, South Fork of Cherry River and North Fork of Cherry River, drain northern Greenbrier County. A gaging station on this river was established at Richwood, Nicholas County, July 3, 1908, and records are available from that date to September 30, 1916, when the station was discontinued. Another gaging station was established on this river at Fenwick, Nicholas County, September, 1929, and records are available for this station to September, 1935. The following records of these stations were taken from the various Water-Supply Papers of the United States Geological Survey previously quoted under the description of Greenbrier River:

Cherry River at Richwood, W. Va.

Location.—At highway bridge at Richwood, Nicholas County, half a mile below junction of North and South Forks.

Drainage area.—90 square miles.

Records available.—July 3, 1908, to September 30, 1916, when station was discontinued.

Gage.—Chain gage on bridge; read by Floyd Artrip.

Discharge measurements.—Made from bridge or by wading.

Channel and control.—Channel straight above and below gage. Right bank subject to overflow and water passes around station at extremely high stages. Bed composed of gravel and boulders. Control practically permanent. The removal of stones from the control in 1909 and 1911 for building purposes changed the stage-discharge relation.

Ice.—Stage-discharge relation affected by ice for short periods in severe winters.

Extreme of discharge.—1908-1916: Maximum stage recorded, 9.0 feet October 1, 1915 (discharge, about 6,600 second-feet); minimum stage recorded, 1.66 feet July 1, 1914 (discharge, 5.2 second-feet); minimum discharge recorded, 4.8 second-feet October 8, 9, 1908 (gage height, 2.12 feet; before change in control).

Accuracy.—Stage-discharge relation practically permanent. Removal of stones for building purposes from primary control in July and August, 1909, and May to August, 1911, changed stage-discharge relation. Date of changes in stage-discharge relation not definitely known but are assumed to have occurred August 15, 1909, and June 30, 1911. Rating curve used July 3, 1908, to August 15, 1909, and curve used August 16, 1909, to June 30, 1911, are both based on only a few measurements and the form of all the rating curves; they are considered only fairly well defined. Rating curve used July 1, 1911, to September 30, 1916, is well defined to 2,400 second-feet and is an extension above that point. Gage read twice daily to half-tenths. Daily discharge ascertained by applying mean daily gage height to rating table except as noted in foot-note to table of daily discharge. Records fair, July 3, 1908, to August, 1911, except for July and August, 1909, and May to August, 1911, which are probably poor. Beginning with September, 1911, records good.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1008-9.												
1.....	15	120	42	368	340	327	220	525	109	308	28	20
2.....	12	100	32	222		408	308	465	100	500	32	20
3.....	8.7	82	28	184		025	280	337	98	209	22	16
4.....	8.6	40	35	150		625	251	273	132	203	15	16
5.....	8.0	39	32	1,140		558	436	230	132	138	14	216
6.....	8.0	35	30	970		525	525	184	214	408	12	76
7.....	8.0	30	30	800		525	387	157	123	465	22	36
8.....	4.8	28	286	890		465	290	308	163	280	14	82
9.....	4.8	25	264	300		660	247	203	120	203	6.4	35
10.....	6.8	24	243	268		816	199	590	247	157	8.0	142
11.....	02	25	222	235	377	1,220	176	558	200	114	8.0	173
12.....	37	46	263	280			157	372	214	92	0.0	01
13.....	23	30	184	222			203	281	170	126	6.0	00
14.....	15	20	167	184			1,740	222	150	126	6.0	46
15.....	14	15	144	590			735	181	222	87	106	43
16.....	12	22	132	890		1,220	366	150	188	08	223	39
17.....	11	22	222	590		600	286	126	170	64	123	75
18.....	11	27	735	525		436	230	103	356	55	61	55
19.....	10	55	660	525		495	102	82	211	46	82	40
20.....	9.4	82	525	408		772	486	184	78	153	32	64
21.....	0.4	82	398	380		406		250	150	123	28	50
22.....	0.4	57	299	351		465		590	243	100	27	46
23.....	11	55	214	361		398		660	123	05	28	49
24.....	04	48	150	328		308		525	98	273	73	30
25.....	68	42	150	305		308		436	98	273	04	32
26.....	63	30	184	282		366		290	308	222	46	32
27.....	23	37	184	233		327		243	350	157	42	26
28.....	23	37	184	184		337	810	346	268	144	80	22
29.....	98	33	120				405	268	207	157	57	20
30.....	192	42	184	340			398	337	157	273	42	20
31.....	132		406				264		126		57	20
1009-10.												
1.....	30	01	80	250	115	1,300	86	160	110	115	43	82
2.....	35	86	82	455	115	876	30	131	115	01	42	130
3.....	32	82	75	1,360	110	620	80	123	123	86	30	202
4.....	32	68	75	950	131	455	136	110	115	165	42	351
5.....	30	64	64	425	115	351	136	98	407	216	42	520
6.....	26	64	64	316	110	330	123	80	875	136	31	351
7.....	26	04	08	1,300	90	860	123	36	443	160	28	244
8.....	20	04	151	056	140	223	123	123	290	290	36	166
9.....	26	315	75	340		110	184	110	131	223	173	40
10.....	20	655	75	223		151	100	110	130	290	110	42
11.....	281	488	75	202		136	136	08	123	426	105	39
12.....	340	316	75	166			115	115	443	585	86	28
13.....	130	236	407	130			110	281	305	655	216	151
14.....	110	173	838	151			123	105	207	552	173	45
15.....	08	100	340	136			123	166	105	585	123	35
16.....	110	136	223	130		184	110	160	160	2,296	91	42
17.....	106	116	105	123		620	110	236	136	1,280	01	32
18.....	80	105	160	195	1,030	08	308	184	585	123	20	75
19.....	110	98	32	725	585	86	207	160	520	130	26	04
20.....	110	80		368	315	86	244	136	340	91	35	04
21.....	61	86		1,190	267	123	290	210	336	75	26	55
22.....	36	80	125	585	520	110	330	196	443	68	80	40
23.....	123	98		368	305	110	725	160	267	50	43	43
24.....	281	115		267	267	110	055	151	184	46	32	35
25.....	105	91	128	105	267	110	468	106	131	35	26	875
26.....	184	86		160	202	110	868	142	105	106	26	455
27.....	173	86		184	184	98	260	115	131	105	20	223
28.....	160	82	100	166	875	08	267	105	351	86	26	346
29.....	142	32		142		08	223	01	223	04	20	105
30.....	123	98		142		86	184	36	151	64	18	166
31.....	110			131		86		86		60	16	

Discharge measurements of Cherry River at Richwood, W. Va., during the years 1908-1916.

Date.	Made by—	Gage height.	Discharge.	Date.	Made by—	Gage height.	Discharge.
1908.		Feet.	Sec.-ft.	1912.		Feet.	Sec.-ft.
July 2	Wm. M. O'Neill.....	2.54	54	Mar. 20	C. T. Bailey.....	4.35	1,270
28	W. G. Hoyt.....	4.24	970				
Sept. 26	Wm. M. O'Neill.....	2.25	8	1913.			
1909.				Dec. 2	Peterson and Walters	3.89	019
Mar. 28	H. J. Jackson.....	4.04	744	2do	3.85	894
29do	3.60	500				
Nov. 13	A. H. Horton.....	3.09	225	1914.			
1910.				Nov. 22	J. G. Mathers.....	2.20	05.0
Mar. 12do	2.67	188	22do	2.27	68.5
Aug. 10	J. C. Dort.....	2.27	41.4				
1911.				1916.			
Oct. 25	Bailey and Perwien..	2.74	295	Sept. 5	B. E. Jones.....	1.84	13.6
				5do	1.84	18.7

Daily discharge, in second-feet, of Cherry River at Richwood, W. Va., for the years ending Sept. 30, 1908-1916.

Day.	July.	Aug.	Sept.	Day.	July.	Aug.	Sept.	Day.	July.	Aug.	Sept.
1.....	58	120	46	11.....	120	89	15	21.....	109	37	6.0
2.....	54	106	46	12.....	02	85	15	22.....	808	37	12
3.....	46	02	28	13.....	98	87	15	23.....	350	37	15
4.....	57	80	28	14.....	57	92	15	24.....	525	28	15
5.....	184	92	37	15.....	57	68	12	25.....	419	89	8.0
6.....	480	150	37	10.....	80	68	8.0	26.....	273	294	9.4
7.....	299	120	28	17.....	73	46	8.0	27.....	1,309	120	8.7
8.....	222	92	22	18.....	123	57	8.0	28.....	979	92	9.4
9.....	192	382	15	19.....	177	46	8.0	29.....	465	08	18
10.....	159	203	15	20.....	126	46	8.0	30.....	286	57	18
								31.....	293	40

Daily discharge, in second-feet, of Cherry River at Richwood, W. Va., for the
years ending Sept. 30, 1908-1916—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1010-11.												
1.....	136	110	267	762	655	110	145	123	315	54	24	112
2.....	142	123	223	1,890	585	110	142	123	115	43	18	55
3.....	105	136	166	1,710	395	98	340	110	68	100	14	43
4.....	91	131	142	875	330	110	725	91	46	43	18	50
5.....	75	110	131	488	253	86	1,020	75	202	30	43	152
6.....	64	110	136	368	195	875	838	04	151	50	08	120
7.....	75	91	123	244	106	600	520	64	105	98	32	02
8.....	223	80	110	202	184	407	455	64	86	120	85	59
9.....	308	80	105	166	244	290	585	64	51	55	40	845
10.....	202	75	86	151	223	300	378	51	39	43	34	288
11.....	142	75	86	136	105	395	290	42	43	100	24	246
12.....	131	64	86	620	166	340	300	40	136	87	20	232
13.....	105	64	80	2,610	142	455	407	43	98	55	28	170
14.....	86	64		1,360	123	395	585	39	71	40	30	152
15.....	75	64	90	1,620	115	300	762	32	59	30	36	415
16.....	64	60		1,030	105	244	305	28	55	34	100	810
17.....	64	55	100	520	86	202	315	32	40	47	59	415
18.....	60	55	100	368	98	184	244	32	64	46	36	232
19.....	50	46	160	267	136	300	223	32	64	32	28	190
20.....	46	46	142	202	115	620	620	26	64	24	24	171
21.....	46	43		267	86	368	500	20	46	26	24	130
22.....	142	39	145	585	110	290	520	20	35	36	18	142
23.....	123	32		468	110	368	538	20	32	30	10	02
24.....	86	46	151	340	80	236	368	184	28	30	13	59
25.....	64	290	290	253	86	195	325	64	42	68	16	64
26.....	64	160	166	520	80	151	244	42	46	50	24	55
27.....	68	131	160	655	123	350	210	43	64	36	18	46
28.....	184	425	160	762	151	230	173	32	115	28	14	142
29.....	131	520	875	875		202	106	30	51	24	32	146
30.....	115	315	2,000	2,720		244	128	28	36	20	59	114
31.....	110		762	762		177		136		24	171	
1911-12.												
1.....	80	80	232	505	304	330	445	385	47	64	160	152
2.....	705	08	198	358	211	255	775	320	43	54	120	120
3.....	602	64	160	288	198	211	775	278	47	50	08	100
4.....	385	59	126	232	152	160	538	224	39	87	80	152
5.....	255	59	100	146	171	146	445	190	30	120	64	92
6.....	179	304	106		100	120	330	171	30	73	54	82
7.....	304	570	98		190	120	385	570	36	54	50	64
8.....	538	330	98	125		120	415	538	26	47	43	50
9.....	358	241	98			211	330	415	22	59	43	44
10.....	255	198	120			190	304	330	20	120	68	40
11.....	415	160	126		130	171	224	304	10	445	68	34
12.....	475	190	152	114		100	190	1,870	10	224	08	31
13.....	358	320	126	87		570	160	995	16	136	87	24
14.....	278	232	120	80		385	152	670	16	152	59	22
15.....	775	224	126	68	59	3,150	146	538	15	505	39	22
16.....	602	190	241	80	59	1,490	241	2,030	18	415	34	34
17.....	670	152	278		54	775	190	1,400	50	211	30	30
18.....	1,670	445	224	92	73	635	320	670	92	232	30	34
19.....	775	415	171	255	87	705	255	505	224	330	106	50
20.....	505	304	152	358	182	1,310	224	358	171	211	136	36
21.....	314	269	152	211	775	995	100	269	80	152	100	31
22.....	288	211	160	190	920	882	211	224	50	152	171	22
23.....	330	160	505	152	505	538	260	182	198	120	114	358
24.....	269	190	385	152	385	995	232	146	87	92	68	570
25.....	179	224	385	120	288	845	198	120	171	1,070	01	358
26.....	160	171	385	120	1,490	602	182	114	160	670	64	215
27.....	120	100	635	106	1,310	415	255	87	126	385	92	215
28.....	114	198	505	106	070	330	570	73	171	232	64	163
29.....	106	358	358	190	475	1,670	255	08	98	232	740	130
30.....	92	269	278	635		920	445	59	73	209	385	114
31.....								50		182	215	

Daily discharge, in second-feet, of Cherry River at Richwood, W. Va., for the
years ending Sept. 30, 1908-1916—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1912-13.												
1.....	92	87	54	358	358	445	299	314	602	215	49	39
2.....	82	100	73	288	320	358	250	278	415	90	90	29
3.....	75	82	241	358	415	260	194	219	278	385	61	29
4.....	64	08	228	358	705	215	182	171	278	250	44	29
5.....	55	64	358	304	475	207	156	142	190	117	39	23
6.....	55	04	475	740	330	182	156	130	152	845	31	19
7.....	50	1,490	445	1,070	269	123	139	136	264	309	29	21
8.....	47	1,070	358	1,970	283	123	123	106	358	175	29	31
9.....	44	570	241	995	237	133	109	92	304	126	23	40
10.....	40	358	198	635	175	175	109	80	219	605	23	29
11.....	34	273	179	538	299	810	114	68	179	260	21	19
12.....	34	211	126	635	538	570	330	68	142	198	54	10
13.....	31	179	112	505	330	475	330	59	112	149	215	19
14.....	36	198	126	385	283	1,670	309	59	92	117	126	19
15.....	34	171	112	304	215	920	1,490	63	80	330	61	19
16.....	34	136	98	255	175	670	705	100	08	241	54	15
17.....	28	120	85	211	156	445	475	358	59	149	42	52
18.....	28	112	85	108	123	320	358	294	50	126	29	106
19.....	163	112	90	179	114	250	304	211	43	1,400	186	52
20.....	176	106	06	100	175	215	246	103	40	810	241	40
21.....	109	98	63	228	283	207	190	179	43	358	117	845
22.....	87	86	60	219	445	182	171	171	50	215	475	385
23.....	385	86	80	211	415	146	142	602	106	167	740	182
24.....	190	73	63	505	320	130	136	958	63	107	273	114
25.....	176	73	73	538	215	139	120	505	59	385	175	75
26.....	163	63	98	445	207	250	112	358	358	198	133	59
27.....	163	63	85	385	269	2,600	358	1,400	142	139	00	55
28.....	146	59	85	283	635	920	358	1,400	106	103	66	04
29.....	130	40	90	224	602	385	705	75	06	57	40
30.....	114	54	475	182	445	475	538	59	06	49	52
31.....	92	570	224	358	1,310	57	42
1913-14.												
1.....	55	150	538	171	705	171	740	219	25	5.2	8.0	39
2.....	55	130	920	146	445	1,070	179	25	8.4	8.0	29
3.....	320	100	538	160	330	150	775	142	21	19	8.0	29
4.....	246	100	415	152	269	475	142	21	12	7.2	21
5.....	146	87	304	120	269	358	219	25	14	8.0	19
6.....	106	75	211	114	255	114	260	570	29	11	8.0	15
7.....	96	75	358	106	385	109	232	385	21	8.4	7.2	15
8.....	75	80	385	106	330	100	882	314	21	7.6	14	15
9.....	64	294	320	211	241	87	070	264	17	61	16	12
10.....	100	203	232	304	232	100	445	219	15	21	43	12
11.....	95	182	211	232	190	114	330	186	14	14	24	14
12.....	130	182	182	198	100	92	278	160	14	11	85	31
13.....	100	294	152	232	152	100	215	149	17	8.0	63	26
14.....	80	1,230	130	190	152	92	182	136	14	80	32	19
15.....	75	1,970	126	190	160	203	330	112	12	63	20	14
16.....	70	1,870	120	152	146	538	1,400	98	9.2	40	16	12
17.....	59	670	120	126	146	845	740	85	8.4	30	13	11
18.....	87	505	130	114	152	740	475	80	7.6	32	10	9.6
19.....	100	445	126	120	920	475	358	73	6.4	20	8.0	7.6
20.....	505	320	120	330	920	330	775	63	0.2	14	8.0	7.6
21.....	415	255	120	1,310	475	278	635	63	11	13	59	7.6
22.....	294	194	120	635	330	215	445	54	11	10	30	7.6
23.....	385	163	120	415	330	182	330	54	11	8.0	20	7.6
24.....	1,070	130	152	385	278	190	264	46	15	8.0	22	7.6
25.....	1,230	123	152	995	211	445	219	43	15	8.0	63	9.6
26.....	1,490	100	505	570	190	1,310	1,150	39	17	13	309	12
27.....	740	100	330	445	190	1,310	670	32	12	55	133	9.6
28.....	445	385	255	385	160	1,580	445	30	9.2	24	06	7.6
29.....	320	445	224	602	1,070	330	25	8.4	16	108	7.0
30.....	237	320	198	775	1,070	264	37	6.4	13	66	7.6
31.....	104	182	1,400	995	31	12	54

Monthly discharge of Cherry River at Richwood, W. Va., for the years ending
Sept. 30, 1908-1916.

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1908.					
July	1,300	46	258	2.87	3.31
August	382	28	96.1	1.07	1.23
September	46	6	17.8	.108	.22
1908-9.					
October	192	4.8	32.3	.359	.41
November	120	15	44.8	.498	.56
December	735	28	218	2.42	2.79
January	1,140	150	426	4.73	5.46
February	1,220	327	446	4.96	5.16
March	1,220	264	530	5.80	6.70
April	1,740	157	379	4.21	4.70
May	590	78	244	2.71	3.12
June	356	95	178	1.98	2.21
July	590	27	142	1.58	1.82
August	223	6.0	40.2	.447	.52
September	290	16	70.8	.787	.88
The year	1,740	4.8	228	2.53	34.41
1909-10.					
October	340	20	111	1.23	1.42
November	655	64	140	1.62	1.81
December	838	64	147	1.63	1.88
January	1,360	123	394	4.38	5.05
February	1,030	269	2.99	3.11
March	1,300	86	229	2.54	2.93
April	725	86	235	2.61	1.01
May	443	86	158	1.76	2.03
June	2,290	105	437	4.86	5.42
July	290	35	117	1.30	1.50
August	151	16	38.3	.426	.40
September	875	35	202	2.24	2.50
The year	2,290	16	206	2.20	31.05
1910-11.					
October	368	40	111	1.23	1.42
November	520	32	122	1.36	1.52
December	2,090	86	245	2.72	3.14
January	2,720	136	768	8.53	9.83
February	655	86	191	2.12	2.21
March	875	86	301	3.34	3.85
April	1,620	123	435	4.83	5.39
May	184	20	58.1	.646	.74
June	315	28	79.1	.870	.98
July	126	20	49.5	.550	.63
August	100	13	40.4	.440	.52
September	845	43	105	2.17	2.42
The year	2,720	13	217	2.41	32.65
1911-12.					
October	1,670	80	395	4.39	5.06
November	570	50	227	2.52	2.81
December	635	08	237	2.63	3.03
January	635	68	189	2.10	2.42
February	1,490	54	333	3.70	3.99
March	3,150	120	646	7.18	8.28
April	775	146	322	3.58	3.99
May	2,930	50	480	5.40	6.23
June	224	15	73.4	.816	.91
July	1,070	47	230	2.50	2.95
August	740	30	114	1.27	1.46
September	570	22	122	1.36	1.52

Years ending Sept. 30, 1908-1916—Continued.

Day.	1914.	1915.	1916.	1917.	1918.	1919.	1920.	1921.	1922.	1923.	1924.	1925.	1926.	1927.	1928.	1929.	1930.	1931.	1932.	1933.	1934.	1935.	1936.	1937.	1938.	1939.	1940.	1941.	1942.	1943.	1944.	1945.	1946.	1947.	1948.	1949.	1950.	1951.	1952.	1953.	1954.	1955.	1956.	1957.	1958.	1959.	1960.	1961.	1962.	1963.	1964.	1965.	1966.	1967.	1968.	1969.	1970.	1971.	1972.	1973.	1974.	1975.	1976.	1977.	1978.	1979.	1980.	1981.	1982.	1983.	1984.	1985.	1986.	1987.	1988.	1989.	1990.	1991.	1992.	1993.	1994.	1995.	1996.	1997.	1998.	1999.	2000.	2001.	2002.	2003.	2004.	2005.	2006.	2007.	2008.	2009.	2010.	2011.	2012.	2013.	2014.	2015.	2016.	2017.	2018.	2019.	2020.	2021.	2022.	2023.	2024.	2025.	2026.	2027.	2028.	2029.	2030.	2031.	2032.	2033.	2034.	2035.	2036.	2037.	2038.	2039.	2040.	2041.	2042.	2043.	2044.	2045.	2046.	2047.	2048.	2049.	2050.	2051.	2052.	2053.	2054.	2055.	2056.	2057.	2058.	2059.	2060.	2061.	2062.	2063.	2064.	2065.	2066.	2067.	2068.	2069.	2070.	2071.	2072.	2073.	2074.	2075.	2076.	2077.	2078.	2079.	2080.	2081.	2082.	2083.	2084.	2085.	2086.	2087.	2088.	2089.	2090.	2091.	2092.	2093.	2094.	2095.	2096.	2097.	2098.	2099.	2100.	2101.	2102.	2103.	2104.	2105.	2106.	2107.	2108.	2109.	2110.	2111.	2112.	2113.	2114.	2115.	2116.	2117.	2118.	2119.	2120.	2121.	2122.	2123.	2124.	2125.	2126.	2127.	2128.	2129.	2130.	2131.	2132.	2133.	2134.	2135.	2136.	2137.	2138.	2139.	2140.	2141.	2142.	2143.	2144.	2145.	2146.	2147.	2148.	2149.	2150.	2151.	2152.	2153.	2154.	2155.	2156.	2157.	2158.	2159.	2160.	2161.	2162.	2163.	2164.	2165.	2166.	2167.	2168.	2169.	2170.	2171.	2172.	2173.	2174.	2175.	2176.	2177.	2178.	2179.	2180.	2181.	2182.	2183.	2184.	2185.	2186.	2187.	2188.	2189.	2190.	2191.	2192.	2193.	2194.	2195.	2196.	2197.	2198.	2199.	2200.	2201.	2202.	2203.	2204.	2205.	2206.	2207.	2208.	2209.	2210.	2211.	2212.	2213.	2214.	2215.	2216.	2217.	2218.	2219.	2220.	2221.	2222.	2223.	2224.	2225.	2226.	2227.	2228.	2229.	2230.	2231.	2232.	2233.	2234.	2235.	2236.	2237.	2238.	2239.	2240.	2241.	2242.	2243.	2244.	2245.	2246.	2247.	2248.	2249.	2250.	2251.	2252.	2253.	2254.	2255.	2256.	2257.	2258.	2259.	2260.	2261.	2262.	2263.	2264.	2265.	2266.	2267.	2268.	2269.	2270.	2271.	2272.	2273.	2274.	2275.	2276.	2277.	2278.	2279.	2280.	2281.	2282.	2283.	2284.	2285.	2286.	2287.	2288.	2289.	2290.	2291.	2292.	2293.	2294.	2295.	2296.	2297.	2298.	2299.	2300.	2301.	2302.	2303.	2304.	2305.	2306.	2307.	2308.	2309.	2310.	2311.	2312.	2313.	2314.	2315.	2316.	2317.	2318.	2319.	2320.	2321.	2322.	2323.	2324.	2325.	2326.	2327.	2328.	2329.	2330.	2331.	2332.	2333.	2334.	2335.	2336.	2337.	2338.	2339.	2340.	2341.	2342.	2343.	2344.	2345.	2346.	2347.	2348.	2349.	2350.	2351.	2352.	2353.	2354.	2355.	2356.	2357.	2358.	2359.	2360.	2361.	2362.	2363.	2364.	2365.	2366.	2367.	2368.	2369.	2370.	2371.	2372.	2373.	2374.	2375.	2376.	2377.	2378.	2379.	2380.	2381.	2382.	2383.	2384.	2385.	2386.	2387.	2388.	2389.	2390.	2391.	2392.	2393.	2394.	2395.	2396.	2397.	2398.	2399.	2400.	2401.	2402.	2403.	2404.	2405.	2406.	2407.	2408.	2409.	2410.	2411.	2412.	2413.	2414.	2415.	2416.	2417.	2418.	2419.	2420.	2421.	2422.	2423.	2424.	2425.	2426.	2427.	2428.	2429.	2430.	2431.	2432.	2433.	2434.	2435.	2436.	2437.	2438.	2439.	2440.	2441.	2442.	2443.	2444.	2445.	2446.	2447.	2448.	2449.	2450.	2451.	2452.	2453.	2454.	2455.	2456.	2457.	2458.	2459.	2460.	2461.	2462.	2463.	2464.	2465.	2466.	2467.	2468.	2469.	2470.	2471.	2472.	2473.	2474.	2475.	2476.	2477.	2478.	2479.	2480.	2481.	2482.	2483.	2484.	2485.	2486.	2487.	2488.	2489.	2490.	2491.	2492.	2493.	2494.	2495.	2496.	2497.	2498.	2499.	2500.	2501.	2502.	2503.	2504.	2505.	2506.	2507.	2508.	2509.	2510.	2511.	2512.	2513.	2514.	2515.	2516.	2517.	2518.	2519.	2520.	2521.	2522.	2523.	2524.	2525.	2526.	2527.	2528.	2529.	2530.	2531.	2532.	2533.	2534.	2535.	2536.	2537.	2538.	2539.	2540.	2541.	2542.	2543.	2544.	2545.	2546.	2547.	2548.	2549.	2550.	2551.	2552.	2553.	2554.	2555.	2556.	2557.	2558.	2559.	2560.	2561.	2562.	2563.	2564.	2565.	2566.	2567.	2568.	2569.	2570.	2571.	2572.	2573.	2574.	2575.	2576.	2577.	2578.	2579.	2580.	2581.	2582.	2583.	2584.	2585.	2586.	2587.	2588.	2589.	2590.	2591.	2592.	2593.	2594.	2595.	2596.	2597.	2598.	2599.	2600.	2601.	2602.	2603.	2604.	2605.	2606.	2607.	2608.	2609.	2610.	2611.	2612.	2613.	2614.	2615.	2616.	2617.	2618.	2619.	2620.	2621.	2622.	2623.	2624.	2625.	2626.	2627.	2628.	2629.	2630.	2631.	2632.	2633.	2634.	2635.	2636.	2637.	2638.	2639.	2640.	2641.	2642.	2643.	2644.	2645.	2646.	2647.	2648.	2649.	2650.	2651.	2652.	2653.	2654.	2655.	2656.	2657.	2658.	2659.	2660.	2661.	2662.	2663.	2664.	2665.	2666.	2667.	2668.	2669.	2670.	2671.	2672.	2673.	2674.	2675.	2676.	2677.	2678.	2679.	2680.	2681.	2682.	2683.	2684.	2685.	2686.	2687.	2688.	2689.	2690.	2691.	2692.	2693.	2694.	2695.	2696.	2697.	2698.	2699.	2700.	2701.	2702.	2703.	2704.	2705.	2706.	2707.	2708.	2709.	2710.	2711.	2712.	2713.	2714.	2715.	2716.	2717.	2718.	2719.	2720.	2721.	2722.	2723.	2724.	2725.	2726.	2727.	2728.	2729.	2730.	2731.	2732.	2733.	2734.	2735.	2736.	2737.	2738.	2739.	2740.	2741.	2742.	2743.	2744.	2745.	2746.	2747.	2748.	2749.	2750.	2751.	2752.	2753.	2754.	2755.	2756.	2757.	2758.	2759.	2760.	2761.	2762.	2763.	2764.	2765.	2766.	2767.	2768.	2769.	2770.	2771.	2772.	2773.	2774.	2775.	2776.	2777.	2778.	2779.	2780.	2781.	2782.	2783.	2784.	2785.	2786.	2787.	2788.	2789.	2790.	2791.	2792.	2793.	2794.	2795.	2796.	2797.	2798.	2799.	2800.	2801.	2802.	2803.	2804.	2805.	2806.	2807.	2808.	2809.	2810.	2811.	2812.	2813.	2814.	2815.	2816.	2817.	2818.	2819.	2820.	2821.	2822.	2823.	2824.	2825.	2826.	2827.	2828.	2829.	2830.	2831.	2832.	2833.	2834.	2835.	2836.	2837.	2838.	2839.	2840.	2841.	2842.	2843.	2844.	2845.	2846.	2847.	2848.	2849.	2850.	2851.	2852.	2853.	2854.	2855.	2856.	2857.	2858.	2859.	2860.	2861.	2862.	2863.	2864.	2865.	2866.	2867.	2868.	2869.	2870.	2871.	2872.	2873.	2874.	2875.	2876.	2877.	2878.	2879.	2880.	2881.	2882.	2883.	2884.	2885.	2886.	2887.	2888.	2889.	2890.	2891.	2892.	2893.	2894.	2895.	2896.	2897.	2898.	2899.	2900.	2901.	2902.	2903.	2904.	2905.	2906.	2907.	2908.	2909.	2910.	2911.	2912.	2913.	2914.	2915.	2916.	2917.	2918.	2919.	2920.	2921.	2922.	2923.	2924.	2925.	2926.	2927.	2928.	2929.	2930.	2931.	2932.	2933.	2934.	2935.	2936.	2937.	2938.	2939.	2940.	2941.	2942.	2943.	2944.	2945.	2946.	2947.	2948.	2949.	2950.	2951.	2952.	2953.	2954.	2955.	2956.	2957.	2958.	2959.	2960.	2961.	2962.	2963.	2964.	2965.	2966.	2967.	2968.	2969.	2970.	2971.	2972.	2973.	2974.	2975.	2976.	2977.	2978.	2979.	2980.	2981.	2982.	2983.	2984.	2985.	2986.	2987.	2988.	2989.	2990.	2991.	2992.	2993.	2994.	2995.	2996.	2997.	2998.	2999.	3000.	3001.	3002.	3003.	3004.	3005.	3006.	3007.	3008.	3009.	3010.	3011.	3012.	3013.	3014.	3015.	3016.	3017.	3018.	3019.	3020.	3021.	3022.	3023.	3024.	3025.	3026.	3027.	3028.	3029.	3030.	3031.	3032.	3033.	3034.	3035.	3036.	3037.	3038.	3039.	3040.	3041.	3042.	3043.	3044.	3045.	3046.	3047.	3048.	3049.	3050.	3051.	3052.	3053.	3054.	3055.	3056.	3057.	3058.	3059.	3060.	3061.	3062.	3063.	3064.	3065.	3066.	3067.	3068.	3069.	3070.	3071.	3072.	3073.	3074.	3075.	3076.	3077.	3078.	3079.	3080.	3081.	3082.	3083.	3084.	3085.	3086.	3087.	3088.	3089.	3090.	3091.	3092.	3093.	3094.	3095.	3096.	3097.	3098.	3099.	3100.	3101.	3102.	3103.	3104.	3105.	3106.	3107.	3108.	3109.	3110.	3111.	3112.	3113.	3114.	3115.	3116.	3117.	3118.	3119.	3120.	3121.	3122.	3123.	3124.	3125.	3126.	3127.	3128.	3129.	3130.	3131.	3132.	3133.	3134.	3135.	3136.	3137.	3138.	3139.	3140.	3141.	3142.	3143.	3144.	3145.	3146.	3147.	3148.	3149.	3150.	3151.	3152.	3153.	3154.	3155.	3156.	3157.	3158.	3159.	3160.	3161.	3162.	3163.	3164.	3165.	3166.	3167.	3168.	3169.	3170.	3171.	3172.	3173.	3174.	3175.	3176.	3177.	3178.	3179.	3180.	3181.	3182.	3183.	3184.	3185.	3186.	3187.	3188.	3189.	3190.	3191.	3192.	3193.	3194.	3195.	3196.	3197.	3198.	3199.	3200.	3201.	3202.	3203.	3204.	3205.	3206.	3207.	3208.	3209.	3210.	3211.	3212.	3213.	3214.	3215.	3216.	3217.	3218.	3219.	3220.	3221.	3222.	3223.	3224.	3225.	3226.	3227.	3228.	3229.	3230.	3231.	3232.	3233.	3234.	3235.	3236.	3237.	3238.	3239.	3240.	3241.	3242.	3243.	3244.	3245.	3246.	3247.	3248.	3249.	3250.	3251.	3252.	3253.	3254.	3255.	3256.	3257.	3258.	3259.	3260.	3261.	3262.	3263.	3264.	3265.	3266.	3267.	3268.	3269.	3270.	3271.	3272.	3273.	3274.	3275.	3276.	3277.	3278.	3279.	3280.	3281.	3282.	3283.	3284.	3285.	3286.	3287.	3288.
------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------

Monthly discharge of Cherry River at Richwood, W. Va., for the years ending
Sept. 30, 1908-1916—Continued.

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1912-13.					
October	385	28	95.3	1.06	1.22
November	1,490	49	209	2.32	2.59
December	570	54	177	1.07	2.27
January	1,970	160	477	5.30	6.11
February	705	114	313	3.48	3.62
March	2,600	123	469	5.21	6.01
April	1,490	109	294	3.27	3.65
May	1,490	59	366	4.07	4.69
June	602	40	166	1.84	2.05
July	1,490	57	284	3.16	3.64
August	740	21	118	1.31	1.51
September	845	15	84.3	.087	1.04
The year	2,600	15	255	2.83	38.40
1913-14.					
October	1,490	55	303	3.37	3.88
November	1,970	75	373	4.14	4.62
December	920	120	258	2.87	3.31
January	1,490	106	370	4.11	4.74
February	920	146	313	3.48	3.62
March	1,580	87	434	4.82	5.56
April	1,400	182	525	5.83	6.50
May	570	25	187	1.52	1.75
June	29	6.4	14.9	.166	.18
July	80	5.2	21.3	.237	.27
August	309	7.2	43.1	.479	.55
September	39	7.6	14.7	.103	.18
The year	1,970	5.2	233	2.59	35.16
1914-15.					
October	358	4.8	61.9	.688	.79
November	241	30	66.8	.742	.83
December	1,670	120	325	3.61	4.16
January	2,380	92	480	5.33	6.14
February	2,380	152	548	6.09	6.34
March	182	68	115	1.28	1.48
April	705	54	212	2.36	2.03
May	255	36	104	1.16	1.34
June	260	0	85.6	.951	1.06
July	475	26	110	1.22	1.41
August	635	24	124	1.38	1.59
September	314	15	91.5	1.02	1.14
The year	2,380	4.8	192	2.13	28.91
1915-16.					
October	4,330	.32	284	3.16	3.64
November	635	14	162	1.80	2.01
December	2,710	31	358	3.98	4.59
January	1,150	156	427	4.74	5.46
February	882	123	340	3.78	4.08
March	1,310	130	472	5.24	6.04
April	995	120	288	3.20	3.56
May	278	37	128	1.42	1.64
June	602	40	198	2.20	2.45
July	995	21	213	2.37	2.73
August	658	19	218	2.42	2.79
September	2,070	8	155	1.72	1.93
The year	4,330	8	271	3.01	40.92

Cherry River at Fenwick, W. Va.

Location.—Chain gage at highway bridge at Fenwick, Nicholas County, 1,000 feet below mouth of Laurel Creek.

Drainage area.—150 square miles.

Records available.—September, 1929, to September, 1930.

Extremes.—Maximum gage height during year, 12.04 feet Oct. 2 (discharge not determined); minimum, 0.1 second-foot Sept. 20 (gage height, 2.62 feet).

Remarks.—Records good.

Daily and monthly discharge, in second-feet, 1929-30.

Day.	Sept.	Oct.	Nov.	Dec.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....		33	190	163	242	125	42	5.8	2.0	2.4
2.....		6,120	178	158	455	125	32	5.8	1.8	1.7
3.....		3,510	885	150	960	58	34	5.5	.5	1.4
4.....		1,700	780	158	1,340	37	26	4.1	.5	1.1
5.....		920	780	181	850	95	21	2.8	.7	1.1
6.....		590	a 600	204	780	123	23	2.0	1.4	1.0
7.....		340	a 450	211	1,000	105	23	2.0	2.4	.7
8.....		253	360	208	960	82	30	1.7	2.2	.4
9.....		218	380	184	1,250	815	61	22	.7	1.1	.5
10.....		181	a 250	172	920	745	60	31	.7	.7	.7
11.....		185	255	103	960	620	58	42	.5	1.0	1.0
12.....		66	222	158	885	480	75	25	.5	2.2	.4
13.....		51	204	150	780	360	79	18	1.1	.7	.5
14.....	55	47	194	150	885	590	88	14	3.0	3.0	.4
15.....	51	39	228	197	745	455	140	14	8.8	6.4	.3
16.....	39	69	530	246	650	340	505	12	8.2	14	.4
17.....	58	95	a 1,200	239	590	300	280	10	7.0	10	.4
18.....	58	69	5,240	680	280	340	11	10	7.0	.3
19.....	43	57	1,890	1,610	300	405	10	3.9	5.5	.2
20.....	34	50	1,000	960	253	405	10	2.0	6.4	.1
21.....	21	54	a 750	680	222	320	9.6	1.5	5.5	.2
22.....	19	a 300	a 550	480	360	230	7.9	2.2	3.4	.2
23.....	18	a 500	a 400	300	280	106	6.4	1.2	3.0	.2
24.....	16	a 400	280	300	225	140	7.0	1.7	3.2	.2
25.....	16	a 300	256	280	200	123	4.1	.7	4.1	7.6
26.....	14	a 250	246	300	181	91	4.9	.8	3.0	4.9
27.....	12	a 250	239	236	150	88	6.1	2.2	2.8	2.2
28.....	12	253	208	214	138	75	8.2	2.6	2.0	2.0
29.....	12	239	184	250	131	64	7.6	2.8	1.4	2.2
30.....	20	218	175	300	125	58	5.8	1.0	1.1	.8
31.....		204	256	51	1.4	1.4

Month	Maximum	Minimum	Mean	Per square mile	Run-off in inches
1929					
September 14-30	58	12	29.3	0.195	0.12
1929-30.					
October	6,120	33	565	3.77	4.35
November	5,240	175	637	4.25	4.74
December 1-17	246	150	182	1.21	.76
March 9-31	1,610	214	634	4.23	3.62
April	1,340	125	471	3.14	3.50
May	505	37	150	1.00	1.15
June	42	4.1	17.3	.115	.13
July	10	.5	3.04	.020	.02
August	14	.5	3.30	.022	.03
September	7.6	.1	1.18	.0079	.009

Cherry River at Fenwick, W. Va.

Location.—Chain gage at highway bridge at Fenwick, Nicholas County, 1,000 feet below mouth of Laurel Creek.

Drainage area.—150 square miles.

Records available.—September, 1929, to September, 1931.

Extremes.—Maximum discharge during year, 4,900 second-feet Apr. 4 (gage height, 9.44 feet); minimum, 0.3 second-foot Oct. 10, 14 (gage height, 2.70 feet).

1920-1931: Maximum gage height, 12.04 feet Oct. 2, 1920 (discharge not determined); minimum discharge, 0.1 second-foot Sept. 22, 1930 (gage height, 2.02 feet).

Remarks.—Records good.

Daily and monthly discharge, in second-feet, 1930-31.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	0.8	1.4	123	90	300	710	080	340	300	34	77	194
2.....	.7	1.2	84	45	250	1,000	560	280	250	100	1,080	148
3.....	.5	1.4	40	77	211	530	1,160	246	228	155	2,400	228
4.....	.8	1.8	27	75	190	430	4,680	236	187	84	1,610	222
5.....	.8	2.0	20	405	117	320	2,200	184	152	75	745	125
6.....	.4	2.2	405	1,000	90	200	1,700	166	172	88	430	07
7.....	.8	2.0	455	780	138	340	000	228	320	115	320	03
8.....	1.2	2.8	54	480	300	430	020	1,160	505	86	280	74
9.....	.5	3.6	218	250	1,800	340	1,700	780	430	88	280	55
10.....	.3	3.2	84	211	1,340	280	2,300	710	340	131	250	45
11.....	.0	3.4	72	175	680	222	2,600	455	300	184	222	71
12.....	.0	3.2	100	100	530	160	2,290	430	107	125	242	81
13.....	.8	4.7	117	160	360	155	1,430	560	225	88	300	63
14.....	.5	6.7	101	84	480	250	1,000	1,520	620	74	107	30
15.....	.5	5.8	66	71	815	1,610	815	1,250	1,520	208	120	34
16.....	.5	0.4	32	23	405	1,000	710	780	1,000	09	05	64
17.....	1.1	7.6	40	125	530	050	680	710	560	72	74	100
18.....	1.7	5.5	47	00	1,080	560	300	1,120	300	77	00	52
19.....	1.1	6.7	41	138	1,120	780	300	1,160	250	40	84	60
20.....	.7	7.3	33	300	745	710	253	1,250	100	55	320	34
21.....	.7	5.8	22	155	530	560	232	1,900	230	50	1,340	36
22.....	.8	7.0	10	31	430	020	260	2,000	218	43	1,430	50
23.....	.7	7.0	23	01	340	530	430	2,030	178	42	2,190	39
24.....	1.2	5.5	20	125	320	650	430	1,700	160	320	1,120	06
25.....	1.3	5.8	17	181	300	850	530	1,120	105	150	680	101
26.....	1.4	7.0	155	320	240	710	780	680	88	88	505	480
27.....	1.4	7.0	360	745	223	080	745	530	09	75	360	505
28.....	1.8	7.9	211	900	320	1,520	710	480	58	60	300	430
29.....	1.8	7.0	150	080	2,000	560	380	50	54	480	340
30.....	2.0	11	99	505	1,160	280	246	46	42	380	250
31.....	1.7	03	405	780	280	26	260

Month	Maximum	Minimum	Mean	Per square mile	Run-off in inches
October	2.0	0.3	0.97	0.0065	0.007
November	11	1.2	5.08	.034	.64
December	455	17	109	.727	.84
January	1,000	23	323	2.15	2.48
February	1,890	99	613	3.42	3.56
March	2,090	155	674	4.49	5.18
April	4,680	232	1,080	7.20	8.03
May	2,030	100	838	5.59	6.44
June	1,520	40	311	2.07	2.31
July	320	26	90.7	.645	.74
August	2,490	60	593	3.95	4.55

ending Sept. 30, 1918-1922—Concluded.
[Drainage area 1,340 square miles.]

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1920-21					
October	620	128	215	.160	.18
November	3,690	158	945	.705	.79
December	10,600	1,200	2,780	2.07	2.39
January	9,760	980	2,550	1.90	2.19
February	9,180	1,130	2,150	1.60	1.67
March	9,180	905	2,640	1.90	2.19
April	1,430	592	889	.663	.74
May	3,160	499	1,260	.940	1.08
June	2,600	216	592	.442	.49
July	1,200	144	343	.256	.30
August	537	100	247	.184	.21
September	1,200	95	279	.208	.23
The year	10,600	95	1,230	.918	12.46
1921-22					
October	371	100	152	.113	.13
November	17,900	662	3,420	2.55	2.84
December	18,200	4,800	3.58	4.13
January	13,900	080	3,070	2.29	2.64
February	20,700	1,280	4,890	3.65	3.80
March	13,900	1,580	6,070	4.53	5.22
April	5,120	1,060	2,370	1.77	1.98
May	9,180	1,130	2,860	2.13	2.46
June	5,410	760	2,180	1.63	1.82
July	7,150	402	1,340	1.00	1.15
August	2,690	254	630	.470	.54
September	1,580	123	378	.282	.31
The year	20,700	100	2,670	1.99	27.02
1922-23					
October	314	111	156	0.116	0.13
November	176	111	131	.0078	.11
December	12,100	106	2,450	1.83	2.11
January	14,800	1,200	3,440	2.57	2.96
February	17,600	648	3,800	2.84	2.06
March	14,800	1,130	4,310	3.22	3.71
April	10,900	760	2,710	2.02	2.25
May	2,800	760	1,260	.940	1.08
June	4,250	307	958	.715	.80
July	802	111	245	.183	.21
August	11,500	190	1,890	1.41	1.63
September	1,280	167	451	.337	.38
The year	17,600	106	1,810	1.35	18.33

Monthly discharge of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1923-1934.

[Drainage area 1,340 square miles.]

1923-24					
October	266	117	169	0.126	0.15
November	1,280	144	491	.366	.41
December	9,470	905	2,590	1.93	2.22
January	21,300	1,360	4,670	3.49	4.02
February	6,860	830	2,170	1.62	1.75
March	10,800	1,430	4,620	3.37	3.88
April	6,860	1,200	2,970	2.22	2.48
May	25,000	1,130	4,710	3.52	4.06
June	7,150	537	2,420	1.81	2.02
July	4,830	227	1,060	.749	.86
August	7,440	195	1,310	.975	1.12
September	12,400	210	848	.633	.71
The year	25,000	117	2,330	1.74	23.68

Monthly discharge of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1923-1934—Continued.
[Drainage area 1,340 square miles.]

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1924-25					
October	13,000	170	1,250	0.933	1.08
November	4,540	188	911	.680	.76
December	13,300	366	2,130	1.59	1.83
January	9,470	1,280	3,260	2.43	2.86
February	0,760	1,280	3,950	2.95	3.07
March	13,900	006	2,070	1.54	1.78
April	5,700	537	1,550	1.16	1.29
May	6,570	401	2,250	1.68	1.94
June	1,360	271	684	.510	.57
July	648	128	325	.243	.28
August	194	84	137	.102	.12
September	106	71	85.9	.064	.07
The year	13,900	71	1,540	1.15	15.59
1925-26					
October	5,410	101	1,150	0.858	0.99
November	3,690	740	1,790	1.34	1.50
December	5,120	405	931	.695	.80
January	19,800	3,200	2.39	2.76
February	16,700	1,580	4,440	3.31	3.45
March	8,600	1,580	3,240	2.42	2.79
April	4,250	1,360	2,250	1.68	1.87
May	1,200	376	946	.482	.56
June	2,090	248	836	.624	.70
July	2,380	109	430	.321	.37
August	12,100	112	1,700	1.27	1.46
September	816	128	365	.272	.30
The year	19,800	101	1,730	1.29	17.55
1926-27					
October	2,000	295	1,240	0.925	1.07
November	15,100	816	3,030	2.26	2.52
December	30,400	1,580	6,260	4.67	5.38
January	8,020	2,500	1.91	2.20
February	18,800	2,480	6,940	5.18	5.39
March	7,440	080	2,240	1.67	1.02
April	15,100	2,090	5,550	4.14	4.62
May	10,000	1,060	2,370	1.77	2.04
June	2,920	356	1,340	1.00	1.12
July	934	188	344	.257	.30
August	2,090	212	906	.743	.86
September	1,280	118	275	.205	.23
The year	30,400	118	2,730	2.04	27.65
1927-28					
October	2,920	106	600	0.515	0.58
November	8,380	295	1,900	1.42	1.58
December	0,860	537	2,800	2.09	2.41
January	11,000	905	2,950	2.20	2.54
February	7,440	1,060	2,430	1.81	1.95
March	11,600	760	2,800	2.13	2.46
April	12,700	1,200	2,620	1.96	2.19
May	16,400	732	2,370	1.77	2.04
June	0,800	802	2,340	1.75	1.95
July	9,370	287	1,770	1.32	1.52
August	8,760	242	1,470	1.10	1.27
September	3,970	271	983	.734	.82
The year	16,400	106	2,100	1.57	21.31

Monthly discharge of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1928-1934—Continued.
[Drainage area 1,340 square miles.]

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1928-29					
October	1,660	200	510	0.387	0.45
November	4,830	425	1,150	.858	.96
December	21,400	810	2,780	2.07	2.39
January	9,500	905	2,890	2.16	2.49
February	29,100	905	3,320	2.48	2.58
March	20,700	2,000	5,800	4.37	5.04
April	6,280	1,060	2,750	2.05	2.29
May	17,300	1,360	4,310	3.22	3.71
June	5,410	500	1,280	.955	1.07
July	2,380	128	436	.325	.37
August	704	92	182	.136	.16
September	115	80	94.3	.0704	.08
The year	20,100	80	2,130	1.59	21.50
1929-30					
October	12,200	188	1,960	1.46	1.68
November	26,600	1,160	4,300	3.28	3.66
December	6,600	954	2,440	1.82	2.10
January	3,200	664	1,420	1.06	1.22
February	8,700	620	2,430	1.81	1.88
March	8,100	941	2,450	1.83	2.11
April	6,450	600	1,800	1.34	1.50
May	928	250	546	.407	.47
June	1,790	116	318	.237	.26
July	123	43	67.0	.050	.06
August	61	28	43.6	.033	.04
September	43	30	34.8	.026	.03
The year	26,600	28	1,480	1.10	15.01
1930-31					
October	58	28	36.0	0.027	0.03
November	100	40	67.0	.050	.06
December	493	53	171	.128	.15
January	2,870	202	687	.513	.59
February	3,990	310	1,340	1.00	1.04
March	9,300	610	2,630	1.96	2.20
April	13,800	928	4,020	3.00	3.35
May	12,900	838	3,500	2.61	3.01
June	4,140	413	1,120	.836	.93
July	928	269	481	.359	.41
August	4,650	173	1,150	.858	.90
September	1,300	147	408	.304	.34
The year	13,800	28	1,300	.970	13.16
1931-32					
October	580	116	191	0.143	0.16
November	103	110	127	.095	.11
December	4,020	202	1,540	1.15	1.33
January	13,600	754	3,310	2.47	2.85
February	37,100	1,230	4,910	3.66	3.95
March	10,300	980	5,330	3.98	4.59
April	12,300	790	2,780	2.07	2.31
May	21,800	484	3,670	2.74	3.16
June	13,400	244	1,330	.993	1.11
July	14,800	183	2,200	1.64	1.89
August	310	78	180	.134	.15
September	188	45	89.4	.067	.07
The year	37,100	45	2,130	1.59	21.68

Monthly discharge of Greenbrier River at Alderson, W. Va., for the years ending Sept. 30, 1923-1934—Concluded.
[Drainage area 1,340 square miles.]

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1932-33					
October	2,480	81	490	0.306	0.42
November	13,500	580	2,460	1.84	2.05
December	13,300	370	2,300	1.72	1.98
January	13,100	1,100	3,380	2.52	2.90
February	15,100	2,080	5,140	3.84	4.00
March	23,400	1,260	5,110	3.81	4.39
April	10,200	1,210	4,100	3.06	3.41
May	9,000	995	3,270	2.44	2.81
June	1,480	207	935	.474	.53
July	12,300	197	1,400	1.04	1.20
August	4,950	282	1,250	.933	1.08
September	778	123	292	.218	.24
The year	23,400	81	2,470	1.84	25.01
1933-34					
October	203	88	119	0.089	0.10
November	520	100	226	.169	.19
December	5,780	284	1,054	.787	.91
January	8,960	310	1,705	1.27	1.46
February	548	251	410	.306	.32
March	32,200	520	7,709	5.75	6.63
April	9,270	858	2,380	1.78	1.99
May	2,160	362	720	.542	.62
June	502	141	281	.210	.23
July	993	53	192	.143	.16
August	457	82	203	.151	.17
September	2,550	48	335	.250	.28
The year	32,200	48	1,201	.963	13.66

GREENBRIER RIVER AT ALDERSON, W. VA.

Location.—Water-stage recorder, lat. 37°43'50", long. 80°38'30", 400 feet above highway bridge at Alderson, Monroe County, and half a mile above the mouth of Muddy Creek. Zero of gage is 1,528.97 feet above mean sea level.

Drainage area.—1,357 square miles (revised).

Records available.—July 1895 to June 1906, May 1907 to September 1935.

Average discharge.—38 years (1895-1905, 1907-35), 2,080 second-feet.

Extremes.—Maximum discharge during year, about 48,500 second-feet Jan. 23 (gage height, 16.85 feet); minimum, 154 second-feet Oct. 30 (gage height, 2.04 feet). 1895-1935: Maximum observed discharge, about 70,000 second-feet (revised) Mar. 13, 14, 1918 (gage height, 22.0 feet); minimum, 20 second-feet part of Aug. 12, Oct. 1, 2, 1930 (gage height, 1.65 feet).

Remarks.—Records good except those for Dec. 11-17, Jan. 2-12, Mar. 19-27, and June 5 to July 1, which are fair and were estimated on basis of records for stations at Buckeye.

Rating table, water year 1934-35 (gage height, in feet, and discharge in second-feet)

2.0	136	3.4	1,680	8.0	14,300
2.1	182	3.6	1,990	9.0	17,900
2.2	234	3.8	2,380	10.0	21,600
2.3	290	4.0	2,790	11.0	25,300
2.4	355	4.5	3,910	12.0	29,100
2.6	520	5.0	5,100	13.0	33,000

	Date		Height (Feet)	(second- feet)
1806	Mar. 30		10.5	25,600
	Nov. 6		9.5	21,800
1897	Feb. 23		51,500
	23	17.5 P		54,000
	May 14		12.2	32,300
1808	Jan. 16		7.8	15,700
	Mar. 30		8.0	16,400
	May 7		8.0	16,400
	Aug. 11		14.8	42,900
	Oct. 22		9.8	23,000
1899	Jan. 7		8.6	18,600
	Feb. 27		8.0	18,600
	Mar. 5		15.4	45,300
1900	Feb. 14		7.9	16,000
	Mar. 21		8.2	17,100
	Nov. 26		18.2	56,800
1901	Jan. 12		9.3	21,100
	Apr. 21		9.1	20,400
	May 23		8.4	17,800
	28		8.8	19,300
	June 17		9.0	20,000
	Dec. 15		13.3	30,700
	30		11.1	28,000
1902	Feb. 26		17,800
	Mar. 1		11.8	30,700
	14		7.8	15,700
	17		15,000
	30		7.8	15,700
1903	Jan. 3		9.9	23,100
	Feb. 5		8.9	19,600
	17		10.3	24,900
	Mar. 1		10.0	23,800
	24		11.9	31,100
1904	Jan. 23		7.7	15,400
	May 19		7.8	15,700
1906	Mar. 10		10.5	25,600
	22		7.6	15,000
	May 12		11.2	28,400
1906	Jan. 23		9.3	21,100

1907	June 14		14.4	41,200
	Dec. 11		7.8	15,700
	24		8.9	19,600
1908	Jan. 12		9.5	21,800
	Feb. 16		14.0	39,600
	Mar. 7		10.4	25,300
	Apr. 1		10.6	26,000
	May 8		0.4	21,500
1900	Apr. 15		7.6	15,000
1910	June 17		12.8	34,500
1911	Jan. 4		9.9	23,100
	30		12.9	35,100
	Apr. 5		8.4	17,800
	Oct. 18		8.5	18,200
1912	Feb. 22		8.1	16,800
	27		8.4	17,800
	Mar. 16		11.9	31,100
	29		8.3	17,500
	May 13		8.3	17,500
	17		8.5	18,200
1013	Mar. 15		8.2	17,100
	Mar. 27		14.7	42,500
	Apr. 13		8.8	19,300
	15		17,100
	May 28		7.8	15,700
	Nov. 17		**	14,300
1914	Feb. 20		**	14,000
1915	Jan. 7		11.7	26,300
	Feb. 2		12.2	27,800
	2		14.5 P	34,000
	Oct. 2		9.11	18,500
	2		11.3 P	25,100

	Date		Height (Feet)	(second- feet)
	Dec. 30		9.09	18,500
1916	Dec. 29		8.0	15,200
1917	Mar. 4		27,200
	4	14.2 P		34,000
	13		0.8	20,600
1918	Feb. 14		8.00	15,200
	16		8.14	15,500
	21		8.30	16,100
-	Mar. 14		18.62	48,000
	14		22 P	58,900
	June 26		9.00	17,600
	26		10.8 P	23,600
	Oct. 31		11.15	24,100
	31		11.9 P	26,000
	Dec. 23		8.99	17,600
	23		11.0 P	24,200
1919	Jan. 2		15.90	39,400
	2		16.3 P	40,700
	Dec. 7		14.0 P	33,300
	8		10.52	22,200
1020	Jan. 25		9.49	19,200
	Mar. 20		10.41	21,900
	Dec. 15		6.68**	10,600
1921	Nov. 1		9.12	17,900
	29		8.68	16,700
	Dec. 25		9.25	18,200
1922	Feb. 21		10.00	20,700
	21		10.5 P	22,200
1923	Feb. 2		8.96	17,600
	2		9.58 P	19,500
1924	Jan. 4		8.18	15,100
	17		10.22	21,300
	Mar. 30		9.68	19,800
	May 12		11.40	25,000
	12		13.60 P	32,000
1925	Mar. 20		7.75**	13,900
1926	Jan. 20		0.72	19,800
	20		9.95 P	20,700
	22		8.48	10,000
	Feb. 15		8.08	16,700
	Nov. 17		8.15	15,100
	Dec. 22		10.68	22,900
	26		13.11	30,400
	26		14.50 P	34,900
1927	Feb. 6		9.38	18,800
	20		9.02	17,600
	23		8.28	15,400
	Apr. 10		8.25	15,100
1928	May 1		8.66	16,400
	1		9.0 P	18,000
	Dec. 1		9.95	21,400
1929	Feb. 28		12.22	29,100
	28		13.15 P	32,700
	Mar. 6		9.70	20,400
	May 21		8.70	17,300
	Nov. 18		26,600
	18		14.20 P	36,300
1930	Feb. 6		6.05**	8,700
1931	Apr. 5		7.74**	13,800
1932	Feb. 5		37,100
	5		16.96 P	46,400
	Mar. 18		16,000
	29		19,300
	May 2		21,800
1933	Feb. 21		15,400
	Mar. 20		10.64	23,400
	20		11.68 P	27,300
1934	Mar. 5		12.76	32,200
	5		13.21 P	33,800
	9		16,000
	28		10.85	24,000

Discharge, in second-feet, water year October, 1934 to September, 1935.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	3,800	256	17,500	2,120	1,430	4,020	23,800	872	1,300	1,300	845	256
2.....	1,810	378	12,000	2,300	1,340	3,230	16,800	838	1,080	1,380	725	245
3.....	1,040	430	0,360	1,900	1,440	2,600	8,780	912	800	773	596	245
4.....	681	630	4,260	1,600	1,480	2,140	7,380	1,260	020	737	3,350	528
5.....	520	577	3,450	1,500	1,360	1,020	0,600	1,610	2,000	773	3,120	10,000
6.....	2,050	692	2,680	1,400	1,210	2,120	5,100	1,740	1,000	1,230	1,920	18,300
7.....	4,140	953	2,200	1,300	1,120	2,480	5,850	10,200	1,300	1,560	1,000	7,120
8.....	2,500	1,210	1,830	1,300	1,060	2,440	8,480	13,900	1,100	14,700	12,000	4,140
9.....	1,460	1,040	1,510	1,600	066	2,100	8,190	6,600	1,000	19,400	7,120	2,790
10.....	900	797	1,260	5,500	2,270	1,850	6,350	4,380	1,200	8,190	4,020	2,010
11.....	725	048	1,100	3,800	5,600	3,580	4,980	3,340	1,100	4,020	3,450	1,460
12.....	568	548	850	3,200	4,740	16,800	4,740	2,640	000	2,790	4,140	1,120
13.....	466	466	750	2,680	3,560	20,500	5,600	2,200	800	2,070	2,640	872
14.....	386	415	850	2,280	3,120	10,400	5,850	2,120	800	1,530	1,770	703
15.....	320	385	000	1,900	6,220	6,220	4,740	2,120	700	1,830	1,580	596
16.....	200	336	800	1,700	8,480	4,620	3,800	2,300	1,200	1,400	1,120	530
17.....	262	322	600	4,500	7,120	3,450	3,120	3,010	1,000	1,100	858	466
18.....	240	200	000	9,410	5,350	2,900	2,520	3,460	800	905	681	415
19.....	224	208	602	6,220	4,020	2,800	2,260	2,900	800	785	586	385
20.....	213	262	1,320	4,860	3,120	2,800	2,010	2,320	700	615	530	355
21.....	198	262	1,970	7,380	2,500	2,700	1,900	2,120	600	520	475	330
22.....	102	250	1,880	20,100	2,050	2,700	1,810	2,620	500	053	475	329
23.....	187	426	1,560	41,900	1,950	3,000	1,580	2,900	460	714	568	355
24.....	177	4,500	1,310	18,600	2,420	13,000	1,380	2,000	600	681	586	355
25.....	164	5,100	1,180	7,910	2,640	13,500	1,210	3,460	450	726	466	303
26.....	164	2,900	1,490	5,480	3,500	17,000	1,970	3,450	375	1,940	370	279
27.....	108	1,900	7,380	4,140	7,120	9,300	966	2,700	300	6,480	322	251
28.....	108	1,480	5,850	2,700	5,600	6,100	912	2,220	275	4,140	200	234
29.....	159	11,400	3,910	2,240	4,860	899	1,790	240	2,460	284	218
30.....	164	10,100	2,900	1,090	3,910	912	1,540	300	1,580	268	203
31.....	159	2,340	1,610	9,130	1,490	1,080	273

Month	Second-foot-days	Maximum	Minimum	Mean	Per square mile	Run-off in inches
October	24,559	4,140	154	792	0.584	0.67
November	55,136	16,100	256	1,838	1.35	1.51
December	94,262	17,500	600	3,041	2.24	2.58
Calendar year 1934.....	601,896	32,200	48	1,649	1.22	16.62
January	175,110	41,900	1,300	5,649	4.16	4.80
February	92,840	8,480	966	3,316	2.44	2.54
March	184,170	20,500	1,850	5,041	4.38	5.05
April	149,589	23,800	899	4,086	3.67	4.10
May	96,037	13,900	833	3,098	2.28	2.03
June	25,235	2,000	240	841	.620	.00
July	87,541	19,400	520	2,824	2.08	2.40
August	50,928	12,000	268	1,836	1.35	1.56
September	55,999	18,300	203	1,867	1.38	1.54
Water year 1934-35.....	1,097,412	41,000	154	3,007	2.22	30.07

The following summary of flood stages and discharges for the Greenbrier River at Alderson is taken from the United States Geological Survey Water Supply Paper No. 771, pages 195-196. The base discharge assumed as flood level is 15,000

Meadow River.—Meadow River, which drains about one-sixth of Greenbrier County, has a meandering length of 52.58 miles, of which about 41 miles is within or along the border of the County. It has its source in eastern Summers County at an elevation of approximately 2800 feet and empties into Gauley River at Carnifex Ferry, Nicholas County, at an elevation of about 1180 feet. The rate of fall is not uniform from the source to the mouth as the following table shows:

Gradient of Meadow River.

	Miles	Elevation	Fall, Feet	Fall per Mile, Feet
Source		2800		
Distance	0.3		100	333.0
Summers-Greenbrier line		2700		
Distance	2.7		265	9.8
Grassy Meadows		2435		
Distance	13.3		40	3.0
Rupert		2395		
Distance	6.7		20	3.0
East Rainelle		2375		
Distance	18.3		500	27.3
Corner of Fayette-Greenbrier- Nicholas		1875		
Distance	11.3		695	61.5
Mouth		1180		

The above table emphasizes the local base-leveling along Meadow River.*

* See page 35.

A gaging station was established near Russellville, July 17, 1908, for which the following records are available, being taken from the various Water Supply Papers of the United States Geological Survey previously quoted under the designation of Gauley River. The station was discontinued

Meadow River near Russellville, W. Va.

Location.—At Bays Ferry, one-fourth mile below mouth of Youngs Creek and 3 miles below Russellville, Fayette County.

Drainage Area.—297 square miles.

Records available.—July 17, 1908 to September 30, 1916, when station was discontinued.

Gage.—Chain gage attached to trees on left bank 25 feet above bridge, near former ferry crossing, read by J. R. Bays.

Discharge measurements.—Made from bridge or by wading. Prior to completion of concrete bridge in 1913 high-water measurements were made from boat.

Channel and control.—Channel straight above and slightly curved for 200 feet below gage. Left bank subject to overflow at extremely high stages. Bed rocky and clean. Control permanent.

Extremes of discharge.—1908-1916: Maximum stage recorded, 13.25 feet morning reading February 3, 1915 (discharge about 7,300 second-feet); minimum stage recorded, 2.57 feet August 7, 8, 1914 (discharge, 6.7 second feet).

Ice.—Stage-discharge relation affected by ice for short periods in severe winters.

Accuracy.—Stage-discharge relation practically permanent, occasionally affected by ice. Rating curve well defined between 12 and 4,800 second-feet; beyond these limits the curve is an extension. Gage read to hundredths twice daily. Daily discharge ascertained by applying mean daily gage height to rating table. See foot-note to table of daily discharge for special estimates. Records good.

Discharge measurements of Meadow River near Russellville, W. Va., during the years 1908-1916

Date.	Made by—	Gage height.	Discharge.	Date.	Made by—	Gage height.	Discharge.
1908.		Feet.	Sec.-ft.	1912.		Feet.	Sec.-ft.
July 18	Wm. M. O'Neill.....	4.07	154	Mar. 27	C. T. Balley.....	7.02	1,370
Aug. 7	W. G. Hoyt.....	3.75	68	30do	9.44	3,230
1909.				1913.			
Apr. 6	H. J. Jackson.....	5.82	686	Nov. 18	Peterson and Walters	9.04	2,830
13do	4.55	260	20	M. I. Walters.....	6.50	1,050
1910.				1914.			
Mar. 24	C. T. Balley.....	4.46	233	Oct. 30	Mathers and Morgan.	3.39	40.0
26do	4.30	210	30do	3.36	44.7
Oct. 14do	3.54	62.9				
15do	3.39	47.1	1916.			
1911.				Aug. 21	B. E. Jones.....	5.01	370
July 29do	2.03	15.2	24do	5.70	632

Daily discharge, in second-feet, of Meadow River near Russellville, W. Va., for the years ending Sept. 30, 1908-1916.

Day.	July.	Aug.	Sept.	Day.	July.	Aug.	Sept.	Day.	July.	Aug.	Sept.
1908.				1908.				1908.			
1.....		202	47	11.....		324	22	21.....	138	48	13
2.....		148	41	12.....		226	20	22.....	134	46	13
3.....		118	34	13.....		148	20	23.....	191	46	12
4.....		07	30	14.....		116	19	24.....	148	44	12
5.....		83	28	15.....		96	19	25.....	133	45	12
6.....		92	28	10.....		86	17	26.....	138	00	12
7.....		86	28	17.....		82	16	27.....	406	238	11
8.....		78	27	18.....		75	15	28.....	765	158	11
9.....		100	25	19.....		158	02	20.....	580	100	11
10.....		170	24	20.....		118	50	21.....	804	75	11

Daily discharge, in second-feet, of Meadow River near Russellville, W. Va., for the
years ending Sept. 30, 1908-1916—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1908-9.												
1.....	10	127	47	1,200	020	670	765	2,070	311	202	54	14
2.....	11	91	58	975	715	540	625	2,070	250	605	64	18
3.....	11	62	48	625	580	505	625	1,040	220	878	115	12
4.....	12	47	47	470	505	020	580	1,140	226	208	08	14
5.....	12	89	45	765	505	920	715	865	286	202	58	26
6.....	12	34	44	2,150	580	865	715	625	311	337	44	25
7.....	12	30	70	1,710	580	1,260	625	505	311	975	36	37
8.....	12	28	180	1,200	540	2,150	505	470	250	765	31	33
9.....	12	28	250	815	505	2,390	406	470	220	640	29	33
10.....	13	26	238	540	1,140	3,000	378	670	274	400	33	68
11.....	15	22	202	438	1,570	2,470	324	1,570	364	250	32	250
12.....	16	26	715	406	1,200	1,570	286	1,260	438	191	28	238
13.....	17	28	920	364	1,030	1,140	262	075	378	100	28	158
14.....	20	32	580	324	920	1,200	2,550	070	505	286	22	88
15.....	34	32	438	1,200	865	1,200	2,550	470	580	238	20	50
16.....	37	30	837	1,990	1,640	920	1,640	378	505	169	214	40
17.....	28	30	250	1,780	1,850	715	1,080	324	378	125	505	42
18.....	28	35	208	1,440	1,320	540	715	274	406	97	337	46
19.....	26	34	580	1,080	1,030	438	505	226	350	115	202	71
20.....	24	44	580	815	1,080	400	438	202	202	100	120	54
21.....	23	120	438	705	1,080	378	406	202	214	88	80	38
22.....	22	138	304	815	1,440	505	865	274	101	73	56	32
23.....	20	111	337	815	1,380	540	1,570	324	158	73	40	29
24.....	20	91	286	865	1,260	580	1,710	286	180	07	35	83
25.....	19	73	274	715	1,380	1,140	1,320	238	202	127	28	148
26.....	19	62	438	025	1,200	1,990	975	304	180	44	23	86
27.....	19	57	438	580	1,030	1,920	715	765	130	73	22	62
28.....	19	53	406	505	865	2,550	625	815	118	70	21	48
29.....	37	51	406	470	2,150	580	765	102	59	19	43
30.....	47	47	470	438	1,500	705	540	131	61	17	23
31.....	158	1,260	920	1,030	400	54	10
1900-10.												
1.....	25	89	85	106	600	2,150	133	580	202	214	67	29
2.....	24	83	82	148	600	2,150	125	438	262	169	52	73
3.....	22	75	80	2,200	865	1,710	133	364	238	158	50	180
4.....	19	72	78	2,150	865	1,200	148	337	226	202	48	138
5.....	17	62	75	1,500	625	020	337	298	286	180	46	350
6.....	16	56	67	1,080	700	670	324	274	2,630	250	42	337
7.....	10	50	67	2,470	700	580	286	238	2,710	350	38	220
8.....	16	47	86	1,920	765	470	250	274	1,570	438	37	148
9.....	16	58	109	1,320	670	406	214	470	1,500	262	36	116
10.....	18	78	04	1,080	625	378	180	670	715	286	33	106
11.....	27	364	96	1,250	580	350	169	625	1,080	226	27	89
12.....	88	337	124	1,250	505	324	214	765	1,380	180	28	72
13.....	214	250	202	1,440	470	324	070	1,140	1,020	226	41	109
14.....	158	214	815	1,200	438	540	865	805	1,710	286	45	180
15.....	122	169	815	1,030	406	625	670	625	1,640	274	37	202
16.....	89	148	715	765	505	505	540	470	2,790	226	79	169
17.....	75	438	765	2,470	438	470	406	3,510	214	64	106
18.....	72	470	765	2,630	378	1,030	378	2,390	262	37	65
19.....	76	505	1,990	2,150	311	1,200	406	1,640	505	35	53
20.....	58	378	1,570	1,570	262	1,140	378	020	540	40	51
21.....	52	274	1,780	1,080	274	1,570	438	580	406	31	47
22.....	50	214	2,470	1,140	262	1,920	505	470	274	42	45
23.....	57	180	1,640	1,440	250	1,900	580	406	202	57	41
24.....	99	85	148	1,260	1,200	226	1,990	505	378	180	45	30
25.....	220	138	765	920	214	1,570	505	350	120	33	40
26.....	226	129	580	670	191	1,140	505	274	109	30	51
27.....	214	113	580	540	180	920	438	262	88	28	79
28.....	191	111	765	715	169	920	378	311	180	23	148
29.....	148	106	715	158	865	311	337	180	20	138
30.....	133	100	580	148	715	274	274	127	20	202
31.....	104	97	505	148	286	88	22

Daily discharge, in second-feet, of Meadow River near Russellville, W. Va., for the
years ending Sept. 30, 1908-1916—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1910-11.												
1.....	202	66	470	2,310	2,880	350	670	378	202	33	10	80
2.....	158	73	540	2,690	3,390	350	625	406	238	25	9.4	127
3.....	106	78	920	4,700	1,640	311	865	470	138	20	8.8	75
4.....	61	86	540	3,420	1,200	298	2,150	470	94	25	8.0	46
5.....	48	94	406	1,920	975	274	3,600	406	82	58	7.5	28
6.....	38	96	337	1,200	715	2,070	3,150	337	76	41	7.2	26
7.....	37	91	286	815	625	2,970	2,230	298	102	116	12	24
8.....	51	85	286	625	540	2,390	1,710	274	68	75	43	22
9.....	158	76	286	438	975	1,440	1,920	250	52	58	41	16
10.....	202	75	262	400	1,200	1,440	1,850	238	40	37	23	13
11.....	131	73	238	364	1,080	1,710	1,440	214	37	37	28	15
12.....	92	71	191	337	865	1,440	1,080	169	37	138	28	30
13.....	71	68	191	670	670	1,140	975	109	60	110	20	36
14.....	54	66	169	1,380	505	920	920	148	120	71	14	34
15.....	47	66	158	1,380	438	715	1,570	131	92	44	12	33
16.....	41	66	148	1,570	337	625	1,710	125	70	28	12	40
17.....	36	68	131	1,260	298	505	1,380	111	57	24	10	111
18.....	33	60	129	815	274	470	975	107	57	23	10	83
19.....	30	53	158	625	311	470	715	100	59	18	14	56
20.....	28	44	202	505	470	1,380	1,200	80	76	16	14	40
21.....	26	44	101	540	540	1,320	1,570	80	68	15	13	33
22.....	33	49	191	1,440	540	975	1,500	70	59	19	10	92
23.....	48	51	226	1,380	438	815	1,640	64	44	17	8.8	09
24.....	71	59	364	1,140	410	580	1,260	58	35	19	8.4	89
25.....	54	88	670	920	378	470	920	59	30	20	7.8	04
26.....	46	214	580	815	364	406	670	71	28	18	7.6	47
27.....	40	214	505	1,320	350	470	505	64	54	16	7.4	32
28.....	57	470	438	1,380	350	670	438	57	56	15	7.2	30
29.....	67	920	715	1,990	625	378	238	44	15	10	27
30.....	97	670	3,890	6,880	670	364	180	40	14	14	23
31.....	70	3,060	4,700	670	100	12	19
1911-12.												
1.....	22	115	438	920	1,200	975	1,640	1,260	104	262	100	100
2.....	33	99	378	920	975	715	1,850	920	91	148	85	158
3.....	80	86	337	765	670	505	2,550	765	79	111	68	118
4.....	180	76	274	625	580	378	2,070	580	68	148	48	91
5.....	158	70	262	000	540	364	1,320	438	60	250	44	88
6.....	158	197	250	580	438	324	920	438	51	324	38	82
7.....	109	1,440	202	540	406	298	670	815	40	350	33	71
8.....	191	1,260	180	470	378	298	505	020	38	298	30	59
9.....	214	865	191	406	311	715	670	815	36	169	28	44
10.....	180	670	180	337	238	920	540	625	32	226	27	37
11.....	250	470	180	215	226	865	470	505	23	202	26	33
12.....	378	337	101		214	975	378	3,060	20	119	25	30
13.....	298	406	202	214	214	2,230	350	3,150	18	158	23	29
14.....	238	438	202		202	2,150	324	2,390	10	109	23	21
15.....	378	438	191	214	180	4,590	298	1,440	14	169	22	16
16.....	470	378	226	225	180	6,040	350	2,880	14	158	21	16
17.....	378	324	262		180	3,330	406	4,810	13	158	19	10
18.....	2,710	540	364	1,420	191	1,850	540	2,880	21	364	17	17
19.....	1,710	805	350		238	1,380	625	1,440	47	262	23	18
20.....	1,080	670	337	870	337	1,200	540	920	89	118	50	17
21.....	670	505	286	650	1,380	1,500	470	580	102	80	148	16
22.....	406	406	286	505	1,809	1,500	406	406	62	68	324	16
23.....	378	350	406	360	2,230	1,320	540	337	43	52	262	27
24.....	406	350	865		1,380	1,920	580	286	35	44	191	158
25.....	337	479	020	1,030	975	2,630	505	238	28	148	109	202
26.....	274	470	1,030		1,590	2,230	438	214	59	438	71	238
27.....	226	438	1,440	215	3,690	1,380	540	180	107	298	53	180
28.....	191	470	1,440	214	2,310	1,030	1,200	148	865	202	46	158
29.....	169	580	1,080		1,440	2,550	1,030	148	625	133	104	138
30.....	148	505	815	1,640	3,240	1,320	138	438	118	214	118
31.....	124	765	1,500	2,390	118	197	180

Daily discharge, in second-feet, of Meadow River near Russellville, W. Va., for the
years ending Sept. 30, 1908-1916—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1912-13												
1.....	107	113	67	1,320	976	1,030	670	470	815	51	30	20
2.....	88	106	92	975	366	866	505	378	580	41	26	10
3.....	72	00	134	020	075	505	406	337	438	37	25	16
4.....	61	83	286	1,200	2,390	680	360	298	530	337	20	14
5.....	61	72	765	1,200	1,990	470	324	274	1,080	274	18	13
6.....	42	67	975	1,200	1,320	438	208	238	815	406	19	12
7.....	37	226	1,080	2,630	866	878	250	337	920	378	16	14
8.....	34	1,920	865	6,900	865	311	226	311	1,440	202	15	17
9.....	30	1,260	580	2,710	1,080	286	202	274	1,440	122	14	22
10.....	28	715	438	1,640	1,080	350	191	238	1,030	106	12	24
11.....	25	606	860	1,440	1,030	1,080	180	214	626	106	38	28
12.....	24	378	208	1,200	1,200	1,780	274	191	438	124	62	24
13.....	24	311	262	1,080	1,040	1,380	680	180	850	102	88	21
14.....	25	280	214	075	1,640	2,230	580	168	274	76	40	17
15.....	28	262	214	765	1,600	3,700	3,510	148	214	91	33	16
16.....	31	226	202	580	1,140	2,710	2,640	148	180	124	33	14
17.....	37	202	101	506	470	1,040	1,570	878	160	168	30	14
18.....	34	180	169	438	324	1,080	1,030	580	191	131	28	16
19.....	53	160	169	406	298	766	715	438	104	118	33	33
20.....	238	148	148	378	286	540	470	364	82	765	30	46
21.....	226	138	133	438	311	470	406	1,200	71	630	25	88
22.....	180	131	107	606	406	406	350	1,080	71	406	65	640
23.....	220	126	102	605	470	337	311	1,200	76	202	202	298
24.....	406	120	06	765	505	311	274	3,400	75	143	238	180
25.....	438	118	08	1,200	438	298	250	1,710	71	134	124	88
26.....	337	116	100	1,200	378	438	238	1,140	120	102	72	73
27.....	298	100	122	1,080	378	2,300	298	1,990	115	83	49	52
28.....	191	86	134	1,030	705	4,000	488	8,900	115	64	39	44
29.....	180	56	131	020	2,470	505	2,230	78	51	30	35
30.....	148	47	378	815	1,570	505	1,600	64	44	24	40
31.....	127	1,090	765	1,030	1,080	30	20
1913-14												
1.....	104	226	1,200	438	1,780	766	1,380	470	33	7.1	9.0	67
2.....	202	214	2,470	438	1,440	765	1,710	378	30	0.0	9.0	01
3.....	280	180	1,860	438	1,140	815	1,710	324	28	7.1	8.6	54
4.....	070	153	1,320	1,080	865	816	1,320	298	25	7.1	3.0	45
5.....	438	136	076	920	670	815	1,030	324	23	12	7.6	35
6.....	236	122	716	765	640	765	816	540	37	12	7.2	30
7.....	202	113	580	625	075	025	025	1,080	35	10	6.8	26
8.....	148	100	1,080	580	1,260	505	1,030	1,030	33	9.6	6.3	21
9.....	125	238	020	070	1,030	438	2,310	976	30	9.0	17	18
10.....	94	400	766	920	805	337	1,710	365	27	12	27	15
11.....	80	470	670	1,080	670	715	1,140	715	23	9.8	22	14
12.....	71	438	505	920	506	1,440	805	580	10	8.8	28	17
13.....	65	705	438	505	1,080	715	505	16	8.0	34	19
14.....	76	1,140	364	470	1,030	680	438	14	11	31	18
15.....	61	2,150	337	650	500	1,260	505	337	12	29	26	18
16.....	54	2,070	324	500	1,920	2,150	274	10	88	23	19
17.....	62	3,000	208	505	3,330	2,550	250	9.0	82	17	18
18.....	50	2,310	274	625	670	3,600	1,850	220	8.6	02	14	16
19.....	51	1,500	262	625	1,140	1,850	1,570	214	9.0	50	12	14
20.....	113	975	250	670	1,990	1,260	1,710	191	7.9	45	9.0	12
21.....	286	670	250	3,790	2,070	920	1,570	180	8.2	40	58	10
22.....	350	505	238	2,710	1,380	765	1,380	148	7.0	33	32	9.6
23.....	311	304	238	1,710	1,320	975	1,140	129	9.8	27	14	10
24.....	378	311	226	1,260	1,140	670	865	116	9.8	20	12	14
25.....	1,200	262	274	3,240	1,030	1,030	625	102	8.6	15	39	12
26.....	2,390	238	670	2,150	865	2,150	540	89	7.9	12	61	9.0
27.....	1,710	214	670	1,570	815	3,240	920	30	7.6	12	138	8.0
28.....	1,080	298	670	1,500	765	2,790	975	70	7.3	9.8	148	7.7
29.....	625	1,320	540	1,380	2,470	765	57	7.3	9.2	111	7.4
30.....	337	1,030	505	1,320	1,850	580	46	7.1	8.8	86	7.2
31.....	262	470	1,780	1,440	38	9.0	75

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1014-15												
1.....	7.0	37	57	1,080	1,640	580	214	1,570	274	21	19	25
2.....	6.8	37	180	865	6,040	505	214	1,140	324	37	22	22
3.....	6.6	34	406	670	6,880	406	202	920	337	56	27	20
4.....	6.5	33	625	505	3,790	350	202	815	324	73		25
5.....	0.8	31	1,850	378	2,150	337	202	975	274	66		337
6.....	7.5	30	2,310	364	1,710	324	202	865	226	59		337
7.....	7.4	28	1,780	6,700	1,500	364	202	765	191	52		298
8.....	7.2	27	1,030	5,580	1,320	378	202	540	238	56		214
9.....	7.1	25	765	2,470	1,080	364	226	438	202	202	18	134
10.....	7.4	24	765	1,440	865	350	220	406	158	180		97
11.....	8.4	23	765	1,140	765	470	220	364	120	148		51
12.....	8.4	22	715	1,030	580	505	286	837	113	118		40
13.....	8.8	20	650	920	470	470	311	311	116	83		34
14.....	11	20	580	670	406	438	311	208	109	64		30
15.....	20	23	540	865	1,320	438	286	262	540	40	42	24
16.....	43	25	350	865	1,990	470	274	238	470	64	35	23
17.....	138	61	238	2,150	1,570	505	262	220	406	57	25	22
18.....	160	102	169	3,890	1,320	505	250	202	324	61	48	21
19.....	148	92	214	5,580	1,030	470	250	180	250	56	46	20
20.....	102	75	505	4,490	765	470	238	158	191	66	42	23
21.....	72	66	975	2,070	540	505	214	148	148	125	37	40
22.....	60	58	1,380	1,260	378	406	202	122	100	109	31	80
23.....	51	52	1,140	1,030	337	350	191	109	80	68	20	102
24.....	45	49	865	765	350	311	324	106	70	54	22	70
25.....	41	48	715	765	670	274	438	99	58	46	22	53
26.....	41	47	505	670	1,080	262	438	99	49	43	20	49
27.....	58	45	505	580	865	274	406	106	41	39	18	72
28.....	71	44	438	540	625	274	815	99	35	35	17	133
29.....	54	41	406	580		250	2,390	92	29	30	17	120
30.....	46	41	1,320	670		226	1,990	180	24	20	18	99
31.....	39		1,500	1,160		214		214		22	28	
1915-16												
1.....	1,780	50	311	1,500	1,780	865	1,380	438	580		238	78
2.....	3,150	48	298	1,440	1,990	670	1,180	406	540		670	70
3.....	1,730	46	274	1,080	1,710	580	975	378	540		1,320	67
4.....	1,260	44	262	765	1,380	505	670	406	580	125	1,200	60
5.....	625	42	250	670	1,140	505	505	438	540		1,920	56
6.....	470	40	226	580	975	1,140	438	406	540		1,640	40
7.....	365	39	202	670	815	3,090	378	378	540	60	1,260	45
8.....	214	38	180	715	670	3,330	378	324	540	57	1,140	45
9.....	148	37	158	670	865	2,550	378	274	540	54	2,470	62
10.....	120	35	148	925	975	1,640	438	226	580	79	1,990	70
11.....	102	35	136	1,500	920	1,200	1,380	202	580	75	1,640	61
12.....	92	33	134	2,970	920	975	2,880	169	540	274	1,440	58
13.....	85	30	125	2,880	865	715	3,060	158	470	262	1,200	56
14.....	72	42	122	2,390	865	580	2,470	148	438	238	975	50
15.....	61	286	118	1,380	815	815	1,320	148	580	226	865	298
16.....	56	540	120	1,030	715	975	920	138	1,140	250	625	286
17.....	54	540	505	815	625	815	765	148	1,260	765	2,150	262
18.....	51	540	3,330	1,200	580	715	670	148	1,140	715	1,380	238
19.....	58	815	2,710	920	540	625	470	138	1,030	580	1,080	169
20.....	125	975	1,440	840	540	540	378	136	865	470	920	109
21.....	113	865	1,200	705	540	505	324	124	715	438	580	78
22.....	94	670	975	952	540	975	311	113	580	865	438	61
23.....	80	540	715	1,140	625	1,320	298	214	409	1,380	815	59
24.....	75	470	540	975	1,030	1,200	262	670	324	1,200	580	56
25.....	71	406	505	765	3,330	975	350	670	298	580	378	53
26.....	67	378	865	625	3,150	815	580	580	438	378	202	47
27.....	64	350	765	505	2,230	865	625	540	324	324	214	39
28.....	58	337	765	438	1,570	2,630	625	505	262	378	158	39
29.....	54	324	2,550	406	1,030	2,710	580	470	214	350	131	54
30.....	53	311	4,920	1,030		2,310	505	540	148	311	100	49
31.....	51		2,880	1,030		1,780		670		286		

NOTE.—Daily discharge estimated, because of ice or missing gage readings, from observer's notes, climatic data, or by comparison with flow at other stations as follows: Nov. 17-30, Dec. 26-31, 1909; Jan. 3, 11-12, Feb. 1-2, 6-7, Dec. 8, 1010; Jan. 10, Feb. 24, Dec. 5, 1911; Jan. 5, 11-14, 16-21, 23-28, Feb. 22, Dec. 25, 1912; Jan. 8, Feb. 2, 1913; Jan. 10, 1914; Jan. 11, 1915; Jan. 12, 1916; Jan. 13, 1917; Jan. 14, 1918; Jan. 15, 1919; Jan. 16, 1920; Jan. 17, 1921; Jan. 18, 1922; Jan. 19, 1923; Jan. 20, 1924; Jan. 21, 1925; Jan. 22, 1926; Jan. 23, 1927; Jan. 24, 1928; Jan. 25, 1929; Jan. 26, 1930; Jan. 27, 1931; Jan. 28, 1932; Jan. 29, 1933; Jan. 30, 1934; Jan. 31, 1935; Jan. 1, 1936; Jan. 2, 1937; Jan. 3, 1938; Jan. 4, 1939; Jan. 5, 1940; Jan. 6, 1941; Jan. 7, 1942; Jan. 8, 1943; Jan. 9, 1944; Jan. 10, 1945; Jan. 11, 1946; Jan. 12, 1947; Jan. 13, 1948; Jan. 14, 1949; Jan. 15, 1950; Jan. 16, 1951; Jan. 17, 1952; Jan. 18, 1953; Jan. 19, 1954; Jan. 20, 1955; Jan. 21, 1956; Jan. 22, 1957; Jan. 23, 1958; Jan. 24, 1959; Jan. 25, 1960; Jan. 26, 1961; Jan. 27, 1962; Jan. 28, 1963; Jan. 29, 1964; Jan. 30, 1965; Jan. 31, 1966; Jan. 1, 1967; Jan. 2, 1968; Jan. 3, 1969; Jan. 4, 1970; Jan. 5, 1971; Jan. 6, 1972; Jan. 7, 1973; Jan. 8, 1974; Jan. 9, 1975; Jan. 10, 1976; Jan. 11, 1977; Jan. 12, 1978; Jan. 13, 1979; Jan. 14, 1980; Jan. 15, 1981; Jan. 16, 1982; Jan. 17, 1983; Jan. 18, 1984; Jan. 19, 1985; Jan. 20, 1986; Jan. 21, 1987; Jan. 22, 1988; Jan. 23, 1989; Jan. 24, 1990; Jan. 25, 1991; Jan. 26, 1992; Jan. 27, 1993; Jan. 28, 1994; Jan. 29, 1995; Jan. 30, 1996; Jan. 31, 1997; Jan. 1, 1998; Jan. 2, 1999; Jan. 3, 2000; Jan. 4, 2001; Jan. 5, 2002; Jan. 6, 2003; Jan. 7, 2004; Jan. 8, 2005; Jan. 9, 2006; Jan. 10, 2007; Jan. 11, 2008; Jan. 12, 2009; Jan. 13, 2010; Jan. 14, 2011; Jan. 15, 2012; Jan. 16, 2013; Jan. 17, 2014; Jan. 18, 2015; Jan. 19, 2016; Jan. 20, 2017; Jan. 21, 2018; Jan. 22, 2019; Jan. 23, 2020; Jan. 24, 2021; Jan. 25, 2022; Jan. 26, 2023; Jan. 27, 2024; Jan. 28, 2025; Jan. 29, 2026; Jan. 30, 2027; Jan. 31, 2028; Jan. 1, 2029; Jan. 2, 2030; Jan. 3, 2031; Jan. 4, 2032; Jan. 5, 2033; Jan. 6, 2034; Jan. 7, 2035; Jan. 8, 2036; Jan. 9, 2037; Jan. 10, 2038; Jan. 11, 2039; Jan. 12, 2040; Jan. 13, 2041; Jan. 14, 2042; Jan. 15, 2043; Jan. 16, 2044; Jan. 17, 2045; Jan. 18, 2046; Jan. 19, 2047; Jan. 20, 2048; Jan. 21, 2049; Jan. 22, 2050; Jan. 23, 2051; Jan. 24, 2052; Jan. 25, 2053; Jan. 26, 2054; Jan. 27, 2055; Jan. 28, 2056; Jan. 29, 2057; Jan. 30, 2058; Jan. 31, 2059; Jan. 1, 2060; Jan. 2, 2061; Jan. 3, 2062; Jan. 4, 2063; Jan. 5, 2064; Jan. 6, 2065; Jan. 7, 2066; Jan. 8, 2067; Jan. 9, 2068; Jan. 10, 2069; Jan. 11, 2070; Jan. 12, 2071; Jan. 13, 2072; Jan. 14, 2073; Jan. 15, 2074; Jan. 16, 2075; Jan. 17, 2076; Jan. 18, 2077; Jan. 19, 2078; Jan. 20, 2079; Jan. 21, 2080; Jan. 22, 2081; Jan. 23, 2082; Jan. 24, 2083; Jan. 25, 2084; Jan. 26, 2085; Jan. 27, 2086; Jan. 28, 2087; Jan. 29, 2088; Jan. 30, 2089; Jan. 31, 2090; Jan. 1, 2091; Jan. 2, 2092; Jan. 3, 2093; Jan. 4, 2094; Jan. 5, 2095; Jan. 6, 2096; Jan. 7, 2097; Jan. 8, 2098; Jan. 9, 2099; Jan. 10, 2100; Jan. 11, 2101; Jan. 12, 2102; Jan. 13, 2103; Jan. 14, 2104; Jan. 15, 2105; Jan. 16, 2106; Jan. 17, 2107; Jan. 18, 2108; Jan. 19, 2109; Jan. 20, 2110; Jan. 21, 2111; Jan. 22, 2112; Jan. 23, 2113; Jan. 24, 2114; Jan. 25, 2115; Jan. 26, 2116; Jan. 27, 2117; Jan. 28, 2118; Jan. 29, 2119; Jan. 30, 2120; Jan. 31, 2121; Jan. 1, 2122; Jan. 2, 2123; Jan. 3, 2124; Jan. 4, 2125; Jan. 5, 2126; Jan. 6, 2127; Jan. 7, 2128; Jan. 8, 2129; Jan. 9, 2130; Jan. 10, 2131; Jan. 11, 2132; Jan. 12, 2133; Jan. 13, 2134; Jan. 14, 2135; Jan. 15, 2136; Jan. 16, 2137; Jan. 17, 2138; Jan. 18, 2139; Jan. 19, 2140; Jan. 20, 2141; Jan. 21, 2142; Jan. 22, 2143; Jan. 23, 2144; Jan. 24, 2145; Jan. 25, 2146; Jan. 26, 2147; Jan. 27, 2148; Jan. 28, 2149; Jan. 29, 2150; Jan. 30, 2151; Jan. 31, 2152; Jan. 1, 2153; Jan. 2, 2154; Jan. 3, 2155; Jan. 4, 2156; Jan. 5, 2157; Jan. 6, 2158; Jan. 7, 2159; Jan. 8, 2160; Jan. 9, 2161; Jan. 10, 2162; Jan. 11, 2163; Jan. 12, 2164; Jan. 13, 2165; Jan. 14, 2166; Jan. 15, 2167; Jan. 16, 2168; Jan. 17, 2169; Jan. 18, 2170; Jan. 19, 2171; Jan. 20, 2172; Jan. 21, 2173; Jan. 22, 2174; Jan. 23, 2175; Jan. 24, 2176; Jan. 25, 2177; Jan. 26, 2178; Jan. 27, 2179; Jan. 28, 2180; Jan. 29, 2181; Jan. 30, 2182; Jan. 31, 2183; Jan. 1, 2184; Jan. 2, 2185; Jan. 3, 2186; Jan. 4, 2187; Jan. 5, 2188; Jan. 6, 2189; Jan. 7, 2190; Jan. 8, 2191; Jan. 9, 2192; Jan. 10, 2193; Jan. 11, 2194; Jan. 12, 2195; Jan. 13, 2196; Jan. 14, 2197; Jan. 15, 2198; Jan. 16, 2199; Jan. 17, 2200; Jan. 18, 2201; Jan. 19, 2202; Jan. 20, 2203; Jan. 21, 2204; Jan. 22, 2205; Jan. 23, 2206; Jan. 24, 2207; Jan. 25, 2208; Jan. 26, 2209; Jan. 27, 2210; Jan. 28, 2211; Jan. 29, 2212; Jan. 30, 2213; Jan. 31, 2214; Jan. 1, 2215; Jan. 2, 2216; Jan. 3, 2217; Jan. 4, 2218; Jan. 5, 2219; Jan. 6, 2220; Jan. 7, 2221; Jan. 8, 2222; Jan. 9, 2223; Jan. 10, 2224; Jan. 11, 2225; Jan. 12, 2226; Jan. 13, 2227; Jan. 14, 2228; Jan. 15, 2229; Jan. 16, 2230; Jan. 17, 2231; Jan. 18, 2232; Jan. 19, 2233; Jan. 20, 2234; Jan. 21, 2235; Jan. 22, 2236; Jan. 23, 2237; Jan. 24, 2238; Jan. 25, 2239; Jan. 26, 2240; Jan. 27, 2241; Jan. 28, 2242; Jan. 29, 2243; Jan. 30, 2244; Jan. 31, 2245; Jan. 1, 2246; Jan. 2, 2247; Jan. 3, 2248; Jan. 4, 2249; Jan. 5, 2250; Jan. 6, 2251; Jan. 7, 2252; Jan. 8, 2253; Jan. 9, 2254; Jan. 10, 2255; Jan. 11, 2256; Jan. 12, 2257; Jan. 13, 2258; Jan. 14, 2259; Jan. 15, 2260; Jan. 16, 2261; Jan. 17, 2262; Jan. 18, 2263; Jan. 19, 2264; Jan. 20, 2265; Jan. 21, 2266; Jan. 22, 2267; Jan. 23, 2268; Jan. 24, 2269; Jan. 25, 2270; Jan. 26, 2271; Jan. 27, 2272; Jan. 28, 2273; Jan. 29, 2274; Jan. 30, 2275; Jan. 31, 2276; Jan. 1, 2277; Jan. 2, 2278; Jan. 3, 2279; Jan. 4, 2280; Jan. 5, 2281; Jan. 6, 2282; Jan. 7, 2283; Jan. 8, 2284; Jan. 9, 2285; Jan. 10, 2286; Jan. 11, 2287; Jan. 12, 2288; Jan. 13, 2289; Jan. 14, 2290; Jan. 15, 2291; Jan. 16, 2292; Jan. 17, 2293; Jan. 18, 2294; Jan. 19, 2295; Jan. 20, 2296; Jan. 21, 2297; Jan. 22, 2298; Jan. 23, 2299; Jan. 24, 2300; Jan. 25, 2301; Jan. 26, 2302; Jan. 27, 2303; Jan. 28, 2304; Jan. 29, 2305; Jan. 30, 2306; Jan. 31, 2307; Jan. 1, 2308; Jan. 2, 2309; Jan. 3, 2310; Jan. 4, 2311; Jan. 5, 2312; Jan. 6, 2313; Jan. 7, 2314; Jan. 8, 2315; Jan. 9, 2316; Jan. 10, 2317; Jan. 11, 2318; Jan. 12, 2319; Jan. 13, 2320; Jan. 14, 2321; Jan. 15, 2322; Jan. 16, 2323; Jan. 17, 2324; Jan. 18, 2325; Jan. 19, 2326; Jan. 20, 2327; Jan. 21, 2328; Jan. 22, 2329; Jan. 23, 2330; Jan. 24, 2331; Jan. 25, 2332; Jan. 26, 2333; Jan. 27, 2334; Jan. 28, 2335; Jan. 29, 2336; Jan. 30, 2337; Jan. 31, 2338; Jan. 1, 2339; Jan. 2, 2340; Jan. 3, 2341; Jan. 4, 2342; Jan. 5, 2343; Jan. 6, 2344; Jan. 7, 2345; Jan. 8, 2346; Jan. 9, 2347; Jan. 10, 2348; Jan. 11, 2349; Jan. 12, 2350; Jan. 13, 2351; Jan. 14,

Monthly discharge of Meadow River near Russellville, W. Va., for the
years ending Sept. 30, 1908-1916.
[Drainage area, 207 square miles.]

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1908.					
July 10-31	765	133	275	0.926	0.45
August	470	44	121	.407	.47
September	47	11	20.2	.068	.08
1908-9.					
October	158	10	24.7	.083	.10
November	138	25	54.7	.184	.21
December	1,260	44	356	1.20	1.38
January	2,150	324	897	3.02	3.48
February	1,850	505	1,030	3.47	3.61
March	3,060	378	1,230	4.14	4.77
April	2,550	262	861	2.90	3.24
May	2,070	202	705	2.37	2.73
June	580	102	282	.949	1.06
July	975	44	231	.778	.90
August	505	16	77.7	.262	.30
September	250	12	62.3	.210	.23
The year	3,060	10	482	1.62	22.01
1909-10.					
October	226	16	80.1	.290	.33
November	364	47	111	.374	.42
December	815	67	220	.761	.88
January	2,470	106	1,210	4.07	4.69
February	2,630	406	944	3.18	3.31
March	2,150	148	547	1.84	2.12
April	1,990	125	757	2.55	2.84
May	1,140	238	475	1.00	1.84
June	3,510	226	1,100	3.70	4.13
July	540	88	239	.805	.93
August	70	20	39.8	.134	.15
September	350	20	123	.414	.46
The year	3,510	16	484	1.63	22.10
1910-11.					
October	202	26	72.3	.243	.28
November	920	44	141	.475	.53
December	3,890	129	544	1.83	2.11
January	6,880	337	1,640	5.52	6.36
February	2,880	274	777	2.62	2.73
March	2,970	274	934	3.14	3.62
April	3,600	364	1,330	4.48	6.00
May	470	57	191	.643	.74
June	238	28	74.0	.249	.28
July	138	12	38.2	.129	.16
August	43	7.2	14.4	.048	.06
September	127	13	48.9	.165	.18
The year	6,880	7.2	483	1.63	22.04
1911-12.					
October	2,710	22	403	1.36	1.57
November	1,440	70	473	1.59	1.77
December	1,440	180	469	1.58	1.82
January	1,640	530	1.78	2.05
February	3,690	180	848	2.86	3.08
March	6,040	298	1,670	5.62	6.48
April	2,550	298	802	2.70	3.01
May	4,810	118	1,090	3.67	4.23
June	865	13	108	.364	.41
July	438	44	189	.036	.73
August	324	17	79.1	.266	.31
September	262	16	81.4	.274	.31
The year	6,040	13	563	1.90	25.77

Monthly discharge of Meadow River near Russellville, W. Va., for the
years ending Sept. 30, 1908-1916—Continued.

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mle	
1912-13					
October	438	24	123	0.414	0.48
November	1,920	47	270	.930	1.05
December	1,990	67	351	1.18	1.36
January	5,900	378	1,190	4.01	4.62
February	2,390	286	914	3.08	3.21
March	4,090	286	1,160	3.91	4.51
April	3,510	180	615	2.07	2.31
May	3,900	148	851	2.87	3.31
June	1,440	64	421	1.42	1.58
July	765	36	181	.609	.70
August	238	12	48.5	.163	.19
September	540	12	61.5	.207	.23
The year	5,900	12	515	1.73	23.55
1913-14					
October	2,390	50	392	1.32	1.52
November	3,900	106	794	2.67	2.98
December	2,470	226	656	2.21	2.55
January	3,790	438	1,180	3.97	4.58
February	2,070	470	979	3.30	3.44
March	3,000	337	1,370	4.61	5.32
April	2,550	505	1,220	4.11	4.59
May	1,080	38	357	1.20	1.38
June	37	7.1	17.0	.057	.06
July	88	6.9	22.0	.074	.09
August	148	6.8	35.4	.119	.14
September	67	7.2	21.1	.071	.08
The year	3,900	6.8	581	1.97	26.73
1914-15.					
October	169	6.5	42.1	.142	.16
November	102	20	42.0	.141	.16
December	2,310	57	782	2.63	3.03
January	6,760	364	1,670	5.62	6.48
February	6,880	337	1,500	5.05	5.26
March	580	214	389	1.31	1.51
April	2,390	191	406	1.37	1.53
May	1,570	92	399	1.34	1.54
June	540	24	194	.653	.73
July	202	21	69.8	.235	.27
August	48	17	24.5	.082	.09
September	337	20	87.2	.294	.33
The year	6,880	6.5	462	1.56	21.09
1915-16.					
October	3,150	51	369	1.24	1.43
November	975	33	297	1.00	1.12
December	4,920	118	894	3.01	3.47
January	2,970	406	1,070	3.60	4.15
February	3,330	540	1,160	3.91	4.22
March	3,690	505	1,270	4.28	4.93
April	3,060	262	850	2.86	3.19
May	670	113	332	1.12	1.29
June	1,260	148	576	1.04	2.16
July	1,380	54	366	1.23	1.42
August	2,470	89	964	3.25	3.75
September	298	39	91.1	.307	.34
The year	4,920	33	688	2.32	31.47

Location.—Chain gage on highway bridge at Nallen, Fayette County.

Drainage area.—207 square miles.

Records available.—July, 1908, to September, 1916; November, 1928, to September, 1929.

Extremes.—Maximum discharge during year, 5,140 second-feet Feb. 28 (gage height, 12.03 feet); minimum, 8 second-feet Sept. 7 (gage height, 2.05 feet).

1908-1916, 1928-29: Maximum discharge, about 7,300 second-feet Feb. 3, 1915 (gage height, 13.25 feet); minimum, 6.7 second-feet Aug. 7 and 8, 1914 (gage height, 2.57 feet).

Remarks.—Records good. Discharge estimated because of ice Jan. 30 to Feb. 2 and because of missing record Sept. 8.

Daily and monthly discharge, in second-feet, 1928-29.

Day.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....		3,880	264	650	4,690	400	1,060	2,620	250	43	18
2.....		2,780	340	600	3,180	430	1,010	1,300	785	36	14
3.....		1,520	830	570	1,770	392	1,580	875	875	32	12
4.....		1,060	1,150	570	1,520	340	1,840	695	570	124	12
5.....		785	875	510	3,340	830	1,520	610	302	170	11
6.....		570	1,010	570	4,600	875	1,200	430	264	104	9.6
7.....		450	1,200	740	3,520	740	1,010	308	200	73	8.4
8.....		340	1,580	695	2,220	650	920	237	159	54	8.9
9.....		250	1,770	650	1,520	490	785	237	105	39	430
10.....		224	1,350	650	920	410	050	278	100	29	340
11.....		200	785	570	695	340	510	224	00	27	324
12.....		190	650	450	570	308	410	169	124	31	169
13.....		179	530	375	510	250	358	150	212	31	108
14.....		190	740	293	065	224	1,250	169	179	29	75
15.....		740	830	212	1,580	200	1,400	237	150	20	02
16.....		1,100	610	170	1,350	250	1,200	490	132	20	45
17.....		1,150	392	170	1,010	610	1,150	392	108	17	37
18.....		1,150	324	179	875	875	065	278	87	15	85
19.....		1,200	830	224	740	920	1,520	212	72	14	33
20.....		965	470	340	650	920	2,220	159	57	12	32
21.....		830	875	430	570	875	3,020	132	46	10	29
22.....		695	1,010	490	490	1,060	2,140	102	36	9.2	24
23.....		610	1,200	430	1,100	1,100	1,350	83	31	41	21
24.....	570	470	1,300	430	2,540	920	065	124	26	410	18
25.....	490	375	1,300	392	2,060	785	875	450	24	237	16
26.....	410	293	1,640	1,250	1,400	785	740	830	29	116	14
27.....	490	264	1,520	3,180	1,100	740	650	430	60	70	12
28.....	358	224	1,150	4,780	830	785	450	278	70	47	12
29.....	324	212	875		650	1,300	875	324	87	37	11
30.....	1,640	237	800		530	1,250	1,250	308	69	28	11
31.....		190	700		510		2,540		48	24	

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
November 24-30	1,640	324	612	2.06	0.54
December	3,880	179	752	2.53	2.92
January	1,770	264	932	3.14	3.62
February	4,780	179	785	2.47	2.57
March	4,690	490	1,550	5.22	0.02
April	1,300	200	671	2.26	2.52
May	3,020	358	1,210	4.07	4.69
June	2,620	83	438	1.47	1.64
July	875	24	176	.593	.68
August	410	9.2	63.4	.213	.25
		8.4	65.0	.219	.24

Meadow River at Nallen, W. Va.

Location.—Chain gage on highway bridge at Nallen, Fayette County.

Drainage area.—297 square miles.

Records available.—July, 1908, to September, 1916; November, 1928, to September, 1930.

Extremes.—Maximum discharge during year, 6,140 second-feet Oct. 2 (gage height, 13.05 feet); practically no flow Sept. 23-24, 28-30.

1908-1910, 1928-1930: Maximum discharge, about 7,300 second-feet Feb. 3, 1915 (gage height, 13.25 feet); practically no flow Sept. 23-24, 28-30, 1930.

Remarks.—Records good except those above 4,000 second-feet and those estimated because of ice, Nov. 30, Dec. 1-5, 24-27, Jan. 18 to Feb. 2, which are fair.

Daily and monthly discharge, in second-feet, 1929-30.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	11	1,460	310	965	130	670	308	108	69	0.7	0.6	0.4
2.....	2,300	1,400	280	875	130	470	203	103	58	1.4	.6	.4
3.....	4,690	3,610	250	1,010	109	375	392	116	49	1.5	.5	.3
4.....	2,780	3,700	200	1,010	264	324	1,200	104	41	1.4	.4	.3
5.....	1,770	2,460	170	875	2,380	293	1,000	95	34	1.1	.4	.3
6.....	965	1,460	250	695	1,980	410	875	108	30	.9	.4	.2
7.....	650	965	308	010	1,400	1,250	920	159	35	.7	.4	.2
8.....	392	695	430	530	1,060	4,000	1,000	132	33	.5	.4	.2
9.....	264	510	830	490	830	3,180	1,010	124	37	.4	.4	.2
10.....	100	375	740	430	650	2,060	920	102	35	.4	.3	.1
11.....	150	324	650	392	530	1,400	740	91	20	.4	.3	.1
12.....	124	308	570	340	450	1,250	570	86	25	.4	.3	.1
13.....	100	278	400	203	450	1,100	470	73	21	.4	.3	.1
14.....	04	278	430	293	530	1,010	375	100	19	3.4	.3	.1
15.....	38	358	392	293	510	965	340	132	16	10	.5	.1
16.....	83	570	375	278	570	785	308	150	13	9.2	.5	.1
17.....	78	1,000	340	250	510	650	264	150	12	8.4	.5	.1
18.....	73	5,540	340	210	410	530	250	170	15	7.6	.5	.1
19.....	64	4,420	570	180	450	095	237	570	14	6.0	.4	.1
20.....	61	2,780	1,350	140	430	785	224	830	13	4.0	.4	.1
21.....	57	1,400	1,150	160	400	740	224	605	11	2.3	.4	.1
22.....	1,770	920	920	180	490	610	250	490	8.8	.7	1.0	.1
23.....	3,430	095	740	180	530	530	200	375	7.6	6.0	2.0	0
24.....	2,380	510	000	100	695	430	179	308	6.8	4.3	2.5	0
25.....	1,840	410	500	140	1,060	375	159	212	6.0	2.8	1.5	.1
26.....	830	340	450	130	1,010	392	141	159	5.3	1.5	1.2	.1
27.....	610	324	400	125	830	358	132	141	4.0	3.4	1.7	.1
28.....	410	358	650	100	570	358	124	124	3.4	4.6	1.0	0
29.....	340	430	1,400	160	358	116	104	.7	3.7	.7	0
30.....	650	360	1,250	150	368	108	91	.7	1.7	.5	0
31.....	1,460	1,100	150	340	80	1.0	.4

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
October	4,690	11	926	3.12	3.60
November	5,540	278	1,280	4.31	4.81
December	1,400	170	695	2.00	2.31
January	1,010	125	382	1.29	1.40
February	2,380	130	697	2.35	2.45
March	4,600	293	889	2.99	3.45
April	1,200	108	448	1.51	1.68
May	830	73	203	.684	.79
June	69	.7	21.7	.073	.08
July	10	.4	3.11	.010	.01
August	2.5	.3	.69	.0023	.003
September00017	.0005

**Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1895-1917—Continued.**

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1915-16.												
1.....	9,800	253	049	5,030	5,350	2,190	3,030	1,800	1,250	516	700	294
2.....	18,500	253	581	3,710	8,050	2,090	2,290	1,600	1,000	581	7,220	268
3.....	0,400	230	570	4,490	6,400	2,810	2,090	1,340	1,000	777	3,480	253
4.....	3,250	211	526	3,710	4,220	2,810	1,390	1,250	1,000	690	1,890	232
5.....	2,000	107	474	2,000	3,250	2,500	1,700	1,160	1,160	1,000	1,510	204
6.....	1,600	185	403	2,290	3,030	3,250	1,700	1,080	030	818	1,700	190
7.....	1,160	190	423	2,000	2,810	5,300	2,090	930	1,000	452	1,700	190
8.....	1,000	204	328	1,900	3,710	2,390	2,100	860	520	452	2,290	190
10.....	078	100	442	1,340	3,960	4,760	2,100	840	2,810	468	1,800	204
10.....	673	100	442	1,340	3,060	4,760	2,190	846	2,810	463	1,800	204
11.....	020	190	334	4,220	3,480	3,250	5,300	764	2,500	673	1,510	232
12.....	547	100	328	13,400	3,030	2,600	6,680	712	2,000	700	1,420	186
13.....	494	179	311	11,000	3,250	2,100	5,850	660	1,000	615	1,340	160
14.....	442	185	336	8,600	4,220	2,190	4,220	592	1,250	484	1,250	164
15.....	423	218	408	5,300	3,710	3,710	3,250	581	3,710	452	902	232
16.....	364	364	328	3,710	2,810	3,030	2,500	547	6,680	626	1,600	1,250
17.....	364	738	403	2,810	2,500	3,430	2,000	536	12,200	2,300	2,390	860
18.....	330	010	002	2,100	2,090	2,600	1,700	558	8,050	3,480	1,900	660
19.....	364	874	12,500	1,600	1,790	2,090	3,710	558	4,220	2,290	1,340	463
20.....	403	1,800	5,850	1,100	1,420	1,890	1,340	520	2,810	1,420	1,340	355
21.....	413	2,090	3,480	1,340	1,340	1,510	1,160	484	1,990	1,080	1,000	277
22.....	432	1,700	2,290	1,510	1,340	1,600	1,250	452	1,510	1,160	700	225
23.....	474	1,340	1,790	1,700	1,250	3,710	1,160	686	3,250	3,030	2,810	232
24.....	432	1,160	1,510	1,800	1,340	5,030	1,100	2,600	1,080	1,800	1,890	232
26.....	384	1,000	1,100	1,510	6,400	3,480	1,250	2,500	1,080	1,420	1,080	107
26.....	345	840	1,420	1,000	7,500	2,600	2,300	1,700	1,340	1,080	804	226
27.....	336	790	1,700	1,990	4,490	2,100	3,250	5,580	930	840	502	232
28.....	302	764	1,800	2,810	3,250	2,290	3,250	2,500	860	1,100	505	190
29.....	302	764	7,500	3,030	2,500	6,120	2,810	1,600	686	1,600	452	860
30.....	294	712	18,500	6,120	5,580	2,290	1,340	502	1,100	304	2,810
31.....	253	8,320	6,950	3,060	1,420	874	328
1010-17.												
1.....	2,190	328	553	2,500	0,120	12,800	1,700	3,710	2,600	179	345	82
2.....	1,250	311	615	1,990	6,400	11,300	1,510	4,490	2,000	218	302	87
3.....	860	286	649	1,700	4,220	12,500	1,420	3,250	2,090	211	846	82
4.....	687	277	592	2,390	2,390	27,200	1,340	2,290	2,300	239	374	70
5.....	586	260	547	6,580	1,790	24,800	1,340	1,990	2,390	225	277	84
6.....	463	260	547	10,400	1,510	0,800	6,580	1,790	2,000	218	302	00
7.....	304	246	536	8,600	1,420	0,120	6,950	1,420	1,700	197	277	92
8.....	345	232	494	5,030	1,420	5,580	5,030	1,420	1,420	174	240	100
9.....	320	225	526	3,480	1,420	11,600	5,030	1,090	1,160	158	286	218
10.....	294	218	404	2,600	1,030	7,500	3,480	3,030	030	137	311	336
11.....	277	232	494	2,090	910	4,760	2,810	2,810	930	137	336	649
12.....	260	225	470	1,600	888	9,800	2,390	2,390	002	133	345	423
13.....	200	239	403	1,080	002	20,600	2,390	2,090	764	137	208	311
14.....	239	253	448	1,800	738	17,600	2,300	1,700	649	122	218	225
15.....	260	260	432	3,030	777	14,300	2,000	1,510	581	118	185	185
16.....	225	239	526	2,810	846	8,900	1,700	1,250	526	153	174	158
17.....	260	204	277	2,190	700	10,100	1,510	1,080	474	246	148	183
18.....	345	225	442	1,790	030	11,000	1,340	1,000	452	260	133	122
19.....	505	218	442	1,600	1,420	7,500	1,160	874	413	302	133	118
20.....	1,420	185	442	1,420	3,710	5,030	1,080	790	384	616	122	104
21.....	1,420	179	442	1,340	7,500	3,960	1,000	725	442	432	115	100
22.....	1,080	190	725	6,400	5,300	6,120	030	673	505	374	115	05
23.....	860	204	2,190	13,700	4,220	6,400	002	680	463	463	107	07
24.....	712	225	2,100	6,680	12,200	14,600	804	073	364	604	100	02
25.....	604	328	1,890	4,220	12,800	13,700	777	649	311	874	97	90
26.....	558	902	1,420	3,030	6,050	7,220	790	615	277	3,030	97	90
27.....	463	673	1,340	2,390	4,490	4,760	874	1,250	253	1,700	100	95
28.....	432	526	2,810	1,800	4,220	3,480	818	12,500	232	1,080	97	130
29.....	394	494	15,200	1,790	2,810	002	8,050	218	738	00	164
30.....	355	620	8,050	2,290	2,500	2,600	5,580	204	526	84	226
31.....	355	3,480	3,250	1,990	3,480	423	74

NOTE.—Daily discharge interpolated or estimated because of ice from observer's notes.

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1918-1922.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1917-18.												
1.....	232	2,600	403	1,100	3,250	5,580	1,660	3,290	1,430	6,280	2,060	3,970
2.....	190	1,516	394	1,000	2,390	6,466	1,500	2,690	1,200	4,830	1,660	1,910
3.....	164	1,080	423	832	2,090	5,030	1,580	2,280	1,060	3,290	1,200	1,430
4.....	158	860	394		1,990	3,480	1,820	2,090	905	2,280	605	1,000
5.....	148	086	384		1,700	5,856	1,740	1,616	766	1,740	676	816
6.....	126	592	394		1,660	7,780	1,580	1,740	774	1,430	511	980
7.....	118	516	374		1,600	10,400	1,500	1,580	905	1,200	449	1,300
8.....	107	463	336	500	3,480	6,260	1,660	1,580	1,130	080	353	1,430
9.....	107	423	311		3,060	5,580	12,700	1,660	980	830	367	1,280
10.....	104	384	204		10,700	4,490	10,900	2,580	816	746	279	1,000
11.....	100	346	174		12,500	3,960	8,020	2,180	662	662	273	605
12.....	111	311	153	832	11,300	3,250	7,730	1,610	474	606	307	718
13.....	114	264		1,680	12,200	15,200	6,286	1,746	371	090	344	600
14.....	114	277		1,080	15,200	48,000	6,700	1,910	362	718	600	524
15.....	114	253		1,420	12,500	16,000	9,180	2,480	330	634	1,300	490
16.....	114	246		1,256	15,500	10,306	9,470	2,696	314	524	980	626
17.....	114	232		1,250	8,600	6,860	7,440	2,230	573	592	020	524
18.....	114	218	150	1,100	5,300	4,540	5,700	1,910	005	676	606	764
19.....	114	204		1,080	3,710	3,420	4,546	2,180	2,926	1,660	905	7,730
20.....	204	197		930	6,500	2,660	3,970	1,916	2,926	2,660	1,360	3,420
21.....	1,000	190			10,100	2,580	9,760	2,090	1,820	1,740	1,366	2,380
22.....	874	100			7,500	8,310	11,500	2,180	1,500	1,200	080	2,380
23.....	592	179	403		4,700	6,570	7,440	2,920	2,000	880	732	2,000
24.....	404	174	404	690	3,710	4,540	5,120	2,860	1,660	062	606	1,600
25.....	403	158	600		3,480	4,250	4,540	2,380	1,360	537	980	1,360
26.....	403	158	874		13,100	3,970	4,540	6,280	17,606	449	1,200	1,130
27.....	423	158	1,340	030	14,000	3,290	6,860	0,280	10,600	1,130	830	980
28.....	516	153	1,600	7,500	7,220	2,690	7,730	3,090	4,836	774	662	816
29.....	600	148	1,510	11,000		2,280	5,410	2,600	5,120	564	606	704
30.....	3,030	225	1,340	0,950		2,000	3,970	2,280	0,470	524	578	600
31.....	4,760		1,250	4,490		1,740		1,820		1,280	537	
1918-19.												
1.....	537	10,900	3,290	16,000	1,740	4,250	2,800	1,000	1,360	2,000	732	401
2.....	474	5,990	2,480	39,400	1,430	0,280	2,380	7,730	1,740	1,580	1,740	788
3.....	402	3,100	2,180	21,600	1,366	4,540	2,000	5,990	1,300	1,280	1,500	606
4.....	371	2,180	1,740	10,000	1,360	3,420	1,820	3,970	1,130	1,280	1,130	402
5.....	336	1,320	1,580	5,690	1,436	2,920	1,740	2,920	980	980	830	321
6.....	314	1,660	1,430	4,540	1,300	3,160	1,660	2,380	905	830	605	200
7.....	307	1,500	1,280	3,690	1,130	3,690	1,500	2,090	788	2,620	1,130	227
8.....	286	1,300	1,200	3,040	1,660	3,290	1,430	2,060	746	3,160	905	216
9.....	279	1,200	1,130	2,480	1,060	3,420	1,360	6,280	660	2,000	732	190
10.....	270	1,000	1,280	2,180	980	6,280	1,280	14,800	006	1,280	606	210
11.....	280	080	2,180	1,740	830	5,120	1,280	11,500	620	980	474	221
12.....	279	905	4,540	1,820	760	3,690	7,780	6,570	788	830	461	210
13.....	260	830	4,540	1,740	905	2,920	5,900	4,250	830	1,500	401	232
14.....	238	760	3,420	1,660	1,500	2,480	3,970	3,690	905	3,040	1,000	190
15.....	243	676	11,300	1,746	2,186	2,180	3,046	4,830	905	4,540	980	260
16.....	273	718	9,700	1,740	2,180	1,820	2,280	4,540	1,200	8,020	704	105
17.....	286	732	5,410	1,740	1,910	1,660	4,540	3,420	1,500	11,200	578	176
18.....	280	1,360	4,250	4,540	1,600	1,820	5,120	3,970	2,480	6,280	564	167
19.....	254	2,280	2,926	9,760	1,506	2,180	3,690	3,420	2,380	4,540	524	151
20.....	286	2,286	2,386	0,570	1,436	2,280	2,806	2,800	1,660	13,606	437	167
21.....	300	2,000	2,090	4,540	1,360	2,690	2,380	4,250	2,000	8,310	437	171
22.....	381	1,740	8,310	3,420	1,580	1,820	2,000	5,120	2,380	5,900	391	185
23.....	425	1,506	17,600	3,290	2,020	1,666	1,820	3,670	1,580	5,126	336	205
24.....	592	1,360	8,000	12,700	3,070	1,660	1,820	3,160	1,860	5,120	307	216
25.....	816	1,130	6,286	9,470	3,420	1,500	1,740	8,020	4,540	2,020	293	221
26.....	0,570	1,060	5,410	5,760	8,310	1,360	1,660	7,440	8,020	2,180	300	210
27.....	5,690	980	4,250	3,070	7,440	1,745	1,500	4,250	13,900	1,000	371	300
28.....	3,420	1,130	3,100	3,040	4,540	13,000	1,430	3,160	8,890	1,360	273	232
29.....	2,480	6,280	2,580	2,580		7,440	1,360	2,480	5,120	1,200	232	210
30.....	3,296	5,126	2,696	2,180		4,830	1,360	2,000	2,020	980	227	176

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1918-1922—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1910-20.												
1.....	158	906	2,280		1,820	1,740	1,580	1,740	980	425	221	486
2.....	140	9,470	1,910		1,580	1,660	6,280	1,740	830	409	232	413
3.....	144	7,160	1,060		1,500	1,580	8,600	1,820	802	704	221	344
4.....	171	3,000	1,600		2,000	1,600	5,120	1,740	000	2,580	200	314
5.....	140	2,480	1,200	620	3,160	1,910	7,780	1,580	3,970	2,920	196	270
6.....	151	1,010	1,500		2,920	6,570	8,800	1,360	10,300	1,740	185	243
7.....	168	1,600	21,600		2,380	3,070	7,150	1,280	4,830	1,280	232	238
8.....	190	1,280	22,200		2,000	2,800	6,280	1,360	3,040	1,180	210	227
9.....	161	1,060	0,180	1,910	1,820	2,280	6,090	1,430	2,000	1,000	180	227
10.....	151	006	7,440	6,700	1,740	2,090	5,700	1,430	1,740	830	243	210
11.....	151	830	6,800	6,120	2,280	1,820	4,830	1,360	1,430	732	314	210
12.....	273	788	4,330	3,100	2,680	1,910	3,690	1,280	1,060	1,360	353	210
13.....	788	700	3,970	2,380	2,380	0,180	3,420	1,280	330	1,910	362	216
14.....	006	760	8,020	2,180	2,380	11,800	4,830	6,700	774	1,500	437	210
16.....	774	732	5,990	1,500	2,380	0,280	3,690	3,690	718	1,130	1,360	200
10.....	830	690	4,640	1,280	2,280	4,540	2,800	2,600	802	005	1,060	185
17.....	802	676	3,420	1,500	2,000	6,280	2,380	2,000	090	774	090	180
18.....	906	604	2,680	1,130	2,000	8,000	2,090	1,740	550	676	048	107
19.....	732	400	2,090	1,430	1,910	15,100	1,820	1,680	662	830	1,600	164
20.....	602	486	1,820	005	1,680	21,900	2,600	1,600	1,740	620	1,010	144
21.....	490	449	1,430	1,360	1,430	10,300	5,700	1,740	4,540	499	6,990	140
22.....	425	413	1,360	12,700	1,740	6,280	0,760	2,000	3,090	449	3,040	133
23.....	537	437	1,130	15,400	3,690	4,540	6,280	1,740	2,480	301	2,090	130
24.....	1,010	611		16,000	6,670	3,420	4,250	1,680	1,660	307	1,580	130
26.....	3,970	678		10,200	6,280	2,800	3,040	1,660	1,360	321	1,360	171
20.....	2,680	634		9,470	4,250	2,480	2,480	1,740	080	070	1,060	180
27.....	1,820	3,600	850	5,410	2,020	2,480	2,480	1,910	774	611	830	105
28.....	1,430	4,830		3,970	2,380	2,280	2,280	1,060	704	461	600	200
29.....	1,200	3,040		3,040	2,090	2,090	2,180	1,430	578	301	550	227
30.....	905	2,480		2,380		1,820	2,000	1,280	499	203	524	266
31.....	760			2,180		1,580		1,060		243	634	
1020-21.												
1.....	620	158	4,640	1,430	1,820	1,430	905	1,280	2,690	238	260	05
2.....	537	162	6,280	1,580	1,580	4,540	905	1,430	2,090	240	216	05
3.....	461	171	4,830	1,430	1,430	5,900	830	1,280	1,740	232	227	07
4.....	402	185	3,420	1,430	1,430	9,180	980	1,280	1,430	106	511	117
6.....	381	216	2,090	1,430	1,280	6,570	906	1,280	1,180	171	449	144
6.....	321	221	2,090	1,280	1,280	4,830	905	1,280	905	154	300	140
7.....	273	221	2,280	1,130	1,130	3,100	830	1,280	732	147	321	133
8.....	254	227	1,740	1,130	1,280	2,690	816	1,200	678	144	381	147
9.....	227	195	1,430	1,130	2,280	2,280	802	1,200	474	280	314	205
10.....	205	180	1,580	1,200	4,640	2,180	788	1,200	474	307	264	606
11.....	185	180	1,430	1,130	0,180	2,090	700	1,130	425	286	200	1,200
12.....	171	185	1,280	1,130	6,280	1,910	732	1,300	371	286	185	550
13.....	167	200	1,280	1,000	3,690	1,740	676	2,180	353	391	170	321
14.....	154	210	3,970	980	2,690	1,660	606	2,800	321	020	167	203
15.....	151	227	10,600	3,970	1,910	2,000	592	1,910	270	1,200	158	266
16.....	140	353	7,730	4,540	1,740	3,420	648	1,130	232	980	162	227
17.....	140	1,740	2,690	3,290	1,580	3,160	774	1,200	216	905	227	205
18.....	133	2,090	2,380	2,380	1,430	2,480	980	1,060	221	718	440	195
19.....	133	1,280	1,820	1,820	1,280	2,090	1,280	830	260	550	537	185
20.....	120	1,060	1,580	1,500	1,430	1,910	1,200	788	314	353	301	100
21.....	126	980	1,200	1,580	1,580	1,060	1,060	718	314	293	314	362
22.....	126	1,200	1,280	3,970	1,430	1,430	980	606	314	270	243	788
23.....	123	1,500	1,430	9,760	1,280	1,300	830	611	293	216	105	402
24.....	120	1,500	3,420	0,180	1,280	1,280	774	499	266	190	171	227
25.....	126	1,280	2,380	5,120	1,280	1,280	704	578	232	171	151	200
26.....	126	1,130	1,740	3,090	1,430	1,280	662	774	232	158	140	185
27.....	130	1,280	2,180	2,580	1,360	1,200	905	905	216	158	130	210
28.....	137	2,920	2,090	2,180	1,280	1,130	1,200	1,280	216	162	120	221
29.....	147	3,420	1,740	1,910		980	1,430	1,280	216	176	111	190
30.....	154	3,690	1,430	1,010		980	1,200	1,500	227	195	106	176
31.....	154		1,280	2,090		905		3,160		238	100	

ending Sept. 30, 1918-1922—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1921-22.												
1.....	170	17,900	4,830	1,130	1,280	2,280	4,540	1,820	1,500	1,740	402	402
2.....	190	9,180	3,160	980	2,090	3,040	4,540	1,580	1,280	1,500	449	480
3.....	170	3,970	4,830	980	5,410	12,700	3,690	1,580	1,680	1,580	480	830
4.....	176	2,280	5,120	1,130	4,250	8,600	2,920	2,690	1,660	2,090	425	1,430
5.....	176	1,580	4,830	1,430	3,290	5,410	2,280	8,020	1,740	7,150	362	1,580
6.....	176	1,280	3,600	1,580	2,580	3,690	2,000	9,180	2,920	4,540	321	1,130
7.....	170	080	3,160	1,500	2,280	5,700	1,740	6,280	2,800	2,090	273	718
8.....	167	802	2,800	1,280	1,740	10,300	1,680	4,540	2,000	1,010	260	660
9.....	180	704	2,680	1,280	1,500	7,440	1,430	3,420	1,740	1,430	300	402
10.....	200	830		1,130	2,090	10,600	1,360	2,380	1,580	1,000	344	203
11.....	190	1,130		1,130	4,540	12,400	1,280	2,000	1,430	816	449	200
12.....	171	1,280		1,000	9,180	8,310	1,130	1,740	3,070	062	362	264
13.....	162	1,130	720	1,060	8,020	6,570	1,060	1,580	5,410	524	300	238
14.....	154	980		1,060	5,120	5,120	1,280	1,280	3,420	486	254	249
15.....	147	830		1,000	3,290	7,440	2,380	1,200	2,480	760	286	216
16.....	137	718		1,060	2,690	13,900	5,120	1,200	2,000	020	550	238
17.....	126	602	1,280	1,280	2,480	8,890	3,970	1,130	1,740	678	1,500	227
18.....	120	2,180	3,290	2,280	2,480	5,700	2,920	2,920	1,580	1,130	830	200
19.....	114	2,480	12,400	8,600	2,920	4,540	2,580	5,900	1,280	1,740	760	176
20.....	108	3,970	7,440	8,890	6,860	3,160	3,040	4,540	2,000	1,200	449	158
21.....	103	5,120	4,250	8,310	20,700	2,690	2,600	2,920	2,920	080	321	158
22.....	100	3,420	2,690	13,900	14,500	2,280	2,480	1,820	1,820	802	286	151
23.....	106	2,480	8,600	10,600	9,180	2,000	2,280	1,740	1,580	718	254	151
24.....	114	2,000	17,600	7,150	5,990	1,820	1,910	1,910	1,430	905	260	144
25.....	126	1,580	18,200	3,070	4,250	1,740	1,740	1,740	1,130	662	1,500	137
26.....	120	1,430	12,700	2,800	3,160	1,680	1,060	1,580	760	524	2,690	130
27.....	117	1,910	8,800	2,280	2,600	1,010	1,580	1,740	3,420	402	1,740	120
28.....	111	3,070	6,120	1,820	2,480	7,440	1,580	3,160	3,690	524	1,130	123
29.....	103	16,700	3,040	1,580	10,000	2,280	2,580	2,380	718	802	123
30.....	108	9,180	1,910	1,430	0,860	2,000	2,180	2,000	578	662	123
31.....	371	1,430	1,280	3,970	1,820	440	524

NOTE.—Stage-discharge relation affected by ice Dec. 13-22, 1917, Jan. 4-11, 21-26, 1018, Dec. 24, 1910, to Jan. 8, 1920, and Dec. 10-10, 1921; mean discharge estimated by study of weather records. Afternoon gage reading of Dec. 31, 1917, increased 1 foot as it was obviously too low.

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the year ending Sept. 30, 1923.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	117	137	106	2,280	4,830	1,130	1,360	1,000	732	344	2,920	196
2.....	117	130	123	11,200	17,600	1,360	1,060	080	704	314	2,800	171
3.....	117	123	117	5,700	14,500	1,130	760	980	718	270	1,280	167
4.....	117	123	137	3,690	9,180	1,200	1,060	980	1,130	270	1,580	205
5.....	111	123	413	2,480	6,570	1,360	1,360	830	830	266	3,040	293
6.....	111	123	2,000	2,000	4,540	1,500	2,480	760	648	238	2,280	537
7.....	117	120	2,000	1,600	3,040	14,800	4,260	760	550	205	1,740	830
8.....	151	117	3,690	1,820	2,480	10,600	3,420	788	474	232	1,060	810
9.....	171	117	3,690	1,820	2,180	5,410	2,600	830	321	266	080	1,280
10.....	232	117	3,600	1,740	2,000	3,690	2,090	816	307	328	2,600	005
11.....	314	117	3,290	1,500	1,910	3,420	1,660	905	344	273	5,990	690
12.....	260	120	2,180	1,360	1,820	3,970	1,500	980	524	221	5,120	511
13.....	210	117	1,580	1,300	2,090	3,970	3,420	1,360	1,360	200	11,500	461
14.....	200	111	1,360	1,200	7,730	5,410	8,310	1,280	4,250	176	4,540	537
15.....	180	123	1,580	1,360	5,000	3,690	10,900	1,200	3,160	162	2,280	511
16.....	107	137	6,280	1,740	3,420	2,020	8,310	1,280	2,280	154	1,660	425
17.....	167	126	7,440	2,800	2,020	10,600	5,410	1,500	1,740	151	1,200	240
18.....	158	117	12,100	2,090	2,380	8,020	3,690	1,740	1,200	147	980	100
19.....	154	137	6,570	1,580	1,820	5,700	2,690	1,580	905	144	830	190
20.....	151	151	8,420	1,740	1,360	4,830	2,090	1,430	830	140	782	243
21.....	144	151	2,280	2,000	1,360	3,690	1,740	1,280	980	126	620	227
22.....	144	176	1,910	2,690	1,280	2,920	1,580	1,200	816	114	425	260
23.....	144	151	1,580	3,420	1,200	2,920	1,280	1,130	704	111	353	353
24.....	140	151	1,360	3,160	830	6,860	1,200	2,380	1,060	114	336	524
25.....	144	158	1,130	2,580	718	6,570	1,130	2,800	550	117	293	662
26.....	130	144	980	2,180	648	4,540	980	1,910	321	130	266	564
27.....	137	137	788	2,280	1,060	3,690	906	1,740	321	195	266	480
28.....	144	130	802	9,470	830	2,580	1,000	1,500	307	634	273	402
29.....	187	182	1,660	14,800	1,010	1,580	1,130	321	402	238	344

ending Sept. 30, 1924.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	206	154	2,580	6,570	1,820	3,090	6,860	1,360	4,830	502	210	062
2.....	232	151	2,690	6,570	1,580	3,420	4,830	1,910	3,160	537	195	537
3.....	200	144	2,000	0,570	1,580	3,160	3,160	1,580	2,480	486	336	486
4.....	185	301	1,660	15,100	1,500	2,690	2,280	1,430	1,910	413	402	437
5.....	176	802	2,920	8,020	1,430	5,120	3,160	1,280	2,090	371	371	391
6.....	162	062	6,570	5,120	1,500	8,890	3,970	1,200	2,000	344	293	353
7.....	151	746	5,410	3,040	2,480	8,310	5,410	1,130	1,360	578	254	362
8.....	144	788	3,600	2,090	2,480	5,700	5,700	1,130	1,130	980	232	321
9.....	140	620	2,020	1,580	1,010	3,690	4,830	1,280	1,200	4,830	210	336
10.....	137	511	2,580	1,430	1,740	3,160	3,970	1,360	5,120	2,480	205	203
11.....	140	425	2,090	1,430	1,500	2,600	3,420	3,420	7,150	1,430	200	300
12.....	144	371	1,910	8,310	1,300	2,180	2,800	25,000	4,540	1,130	344	426
13.....	151	321	1,740	5,990	1,280	1,820	2,480	20,700	2,920	905	005	353
14.....	140	293	1,660	4,250	1,130	1,580	2,090	12,100	7,150	3,420	905	321
15.....	133	266	1,500	2,600	980	1,430	1,740	9,180	4,830	2,180	648	293
16.....	123	254	1,300	3,040	905	1,430	1,580	7,440	3,070	1,580	480	200
17.....	117	314	1,130	21,300	830	1,500	1,430	5,120	3,160	1,360	353	328
18.....	126	321	1,060	0,700	1,060	1,910	1,740	3,690	2,380	1,130	279	273
19.....	140	293	1,060	5,120	1,500	2,580	3,070	2,020	1,820	905	221	243
20.....	154	266	980	3,970	2,000	3,160	4,540	2,580	1,500	690	1,360	221
21.....	162	243	005	3,000	6,800	3,100	3,420	2,600	1,280	504	3,070	210
22.....	180	227	1,000	3,160	5,120	3,160	2,920	2,920	1,060	511	3,160	216
23.....	105	273	1,360	2,280	3,290	2,920	2,480	2,690	830	499	1,580	1,200
24.....	105	437	2,000	1,530	2,480	2,480	2,090	2,480	704	620	1,280	1,060
25.....	200	409	2,380	1,740	2,000	2,020	1,000	1,910	802	690	7,440	005
26.....	205	1,280	2,230	2,280	1,910	3,290	1,580	1,660	704	499	5,410	670
27.....	190	1,280	2,090	2,000	2,090	4,540	1,360	2,280	620	362	3,420	511
28.....	205	788	2,090	1,000	2,690	5,120	1,200	2,800	564	293	2,090	461
29.....	105	718	0,470	1,430	5,700	14,200	1,200	2,800	537	200	1,660	502
30.....	185	005	5,410	1,360	19,800	1,200	10,300	648	243	1,280	12,400
31.....	171	3,690	1,660	10,300	7,730	227	802

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the year ending Sept. 30, 1925.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	13,000	271	062	1,280	1,280	2,800	5,120	415	400	118	80
2.....	6,570	242	620	1,500	2,200	1,200	2,380	4,250	395	437	115	75
3.....	3,040	218	564	1,740	1,200	2,000	3,420	376	305	115	77
4.....	2,090	212	511	2,090	2,180	1,200	2,000	2,920	336	356	131	77
5.....	1,580	212	405	1,010	2,280	1,130	1,010	2,480	303	318	162	76
6.....	1,280	206	306	2,000	1,800	1,130	1,910	2,280	271	287	184	75
7.....	1,130	200	405	1,910	3,290	1,060	1,740	2,000	1,000	264	104	75
8.....	1,060	188	5,120	1,740	3,970	1,060	1,500	2,090	905	287	184	75
9.....	080	760	13,300	2,280	4,540	1,060	1,280	2,280	802	648	175	72
10.....	302	436	7,440	3,420	6,860	080	1,130	2,480	690	592	170	71
11.....	076	437	4,250	4,830	8,600	980	1,200	2,690	578	564	162	71
12.....	550	395	3,160	5,700	9,760	030	1,130	3,000	511	537	162	71
13.....	511	356	2,530	5,120	8,310	905	080	6,570	461	511	153	71
14.....	486	318	2,180	3,420	5,700	905	005	4,250	415	486	153	77
15.....	437	287	1,820	2,180	3,970	830	774	3,690	376	437	148	87
16.....	395	830	1,580	1,910	6,860	774	704	3,160	461	395	144	95
17.....	346	713	1,430	3,040	7,440	690	634	2,480	620	356	144	95
18.....	270	511	1,740	0,280	5,000	006	788	2,090	005	318	141	95
19.....	236	718	1,430	9,470	4,540	4,540	732	1,740	980	287	137	95
20.....	200	502	1,280	8,020	3,420	13,000	573	1,430	830	250	137	95
21.....	270	449	1,500	6,570	2,480	8,020	437	1,230	746	230	134	95
22.....	303	550	1,430	4,830	2,000	3,160	740	1,060	537	206	131	95
23.....	270	4,540	1,280	3,420	1,740	1,910	980	980	425	184	131	95
24.....	250	3,070	1,130	2,690	1,580	1,740	1,000	830	690	179	128	95
25.....	218	2,920	2,180	2,180	1,500	1,060	1,200	788	1,280	170	122	95
26.....	104	2,180	1,740	2,000	1,430	1,580	1,280	732	1,360	166	115	95
27.....	170	1,530	1,430	1,360	1,430	1,580	676	1,360	157	109	95
28.....	425	1,200	1,200	1,280	1,430	2,380	620	1,060	153	98	101
29.....	405	080	1,060	1,900	1,740	3,070	564	774	144	92	101
30.....	336	816	1,060	2,280	5,700	511	606	137	87	106
31.....	303	1,130	2,800	461	128	84

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the year ending Sept. 30, 1926.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	101	2,480	718		5,120	3,420	1,740	1,200	2,000	200	153	816
2.....	101	2,280	676		5,090	2,600	1,680	1,060	2,090	170	1,200	704
3.....	104	2,090	620	530	4,250	2,480	1,580	980	1,010	166	980	578
4.....	106	1,910	564		3,600	2,000	1,910	980	1,740	148	000	090
5.....	112	1,740	634		2,800	1,740	1,010	905	1,580	134	385	400
6.....	118	1,740	676		2,280	1,580	1,910	816	1,430	2,380	205	850
7.....	125	1,910	020		2,000	1,680	1,740	732	1,300	2,280	250	437
8.....	134	3,040	578		2,000	2,280	1,660	076	1,200	1,360	200	578
9.....	144	3,690	550		1,820	3,290	1,580	034	1,060	980	230	564
10.....	157	3,160	524		1,820	2,380	1,430	620	080	537	248	524
11.....	179	2,920	499	730	1,000	2,090	1,660	592	980	648	104	440
12.....	206	2,690	474		1,580	2,090	2,000	606	905	437	153	396
13.....	248	2,480	449		2,280	1,910	3,100	578	005	336	131	330
14.....	310	2,280	437		10,300	1,910	3,970	550	788	287	112	270
15.....	396	2,090	437		16,700	1,910	3,600	564	670	256	128	248
16.....	537	1,010	425		8,600	1,820	3,160	524	578	224	200	242
17.....	606	1,740	415	802	4,330	1,740	2,800	511	437	184	310	224
18.....	670	1,580	415	2,580	3,420	1,060	2,280	511	356	106	499	200
19.....	760	1,430	415	5,120	3,420	1,660	2,690	409	318	144	005	200
20.....	810	1,360	405	19,800	4,540	3,070	2,280	461	326	141	4,830	188
21.....	006	1,200	020	11,500	3,970	5,700	1,010	437	326	131	5,700	184
22.....	980	1,130	1,360	16,000	3,160	4,830	1,740	415	310	115	3,100	170
23.....	1,000	1,060	5,120	12,700	2,020	4,250	4,250	305	295	100	2,090	157
24.....	1,130	980	4,540	5,700	2,800	5,700	3,600	376	205	128	2,000	144
25.....	1,360	005	2,280	3,690	2,800	5,120	2,920	678	270	218	1,010	128
26.....	5,410	830	1,280	2,480	8,600	5,700	2,090	504	486	310	12,100	212
27.....	5,120	830	980	2,090	6,570	7,730	1,740	486	440	340	6,280	300
28.....	4,250	802		1,910	4,250	8,600	1,500	802	366	205	3,420	340
29.....	3,690	774		1,580		4,250	1,430	676	303	212	1,740	310
30.....	3,100	746	540	1,060		2,380	1,360	634	248	153	1,360	408
31.....	2,690			1,580		1,910		676		125	1,060	

NOTE.—Stage-discharge relation affected by ice Dec. 28 to Jan. 10; discharge estimated from observer's notes and study of weather records.

Daily discharge, in second-feet, of Greenbrier River at Alderson, 1926-27.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	336	1,580	1,740	3,290	5,120	2,920	3,970	10,000	905	318	212	360
2.....	295	1,500	2,280	2,690	3,070	2,480	12,400	7,440	816	356	212	330
3.....	1,740	1,430	2,180	2,280	2,020	2,280	6,570	4,540	1,580	310	370	303
4.....	1,660	1,280	2,000	2,090	2,480	1,820	5,120	3,420	2,380	237	1,500	250
5.....	2,000	1,200	1,010	1,010	2,800	1,060	8,020	2,800	2,480	248	1,010	256
6.....	1,600	1,130	1,010	1,740	18,800	1,580	6,570	2,280	2,280	236	1,200	242
7.....	1,430	1,060	2,690	1,500	11,200	1,660	5,700	2,090	1,910	218	802	206
8.....	1,200	980	2,280	1,360	7,440	2,800	4,250	1,820	1,600	248	732	184
9.....	905	905	2,280	1,200	5,120	7,440	10,000	1,580	1,430	287	1,500	170
10.....	718	830	3,970		3,690	5,120	15,100	1,500	1,280	524	2,090	310
11.....	578	316	5,120		3,200	3,420	9,700	1,430	1,130	634	2,090	1,280
12.....	499	905	4,250		2,800	2,690	6,280	1,360	980	537	1,500	306
13.....	905	1,500	8,020		2,480	2,480	4,540	1,360	905	578	1,130	376
14.....	980	1,580	9,180	850	3,420	2,090	4,540	1,280	1,200	461	788	511
15.....	802	1,430	5,700		3,690	2,690	5,120	1,280	2,920	376	606	376
16.....	718	10,000	3,070		3,200	3,040	3,970	1,360	2,800	405	1,280	326
17.....	774	15,100	2,690		3,160	2,580	3,600	1,660	1,910	366	1,200	318
18.....	905	10,900	2,380	1,360	3,040	2,280	3,290	2,580	1,500	461	005	271
19.....	980	8,310	1,910	1,430	17,600	2,090	3,290	3,420	1,280	437	746	218
20.....	1,130	5,700	1,580	1,430	17,600	1,740	3,420	2,920	1,360	346	1,000	188
21.....	1,360	4,540	3,040	2,920	9,180	1,660	5,700	2,280	1,280	271	980	170
22.....	1,500	3,970	22,900	8,020	6,280	1,580	6,860	1,910	1,060	242	1,430	106
23.....	1,820	3,160	12,700	7,150	15,400	1,580	7,440	1,910	905	320	1,360	153
24.....	1,910	2,180	7,730	6,280	15,100	1,500	5,120	2,090	905	318	1,200	144
25.....	1,820	1,580	7,730	5,410	9,180	1,500	3,690	1,740	830	264	908	137

Daily discharge, in second-feet, of Greenbrier River at Alderson, 1927-28.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	112	303	746	6,860	1,360	1,500	3,160	16,400	1,060	5,120	287	1,360
2.....	112	295	760	3,970	1,200	1,360	2,580	11,000	080	2,920	264	1,360
3.....	106	318	080	4,250	1,060	1,200	2,280	0,570	816	2,690	242	1,200
4.....	157	502	2,920	4,540	1,060	1,060	2,000	4,250	802	2,090	300	1,280
5.....	148	2,480	3,420	3,600	1,130	080	1,820	2,800	830	1,580	474	1,060
6.....	184	1,600	3,070	1,010	1,280	005	1,660	2,180	1,660	1,280	395	080
7.....	179	502	3,290	1,580	1,660	830	1,580	2,090	3,160	980	366	905
8.....	212	1,280	5,700	1,500	1,910	774	1,500	1,820	2,130	830	425	905
9.....	184	2,580	5,120	1,430	4,250	700	1,660	1,580	1,740	746	400	788
10.....	170	2,000	3,290	1,280	3,070	1,060	1,580	1,430	1,580	2,690	376	070
11.....	184	2,600	2,300	1,200	3,040	1,360	1,580	1,280	1,820	2,090	1,910	490
12.....	218	2,380	2,480	1,060	2,480	1,360	3,420	1,130	1,820	2,000	1,360	405
13.....	1,820	2,000	2,380	005	2,000	1,280	3,600	980	1,660	2,020	005	360
14.....	2,920	1,580	6,280	905	1,910	1,280	3,160	005	1,580	0,370	690	326
15.....	1,360	1,360	5,120	980	7,440	1,580	2,800	830	1,500	6,000	400	303
16.....	080	1,130	5,120	1,130	6,570	1,600	2,280	760	1,280	2,380	1,500	287
17.....	760	1,740	6,860	1,130	3,970	2,280	1,910	732	1,130	1,580	8,760	287
18.....	550	8,330	5,120	1,130	3,040	3,070	1,740	930	1,130	1,280	5,410	271
19.....	437	6,720	3,290	4,540	2,580	3,040	1,580	905	080	1,060	3,690	376
20.....	486	3,420	2,280	4,100	2,090	2,480	1,430	905	5,700	816	2,480	3,070
21.....	1,500	2,380	2,090	11,900	1,820	2,580	1,280	905	9,800	648	1,660	3,690
22.....	2,000	1,820	1,740	7,730	1,500	3,420	1,200	788	4,540	504	980	2,180
23.....	1,500	1,500	1,500	3,070	1,600	11,600	1,200	830	3,160	409	830	1,580
24.....	1,130	1,360	1,200	2,280	2,090	10,100	1,500	1,060	3,290	425	788	1,130
25.....	980	1,200	1,060	3,420	2,380	7,440	1,660	1,130	3,690	366	600	802
26.....	746	1,060	802	4,250	2,000	5,700	2,580	1,360	3,420	318	034	690
27.....	606	980	732	3,040	1,660	4,250	2,580	1,010	2,480	205	1,280	611
28.....	511	905	537	2,280	1,740	3,290	5,410	1,740	1,820	320	3,040	449
29.....	437	816	830	1,740	1,060	2,580	5,000	1,580	1,500	310	2,090	415
30.....	366	774	1,660	1,500	3,160	12,700	1,360	3,040	303	1,430	440
31.....	326	2,690	1,360	3,690	1,200	287	1,280

Daily discharge, in second-feet, of Greenbrier River at Alderson, 1928-29.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	648	475	21,400	005	1,660	20,700	2,380	3,070	5,410	500	157	105
2.....	648	438	9,800	1,010	1,660	11,000	2,180	3,070	3,420	648	141	90
3.....	062	425	4,830	1,910	1,200	7,730	2,000	11,900	2,090	2,380	134	94
4.....	525	500	3,160	1,580	1,280	5,700	1,740	8,600	1,820	1,430	138	90
5.....	425	525	2,280	1,580	1,060	10,100	3,040	6,280	1,200	905	218	87
6.....	395	620	1,820	2,000	080	20,400	3,160	4,540	1,430	704	287	84
7.....	376	690	1,580	2,920	905	10,400	2,480	3,420	1,200	578	200	84
8.....	356	620	1,360	2,020	1,130	7,150	2,180	2,580	1,130	525	200	85
9.....	336	648	1,200	1,740	1,430	4,830	1,820	2,180	1,130	385	162	00
10.....	310	062	080	1,820	1,660	3,200	1,660	1,820	1,200	326	144	110
11.....	273	634	830	3,420	1,580	2,580	1,580	1,660	1,200	279	138	101
12.....	248	606	816	3,420	1,430	2,280	1,430	2,430	905	248	131	115
13.....	236	573	005	2,480	1,280	2,090	1,280	1,360	905	538	125	101
14.....	224	578	1,060	2,000	1,200	3,690	1,130	1,430	1,060	402	131	99
15.....	212	550	2,000	1,500	1,130	11,900	1,060	3,690	1,130	475	125	99
16.....	200	578	3,420	1,430	1,130	7,150	1,430	2,300	1,200	405	122	90
17.....	218	550	2,800	1,500	1,060	4,830	5,120	2,280	080	336	128	94
18.....	242	550	2,690	1,430	980	3,420	4,250	1,820	810	279	118	07
19.....	303	525	3,970	2,280	1,060	2,690	3,420	1,740	662	236	112	94
20.....	376	4,830	3,040	4,830	1,500	2,480	2,020	1,820	830	212	105	101
21.....	395	4,250	2,280	3,970	1,820	2,090	3,040	17,300	980	179	97	94
22.....	376	2,690	1,910	2,920	1,430	2,000	3,420	9,200	816	162	92	04
23.....	500	2,000	1,660	2,480	1,360	4,250	3,970	5,410	500	148	112	04
24.....	1,660	1,600	1,800	3,690	1,200	8,020	3,100	3,160	512	134	395	99
25.....	1,430	1,500	1,910	3,970	1,360	5,410	2,580	2,580	1,130	128	704	94
26.....	1,130	1,580	1,360	0,500	6,800	3,690	2,480	2,000	718	134	376	92
27.....	080	1,360	1,130	8,020	24,500	3,040	2,580	1,010	1,060	138	256	87
28.....	732	1,280	1,130	4,250	29,100	2,480	3,040	1,740	830	144	188	84
29.....	524	1,280	1,060	2,690	2,690	6,280	5,900	718	170	157	80	80

Daily discharge, in second-feet, of Greenbrier River at Alderson, 1929-30.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	2,090	2,460	1,620	3,200	620	2,080	880	570	237	123	38	38
2.....	7,150	2,220	1,350	2,970	742	1,800	850	000	225	123	36	38
3.....	12,200	6,470	1,180	3,020	915	1,670	863	620	202	110	38	35
4.....	6,410	12,200	1,020	2,070	3,720	1,400	1,120	631	188	97	36	35
5.....	2,600	7,150	954	2,460	8,700	1,210	1,460	570	173	84	31	35
6.....	1,740	4,080	1,080	2,050	8,100	1,160	1,740	550	173	78	30	31
7.....	1,280	2,780	1,200	1,860	6,100	1,380	5,400	570	192	73	30	31
8.....	980	2,110	1,700	1,760	3,640	8,100	6,450	031	188	70	33	33
9.....	760	1,690	2,300	1,630	2,870	7,650	6,400	760	188	64	38	36
10.....	550	1,440	2,570	1,500	2,380	4,500	4,080	064	207	68	31	35
11.....	462	1,250	2,340	1,440	2,050	3,260	3,230	590	448	56	28	31
12.....	415	1,180	2,120	1,270	1,720	2,940	2,660	560	1,790	53	30	30
13.....	310	1,100	2,220	1,180	1,510	2,770	2,220	520	1,050	51	33	33
14.....	279	1,160	2,530	1,160	1,580	2,660	1,860	502	708	61	35	33
15.....	271	1,400	2,510	1,230	1,000	2,750	1,650	403	630	73	43	33
16.....	236	1,040	2,200	1,240	1,480	2,510	1,510	493	430	01	48	40
17.....	224	2,110	1,970	1,130	1,210	2,300	1,290	550	345	63	48	40
18.....	200	20,000	1,810	1,020	1,160	2,080	1,130	540	206	53	43	33
19.....	200	21,300	3,620	860	1,290	2,220	1,110	610	262	58	53	36
20.....	188	8,700	6,600	814	1,210	3,330	1,040	928	231	58	43	33
21.....	194	6,100	4,350	742	1,160	3,100	954	915	212	70	61	35
22.....	1,980	3,280	2,820	941	1,130	2,530	928	708	197	73	01	36
23.....	5,840	2,460	2,420	1,100	1,310	2,080	980	590	178	61	58	35
24.....	4,830	2,050	2,120	850	1,680	1,770	915	602	151	04	61	33
25.....	2,500	1,770	1,670	766	2,610	1,560	838	430	139	56	61	35
26.....	1,810	1,630	1,480	742	3,120	1,400	778	370	127	48	68	43
27.....	1,380	1,440	1,320	826	2,970	1,410	730	331	123	48	56	40
28.....	1,120	1,600	1,950	876	2,460	1,190	675	310	116	51	53	35
29.....	965	1,720	6,700	876	1,050	631	280	110	56	48	30
30.....	980	1,680	5,250	730	967	600	269	120	51	45	30
31.....	1,600	3,810	664	941	250	43	43

Daily discharge, in second-feet, of Greenbrier River at Alderson, 1930-31.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	28	48	53	275	376	610	4,350	1,720	1,430	354	289	778
2.....	28	43	67	317	730	1,020	6,300	1,410	1,300	303	244	610
3.....	30	40	67	244	580	2,220	5,250	1,240	1,160	209	2,710	611
4.....	30	43	103	231	611	1,040	5,400	1,110	028	310	3,000	430
5.....	31	51	87	244	448	1,560	13,800	905	814	388	2,380	422
6.....	31	58	120	1,160	370	1,230	8,700	850	754	602	2,080	390
7.....	31	58	163	2,870	331	006	0,600	838	1,020	540	1,340	331
8.....	31	50	466	1,080	310	002	6,560	4,050	1,400	475	850	200
9.....	31	51	493	1,060	354	1,260	5,400	5,100	1,540	457	590	225
10.....	31	51	493	754	1,460	1,390	6,300	3,410	1,300	413	475	197
11.....	33	63	354	000	3,280	1,160	6,550	2,460	1,130	370	430	207
12.....	35	56	209	690	1,070	041	5,400	2,180	015	766	370	168
13.....	33	70	225	511	1,400	838	4,200	2,340	778	889	345	161
14.....	31	70	197	396	1,210	876	3,150	2,280	814	631	388	147
15.....	30	73	173	282	1,540	2,140	2,530	2,340	838	511	338	151
16.....	30	76	151	275	1,540	6,000	2,080	2,180	1,800	826	303	183
17.....	31	93	120	202	1,300	4,500	1,760	1,000	4,140	502	262	202
18.....	30	97	76	282	2,330	2,770	1,480	1,770	2,240	430	218	188
19.....	31	100	93	317	3,900	2,120	1,270	6,880	1,480	331	188	108
20.....	30	07	100	310	3,150	1,880	1,110	3,250	1,060	310	173	151
21.....	31	03	87	310	2,340	1,760	954	5,700	814	282	173	275
22.....	33	90	73	303	1,700	1,540	928	9,000	653	302	317	362
23.....	35	87	84	303	1,380	1,490	2,730	12,000	778	310	4,340	282
24.....	51	34	76	324	1,120	2,280	3,730	10,200	742	580	4,650	331
25.....	58	78	76	448	954	3,070	3,230	6,600	686	028	2,420	396
26.....	53	76	76	448	826	3,700	2,730	5,100	697	002	1,540	390
27.....	51	70	110	493	719	3,730	2,680	3,640	610	590	1,080	1,150
28.....	48	66	135	1,210	042	3,580	2,900	2,680	530	430	803	1,300
29.....	51	48	197	2,260	9,300	2,550	2,100	457	254	203	202

Daily discharge, in second-feet, of Greenbrier River at Alderson, 1931-32.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	580	143	202	1,670	7,050	1,180	12,300	15,200	430	2,460	207	76
2.....	457	143	475	5,250	4,350	1,080	9,600	21,800	388	1,600	231	76
3.....	370	159	675	5,400	5,400	980	6,450	8,550	362	1,130	269	116
4.....	317	163	502	3,530	15,000	1,150	4,350	5,100	331	850	282	120
5.....	282	151	404	2,400	37,100	2,220	3,100	3,500	310	12,700	310	116
6.....	250	143	345	2,140	12,800	3,870	2,530	2,640	296	14,800	289	188
7.....	225	135	324	6,300	6,750	6,000	2,300	2,100	345	11,000	256	147
8.....	207	135	310	7,950	4,500	4,500	1,970	1,740	303	6,000	207	147
9.....	192	131	303	8,700	3,550	3,280	2,030	1,480	282	3,810	173	123
10.....	183	123	354	7,350	2,820	2,530	3,780	1,490	269	2,380	147	100
11.....	173	120	1,860	4,050	2,380	1,880	3,700	2,360	244	1,790	135	00
12.....	163	120	2,730	3,200	2,420	1,720	3,120	5,800	256	1,380	163	106
13.....	150	120	4,020	2,400	3,330	1,580	2,700	9,600	331	1,050	159	113
14.....	148	120	3,390	2,050	2,820	1,440	2,200	6,900	1,000	802	135	87
15.....	130	120	3,840	1,760	2,320	1,210	1,900	4,800	1,080	664	178	70
16.....	135	116	2,440	1,540	2,050	1,040	1,610	3,470	730	600	108	61
17.....	135	116	2,220	1,380	2,220	2,930	1,440	2,570	653	580	159	53
18.....	135	116	1,610	1,240	3,330	16,000	1,270	2,070	550	403	155	56
19.....	131	123	1,210	1,110	3,390	9,900	1,160	1,700	570	413	163	48
20.....	131	127	080	980	2,700	6,450	1,060	1,410	600	370	244	51
21.....	131	123	838	876	2,220	5,100	954	1,210	778	317	192	45
22.....	135	123	954	802	2,130	6,750	876	1,240	1,610	289	178	48
23.....	139	110	2,430	754	2,400	9,000	850	1,320	876	262	202	48
24.....	127	116	2,820	754	2,100	7,050	700	1,130	642	256	188	51
25.....	123	116	2,590	814	1,770	4,650	802	915	475	250	173	48
26.....	120	110	2,220	1,040	1,540	3,280	2,030	790	370	244	127	51
27.....	120	110	1,770	1,100	1,390	2,680	2,750	697	331	250	116	127
28.....	116	110	1,390	1,380	1,310	17,500	2,260	708	6,690	231	103	116
29.....	123	116	1,260	2,030	1,230	19,300	1,900	610	13,400	102	93	100
30.....	135	143	1,150	8,110	9,900	1,650	530	5,250	183	87	93
31.....	135	1,080	13,600	8,250	484	183	78

Daily discharge, in second-feet, of Greenbrier River at Alderson, 1932-33.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	84	580	560	4,650	2,510	3,280	2,340	1,110	1,480	580	2,080	540
2.....	07	1,270	620	3,440	2,730	2,660	2,770	1,020	1,160	413	1,600	457
3.....	93	1,440	620	2,680	3,550	2,200	3,610	1,020	941	413	2,050	422
4.....	84	1,100	580	2,320	3,670	1,880	4,650	005	802	413	3,410	430
5.....	81	814	550	2,030	3,150	1,600	4,350	1,110	686	448	4,050	642
6.....	106	653	520	1,720	2,340	1,360	3,730	1,920	620	540	2,820	778
7.....	120	000	466	1,480	2,080	1,260	0,300	4,950	620	370	1,740	620
8.....	147	642	439	1,260	12,000	1,490	7,650	4,140	530	310	1,210	484
9.....	422	1,420	413	1,490	12,000	2,440	5,700	3,230	708	250	915	306
10.....	310	9,520	383	2,070	6,600	2,550	4,110	5,550	675	225	754	324
11.....	225	6,000	370	2,120	4,650	2,220	3,120	6,800	675	296	719	289
12.....	173	3,610	396	1,970	3,230	1,900	5,730	5,250	1,120	338	1,360	250
13.....	143	2,260	820	1,770	2,480	1,810	10,200	6,000	863	354	1,990	231
14.....	123	1,600	1,880	1,560	2,160	1,850	6,450	5,400	826	396	1,320	212
15.....	113	1,210	1,940	1,300	2,770	6,350	4,650	4,350	766	296	889	212
16.....	120	967	1,600	1,240	3,930	10,500	3,580	4,650	580	310	742	197
17.....	244	850	928	1,130	3,550	7,200	3,780	9,900	475	296	607	197
18.....	2,010	838	850	1,100	3,760	4,050	6,000	7,350	404	262	686	183
19.....	2,480	2,260	803	1,150	3,990	12,700	4,050	4,650	354	324	600	202
20.....	1,410	13,500	1,160	1,230	11,400	23,400	3,870	3,170	317	303	466	202
21.....	876	7,200	1,020	1,480	15,400	17,400	5,400	2,460	280	250	388	188
22.....	620	3,990	350	13,100	8,700	12,000	4,350	2,240	262	231	430	163
23.....	475	2,530	967	10,800	5,700	8,400	3,390	1,850	225	197	331	155
24.....	388	1,900	1,530	6,600	4,200	5,700	2,590	1,480	207	202	206	151
25.....	324	1,510	3,440	4,500	3,280	4,110	2,220	1,290	225	262	282	151
26.....	303	1,310	5,700	6,000	3,810	3,330	2,010	1,150	443	202	1,700	150
27.....	396	1,150	4,350	7,050	5,550	2,840	1,760	1,130	580	2,440	1,240	139
28.....	754	067	11,400	6,300	4,200	2,460	1,540	1,810	766	11,100	826	131
29.....	1,000	742	12,300	4,800	2,550	1,340	1,580	742	12,300	686	123

Daily discharge, in second-feet, of Greenbrier River at Alderson, 1933-34.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1.....	125	117	284	626	370	520	3,400	749	348	141	177	76
2.....	121	113	303	2,540	392	520	2,750	681	329	168	224	72
3.....	117	100	316	3,450	502	7,930	2,260	687	290	192	303	69
4.....	113	113	342	2,600	484	25,700	1,940	615	268	154	457	63
5.....	117	113	378	1,990	511	32,200	1,770	596	240	168	408	79
6.....	109	125	457	1,860	493	22,500	1,650	577	234	159	284	85
7.....	102	141	586	2,950	415	11,500	1,540	548	240	136	310	76
8.....	98	187	692	8,960	400	14,400	1,460	520	290	117	234	63
9.....	95	182	725	5,780	415	16,000	1,430	475	329	145	213	01
10.....	92	187	586	3,640	316	8,140	1,630	448	348	234	177	56
11.....	92	203	539	2,500	251	5,250	1,560	439	303	475	145	53
12.....	88	182	484	1,900	290	3,570	1,630	520	268	251	141	48
13.....	92	164	362	1,740	329	2,680	1,680	833	342	187	136	50
14.....	92	150	370	1,390	303	2,340	1,740	761	303	154	121	48
15.....	92	154	342	1,260	310	2,100	1,630	670	290	136	125	48
16.....	88	154	422	1,060	362	1,920	1,680	1,160	268	117	132	240
17.....	98	145	539	899	385	1,770	3,460	2,160	229	98	208	2,550
18.....	113	128	1,430	737	475	1,650	9,270	1,720	218	88	213	2,080
19.....	159	168	1,680	615	548	1,740	6,460	1,360	234	82	310	980
20.....	168	240	2,990	626	502	2,790	4,730	1,080	362	72	362	568
21.....	177	329	5,780	626	439	3,900	3,610	886	502	01	251	302
22.....	203	400	3,660	558	457	3,710	2,810	737	430	61	203	206
23.....	177	302	2,360	548	439	3,190	2,280	659	322	61	159	245
24.....	154	302	1,720	548	378	2,810	1,920	615	256	61	145	208
25.....	136	520	1,310	548	422	2,730	1,630	539	234	66	136	182
26.....	121	457	1,070	548	378	3,100	1,360	475	224	56	164	173
27.....	113	385	886	530	457	5,490	1,190	457	234	53	145	198
28.....	109	322	615	558	457	24,600	1,080	457	192	993	121	154
29.....	109	284	475	548	13,100	980	400	159	670	106	154
30.....	109	268	392	400	6,600	858	378	141	362	88	094
31.....	113	568	310	4,470	362	245	82

Monthly discharge of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1895-1917.
[Drainage area 1,340 square miles.]

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1895.					
August	390	125	252	0.188	0.22
September	755	82	164	.122	.14
1895-99.					
October	101	70	82.8	.062	.07
November	238	101	151	.113	.13
December	3,150	137	474	.354	.41
January	4,620	280	1,020	.701	.88
February	10,600	700	2,990	2.23	2.40
March	25,600	810	4,620	3.45	3.98
April	12,900	870	2,740	2.04	2.28
May	5,690	555	1,510	1.13	1.30
June	3,610	390	894	.667	.74
July	12,900	573	2,360	1.70	2.03
August	5,420	250	1,000	.701	.91
September	6,520	182	445	.332	.37
The year	25,600	70	1,530	1.14	15.51
1896-97.					
October	11,600	289	1,040	0.770	0.80
November	21,800	262	2,800	2.09	2.33
December	6,240	125	1,400	1.04	1.20
January	1,640	470	870	.049	.76
February	51,500	528	7,690	5.74	6.98
March	12,300	1,460	4,160	3.10	3.57
April	6,240	700	2,170	1.62	1.81
May	32,300	080	4,010	2.99	3.45
June	1,900	510	1,020	.761	.85
July	0,240	510	2,080	1.55	1.79
August	1,000	244	473	.353	.41
September	232	110	147	.110	.12
The year	51,500	110	2,290	1.71	23.15
1897-98.					
October	187	113	146	.109	.13
November	573	125	251	.137	.21
December	4,880	390	1,240	.925	1.07
January	15,700	620	3,500	2.08	3.00
February	5,690	755	2,000	1.49	1.55
March	16,400	1,070	3,690	2.75	3.17
April	13,300	1,550	3,970	2.96	3.30
May	16,400	1,070	3,520	2.03	3.03
June	1,300	430	761	.568	.63
July	2,930	238	705	.526	.61
August	42,900	430	4,390	3.28	3.78
September	510	209	300	.224	.25
The year	42,900	113	2,060	1.54	20.82
1898-99.					
October	23,000	204	1,920	1.43	1.65
November	10,300	810	2,420	1.81	2.02
December	8,810	630	2,210	1.65	1.90
January	18,600	1,380	3,320	2.48	2.86
February	18,600	755	4,750	3.54	3.69
March	45,300	2,190	8,420	6.23	7.24
April	6,240	870	2,060	1.54	1.72
May	14,000	650	2,950	2.20	2.54
June	2,500	390	1,030	.769	.86
July	336	165	219	.163	.19
August	810	75	157	.117	.13
September	280	95	155	.116	.13

Monthly discharge of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1895-1917—Continued.
[Drainage area 1,340 square miles.]

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1899-1900.					
October	157	75	105	.078	.00
November	268	113	156	.116	.13
December	1,000	125	345	.257	.30
January	9,700	1,090	1.26	1.45
February	16,000	390	3,390	2.53	2.64
March	17,100	1,220	5,010	3.74	4.31
April	3,850	700	1,850	1.38	1.54
May	1,720	301	661	.493	.57
June	7,360	268	1,140	.851	.95
July	4,100	133	813	.607	.70
August	1,000	110	226	.169	.19
September	414	70	145	.108	.12
The year	17,100	70	1,280	.955	12.99
1900-1.					
October	3,380	104	387	.289	.33
November	56,800	145	3,630	2.71	3.02
December	12,900	555	1,940	1.45	1.07
January	21,100	620	2,380	1.78	2.05
February	2,390	000	1,110	.828	.86
March	12,300	600	2,690	2.01	2.32
April	20,400	1,550	6,410	4.78	5.33
May	19,300	870	4,470	3.34	3.85
June	20,000	930	3,970	2.96	3.30
July	4,100	280	1,270	.948	1.09
August	2,500	238	849	.634	.73
September	2,090	280	693	.517	.58
The year	56,800	104	2,480	1.85	25.13
1901-2.					
October	315	153	228	0.170	0.20
November	422	119	188	.140	.16
December	36,700	238	4,850	3.62	4.17
January	12,900	470	2,630	1.96	2.26
February	29,600	600	4,410	3.29	3.43
March	30,700	1,640	7,500	5.60	6.46
April	7,940	755	3,170	2.37	2.64
May	2,030	390	828	.618	.71
June	4,100	301	982	.509	.57
July	930	165	414	.309	.36
August	280	66	138	.103	.12
September	164	70	91.4	.608	.08
The year	36,700	66	2,090	1.56	21.16
1902-3.					
October	415	76	151	.113	.13
November	3,100	51	456	.340	.38
December	7,360	598	2,730	2.04	2.35
January	23,100	576	3,840	2.87	3.31
February	24,900	1,720	6,880	5.13	5.34
March	31,100	1,160	6,300	4.70	5.42
April	10,000	1,920	3,410	2.54	2.83
May	2,620	315	870	.649	.75
June	8,810	396	1,750	1.31	1.46
July	2,500	128	760	.567	.65
August	788	58	238	.178	.21
September	301	70	161	.120	.13
The year	31,100	51	2,270	1.69	22.96

Monthly discharge of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1895-1917—Continued.
[Drainage area 1,340 square miles.]

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1903-4.					
October	246	101	149	.111	.13
November	246	70	150	.112	.12
December	490	101	183	.137	.16
January	15,400	217	1,580	1.18	1.36
February	0,400	246	2,330	1.74	1.88
March	10,000	1,420	3,660	2.73	3.15
April	8,810	659	2,280	1.70	1.90
May	15,700	788	2,800	2.00	2.41
June	7,860	396	1,360	1.01	1.13
July	1,160	140	352	.263	.30
August	188	86	141	.105	.12
September	188	46	86.4	.064	.07
The year	15,700	46	1,250	.933	12.73
1904-5.					
October	101	46	60.5	.045	.05
November	101	46	83.4	.062	.07
December	2,380	70	415	.310	.36
January	5,690	315	1,170	.873	1.01
February	1,520	246	693	.517	.54
March	25,600	1,000	7,030	5.25	6.05
April	3,600	855	1,610	1.20	1.34
May	28,400	720	3,920	2.93	3.38
June	5,960	246	1,020	.761	.85
July	12,900	544	2,510	1.87	2.16
August	1,080	315	616	.460	.53
September	928	101	287	.214	.24
The year	28,400	46	1,640	1.22	16.58
1905-6.					
October	1,620	70	274	.204	.24
November	598	188	321	.240	.27
December	11,800	490	2,420	1.81	2.09
January	21,100	1,080	4,000	3.72	4.29
February	2,140	315	841	.628	.66
March	13,300	544	4,750	3.54	4.08
April	7,940	1,240	3,490	2.60	2.90
May	2,500	855	1,360	1.01	1.16
June	2,860	396	1,070	.799	.89
1907.					
May 10-31	5,060	720	1,680	1.25	1.02
June	41,200	1,100	6,050	4.51	5.03
July	6,240	490	1,430	1.07	1.23
August	4,880	164	781	.583	.67
September	3,350	315	963	.719	.80
1907-8.					
October	7,940	280	976	.728	.84
November	9,400	306	3,470	2.59	2.89
December	19,600	598	4,260	3.18	3.67
January	21,800	1,420	4,610	3.44	3.97
February	39,600	1,000	5,270	3.93	4.24
March	25,300	1,520	7,700	5.75	6.63
April	26,000	2,040	5,450	4.07	4.54
May	21,500	1,720	5,280	3.94	4.54
June	5,150	396	1,300	.970	1.08
July	4,620	188	1,310	.978	1.13
August	2,140	217	675	.504	.58
September	443	101	183	.137	.15
The year	39,600	101	3,370	2.51	34.26

Monthly discharge of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1898-1917—Continued.
[Drainage area 1,340 square miles.]

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1908-9.					
October	720	101	183	.137	.16
November	598	140	261	.195	.22
December	5,150	164	1,080	.806	.93
January	11,300	1,240	3,870	2.89	3.33
February	12,300	788	4,130	3.08	3.21
March	10,600	1,520	4,100	3.06	3.53
April	15,000	1,160	3,790	2.83	3.16
May	7,080	855	3,280	2.45	2.82
June	2,140	490	1,080	.806	.99
July	3,600	188	753	.562	.65
August	2,380	140	422	.315	.36
September	544	101	193	.144	.16
The year	15,000	101	1,920	1.43	19.43
1909-10.					
October	1,520	86	391	.292	.34
November	1,720	188	447	.334	.37
December	9,700	246	1,190	.888	1.02
January	14,500	490	3,150	2.35	2.71
February	14,300	720	2,960	2.21	2.30
March	11,300	696	2,390	1.78	2.05
April	3,480	598	1,870	1.40	1.56
May	2,380	855	1,350	1.01	1.16
June	34,500	747	5,750	4.29	4.79
July	4,880	380	1,250	.933	1.08
August	696	178	268	.200	.23
September	1,820	164	610	.455	.51
The year	34,500	86	1,790	1.34	18.12
1910-11.					
October	622	178	284	.212	.24
November	788	150	230	.172	.19
December	8,660	200	1,010	.754	.87
January	35,100	1,160	7,040	5.25	6.05
February	9,109	1,130	2,910	2.17	2.26
March	9,400	1,240	4,380	3.27	3.77
April	17,800	1,720	5,780	4.31	4.81
May	1,720	315	747	.557	.64
June	971	315	522	.390	.44
July	471	101	231	.172	.20
August	443	86	196	.146	.17
September	3,850	246	975	.728	.81
The year	35,100	86	2,020	1.51	20.45
1911-12.					
October	18,200	315	2,600	1.94	2.24
November	10,300	415	2,200	1.64	1.83
December	6,800	708	2,200	1.64	1.89
January	10,000	2,230	1.66	1.91
February	17,800	659	3,510	2.62	2.83
March	31,109	1,400	6,950	5.19	5.98
April	9,700	1,140	2,960	2.21	2.47
May	18,200	587	4,010	2.99	3.45
June	2,040	188	498	.372	.42
July	6,800	396	1,040	.776	.89
August	490	105	213	.159	.18
September	2,980	68	395	.295	.33
The year	31,109	68	2,409	1.79	24.42

Monthly discharge of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1895-1917—Continued.
[Drainage area 1,340 square miles.]

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1912-13.					
October	400	145	257	.192	.22
November	5,690	183	759	.506	.63
December	12,900	178	1,050	.784	.90
January	14,300	1,240	4,270	3.19	3.68
February	7,300	774	2,100	1.63	1.70
March	42,500	870	0,430	4.80	5.53
April	19,300	013	3,900	2.91	3.25
May	15,700	544	2,770	2.07	2.39
June	5,690	356	1,660	1.24	1.38
July	8,230	380	1,150	.858	.99
August	2,860	164	645	.481	.55
September	443	113	189	.141	.16
The year	42,500	113	2,110	1.57	21.88
1913-14.					
October	7,080	164	1,200	.806	1.03
November	14,800	380	2,930	2.19	2.44
December	6,520	696	2,270	1.69	1.95
January	9,700	1,230	3,280	2.45	2.82
February	14,000	1,130	4,420	3.30	3.44
March	13,300	1,480	4,670	3.40	4.02
April	12,600	1,920	4,780	3.57	3.08
May	3,220	308	1,220	.910	1.05
June	659	140	241	.180	.20
July	1,000	164	305	.228	.26
August	1,130	124	287	.214	.25
September	512	95	217	.162	.18
The year	14,800	95	2,130	1.59	21.62
1914-15.					
October	1,080	64	270	.201	.23
November	311	111	167	.125	.14
December	8,900	169	3,660	2.73	3.15
January	26,300	1,080	7,030	5.25	6.05
February	27,800	1,340	6,320	4.72	4.92
March	2,500	700	1,330	.003	1.14
April	1,250	526	802	.599	.67
May	1,250	239	606	.452	.52
June	7,500	179	1,300	.970	1.08
July	374	122	223	.166	.19
August	916	120	456	.340	.39
September	1,000	158	431	.322	.36
The year	27,800	64	1,860	1.39	18.84
1915-16.					
October	18,500	253	1,730	1.29	1.49
November	2,090	179	634	.473	.53
December	18,500	294	2,450	1.83	2.11
January	13,400	1,160	3,650	2.72	3.14
February	8,050	1,250	3,530	2.63	2.84
March	7,500	1,510	3,250	2.43	2.80
April	6,680	1,160	2,600	1.94	2.16
May	5,580	452	1,250	.033	1.08
June	12,200	526	2,370	1.77	1.98
July	3,480	423	1,120	.836	.96
August	7,220	328	1,620	1.21	1.40
September	2,810	164	409	.305	.34
The year	18,500	164	2,050	1.53	20.83

**Monthly discharge of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1895-1917—Concluded.**
[Drainage area 1,340 square miles.]

Month	Discharge in second-feet				Run-off in inches
	Maximum	Minimum	Mean	Per square mile	
1916-17.					
October	2,190	225	509	0.447	0.52
November	902	179	306	.228	.25
December	15,200	277	1,000	1.10	1.37
January	13,700	1,080	3,580	2.07	3.08
February	12,800	738	3,480	2.60	2.71
March	27,200	1,990	9,000	7.39	8.52
April	6,950	777	2,090	1.56	1.74
May	12,500	615	2,450	1.83	2.11
June	2,600	204	940	.701	.78
July	3,030	118	462	.345	.40
August	846	74	216	.101	.19
September	649	79	164	.122	.14
The year	27,200	74	2,150	1.60	21.81

**Monthly discharge of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1918-1922.**
[Drainage area 1,340 square miles.]

1917-18.					
October	4,760	100	515	0.384	0.44
November	2,600	148	447	.334	.37
December	1,600	481	.359	.41
January	11,000	1,700	1.27	1.46
February	16,100	1,600	7,500	5.60	5.83
March	48,000	1,740	7,100	5.30	0.11
April	12,700	1,500	5,720	4.27	4.76
May	6,280	1,580	2,520	1.88	2.17
June	17,600	314	2,510	1.87	2.09
July	6,280	449	1,340	1.00	1.15
August	2,090	273	802	.598	.69
September	7,730	449	1,520	1.13	1.26
The year	48,000	100	2,640	1.97	20.74
1918-19					
October	24,100	238	1,700	1.31	1.51
November	10,900	676	2,100	1.61	1.80
December	17,000	1,130	4,240	3.16	3.64
January	39,400	1,660	6,310	4.71	5.43
February	8,310	760	2,100	1.63	1.70
March	13,000	1,300	3,510	2.62	3.02
April	7,730	1,280	2,520	1.88	2.10
May	14,800	1,660	4,660	3.48	4.01
June	13,900	606	2,480	1.85	2.06
July	13,900	830	3,480	2.60	3.00
August	1,740	227	641	.478	.55
September	788	151	258	.193	.22
The year	39,400	151	2,870	2.14	29.04
1919-20					
October	3,970	140	787	.587	.68
November	6,470	413	1,810	1.35	1.51
December	22,200	4,040	3.01	3.47
January	19,200	4,010	2.99	3.45
February	6,570	1,430	2,580	1.93	2.08
March	21,900	1,500	4,950	3.69	4.25
April	9,760	1,580	4,530	3.38	3.77
May	5,700	1,060	1,810	1.35	1.56
June	10,300	499	1,870	1.40	1.56
July	2,920	243	908	.678	.78
August	6,990	180	930	.701	.81
September	480	130	223	.166	.19
The year	22,200	130	2,370	1.77	24.11

STREAMS	Square Miles.
Meadow Creek.....	23.21
Laurel Run.....	4.96
North Fork of Anthony Creek (entire).....	22.89
North Fork of Anthony Creek (above Poeahontas line)	7.20
Onemile Run.....	1.62
Twomile Run.....	2.30
Fourmile Run.....	1.56
Hoffman Run.....	1.04
Coles Run.....	1.99
Pondliek Run.....	0.67
Sugar Run.....	1.92
Bear Run.....	1.08
Laurel Run.....	12.31
Boardhouse Run.....	1.52
Spring Creek.....	76.31
Dry Run.....	5.23
Robbins Run.....	11.41
Boggs Run.....	2.24
Roekeamp Run.....	2.75
Panther Camp Creek.....	4.88
Board Liek Run.....	1.40
Wolfpen Run.....	1.98
Boggs Run.....	2.36
Big Run.....	1.49
Snodgrass Run.....	3.49
Slabeamp Run.....	5.81
Red Run.....	0.91
Kineaid Run.....	2.29
Davy Run.....	2.90
Splee Run.....	8.38
Milligan Creek (surface area only).....	25.87
Culverson Creek (surface area only).....	35.13
Spiee Run.....	2.41
Burns Run.....	2.78
Indian Creek.....	2.59
Sinking Creek (surface area only).....	41.39
Hughart Creek.....	7.55
Flynn Creek.....	2.59
Roaring Creek (surface area only).....	7.94
Little Roaring Creek.....	2.83
Meadow River (entire).....	360.77
Meadow River (above Nicholas County line).....	282.16
Anglins Creek (entire).....	32.98
Anglins Creek (in Greenbrier County).....	4.21
Youngs Creek.....	10.96
North Prong.....	1.78
Spring Creek.....	1.01
Haynes Branch.....	0.67
Burdette Creek.....	5.46
Piney Creek.....	1.58
Toms Creek.....	3.20
	10.40

Areas of Drainage Basins (Continued).

STREAMS	Square Miles.
Sewell Creek (entire).....	40.55
Sewell Creek (in Greenbrier County).....	22.45
Little Sewell Creek.....	16.82
Boggs Creek.....	10.16
Wolf Pen Creek.....	2.93
Little Creek.....	3.54
Laurel Creek.....	5.91
Mill Creek.....	6.94
Big Clear Creek.....	51.89
Brown Creek.....	6.90
South Fork.....	18.90
Smokehouse Branch.....	3.18
Old Field Branch.....	4.27
Old Knob Branch.....	4.38
Sam Creek.....	2.74
Elijah Branch.....	1.72
Road Branch.....	1.18
North Fork.....	5.31
Little Clear Creek.....	33.40
Beaver Creek.....	7.20
Stony Run.....	1.34
Rader Run.....	2.10
Laurel Creek.....	4.46
Kuhn Branch.....	2.59
Otter Creek.....	15.69
Methodist Branch.....	2.27
Smoot Branch.....	1.76
Eagle Branch.....	2.98
Buffalo Creek.....	5.78
Morris Branch.....	3.69
Patterson Creek.....	3.18
(Gauley River)	
Hominy Creek (entire).....	104.81
Hominy Creek (in Greenbrier County).....	10.98
Price Fork.....	2.93
Peaser Branch.....	2.30
Cherry River (entire).....	171.90
Cherry River (in Greenbrier County).....	122.64
Laurel Creek (entire).....	42.02
Laurel Creek (in Greenbrier County).....	32.77
McMillion Creek.....	3.56
Mill Branch.....	1.44
Beech Run.....	3.18
Hogcamp Run.....	2.58
Manning Branch.....	1.91
Middle Branch.....	2.37
Cold Spring Branch.....	2.95
Linn Branch.....	2.12
Little Laurel Creek (entire).....	18.40
Little Laurel Creek (in Greenbrier County).....	14.26
Baber Branch.....	1.08
Improvement Branch.....	2.21

Areas of Drainage Basins (Concluded).

STREAMS	Square Miles.
South Fork of Cherry River (in Greenbrier County)	55.18
Shiras Run.....	1.65
Elklick Run.....	1.18
Rooky Run.....	8.46
Little Rocky Run.....	1.81
Becky Run.....	3.46
Cold Knob Fork.....	8.77
Blizzard Run.....	1.23
Little Blizzard Run.....	0.65
Big Run.....	1.05
North Fork of Cherry River (entire).....	36.98
North Fork of Cherry River (in Greenbrier County)	20.43
Coats Run.....	1.84
Little Lick Run.....	1.21
Windy Run.....	0.74
Armstrong Run.....	0.64
Hamrick Run.....	0.78
Rabbit Run.....	0.69
Carpenter Run.....	0.93
Deacon Run.....	0.61
Fallen Timber Run.....	0.78
Bear Run.....	3.66
Dogway Fork (of Cranberry River).....	9.78
Dogway Fork (in Greenbrier County).....	0.45

DESCRIPTION OF DRAINAGE BASINS.

Greenbrier River.—Greenbrier River, the stream that carries the greater part of Greenbrier County's rainfall, has its source in two forks heading in the extreme northern end of Pocahontas County. West Fork heads east of Shavers Mountain about two miles northeast of Wildell with an elevation of 3,625 feet. East Fork heads at Blister Swamp on the west slope of Allegheny Mountain with an elevation of 3,875 feet and flows in a southwest direction to join the West Fork at Durbin where it makes the Greenbrier River proper. The Greenbrier flows in a comparatively straight line in a southwest direction across Pocahontas and Greenbrier Counties to a point south of Lewisburg where it turns westward to form part of the Greenbrier-Monroe County line. Here it enters Summers County and after much meandering joins New River

From the Table of Stream Data on page 39, it can be seen that from its mouth to its East Fork source it has a meandering length of 164.8 miles with an air-line distance of 98.64 miles, or a ratio of 1.67. It has a total fall of 2,500 feet or at a rate of 15.2 feet per mile. From its mouth to its West Fork source it has a meandering length of 162.9 miles with an air-line distance of 97.14 miles or a ratio of 1.67 also. The fall is much more rapid near its source than at the mouth as the following gradient table shows:

Gradient of Greenbrier River.

	Miles.	Elevation.	Fall. Feet.	Rate per mile. Feet.
Source of East Fork.....		3875		
Distance	18.8		1175	62.5
Durbin (River forks).....		2700		
Source of West Fork.....		3625		
Distance	16.9		925	54.7
Durbin (River forks).....		2700		
Distance	14.9		275	18.4
Cass		2425		
Distance	9.0		155	17.2
Clover Liek.....		2270		
Distance	16.8		155	9.2
Marlinton		2115		
Distance	21.5		162	7.53
Pocahontas-Greenbrier line.....		1953		
Distance	22.7		173	7.62
Anthony		1780		
Distance	17.6		135	7.67
Ronceverte		1645		
Distance	15.1		120	7.94
Alderson		1525		
Distance	28.4		150	5.28
Mouth (Bellepoint) (empties into New River 1½ miles south of Hinton)		1375		

According to Reger¹⁷ Greenbrier River has a total drainage area of 1629.43 square miles. In Greenbrier County it has a drainage area of 679.02 square miles. The principal tributaries in Greenbrier are Muddy Creek, Second Creek, Howard Creek, Anthony Creek, and Spring Creek.

A gaging station was established on the Greenbrier River at Alderson August 1, 1895, by C. C. Babb and D. C. Humphreys, of the United States Geological Survey, and since that date until the present time, with few interruptions the gage has been read daily by local observers. Mr. W. J. Hancock and Mr. W. C. England are accredited with most of this detail. Prior to October 15, 1929, the gage was located at the highway bridge at Alderson, half a mile above the mouth of Muddy Creek and thereafter 400 feet above the bridge. The non-recording gage was read to half tenths once daily prior to April 1, 1910; to half tenths twice daily, April 1, 1910 to December 31, 1911; to hundredths twice daily January 1, 1911 to October 14, 1929; recording gage thereafter. Zero of gages is 1,528.97 feet above mean sea-level. Channel described as practically permanent at the bridge and as "shifts occasionally" at the recording gage. Affected during 1914-15 by construction of new highway bridge. Sometimes affected by ice. Rating well defined to about 25,000 second-feet. Discharge measurements have been made from time to time by government officials, the work having been done partly in cooperation with the West Virginia Geological Survey.

The records for the years 1895 to 1935 are taken directly from the United States Water-Supply Papers, as follows:

1895-1920, from No. 536, pages 177-196.
1920-1922, from No. 543, pages 68-71.
1922-1923, from No. 563, pages 57-59.
1923-1924, from No. 583, pages 94-95.
1924-1925, from No. 603, pages 109-110.
1925-1926, from No. 623, page 95.
1926-1927, from No. 643, page 58.
1927-1928, from No. 663, page 62.
1928-1929, from No. 683, page 71.
1929-1930, from No. 698, page 75.
1930-1931, from No. 713, page 77.
1931-1932, from No. 728, page 113.
1932-1933, from No. 743, page 111.
1933-1934, from No. 758, page 119.
1934-1935, from No. 783, page 116.

Date	Made by—	Gage h	Disch	Date	Made by—	Gage h	Disch
1895.		Feet.	Sec.-ft.	1908.		Feet.	Sec.-ft.
July 30	O. C. Babb.....	1.80	457	Apr. 21	Follansbee and Bar-		
Sept. 4	D. C. Humphreys	1.36	106		rows	8.04	3,880
1890.				Aug. 4	W. C. Hoyt.....	2.15	425
June 10do	2.80	714	Aug. 8do	2.20	456
July 22do	2.07	065dodo	2.88	1,400
Aug. 13do	1.05	520	Sept. 18	W. M. O'Neill.....	1.70	176
Dec. 18do	2.80	1,480	1009.			
1897.				Nov. 10	A. H. Horton.....	2.12	413
Mar. 30do	2.75	1,640	Dec. 1	C. L. Parket.....	2.00	334
May 8	F. H. Anschutz.....	4.88	6,180	1910.			
4do	4.88	5,450	Aug. 23	J. C. Dort.....	1.78	163
4do	4.29	4,960	Oct. 11	C. T. Bailey.....	2.20	572
0do	3.50	3,160	1011.			
18do	12.30	32,000	Nov. 4	Bailey and Perwien..	2.22	513
14do	12.32	33,000	1912.			
14do	0.52	20,700	Mar. 24	C. T. Bailey.....	5.88	9,200
31do	2.26	681	1913.			
Oct. 12do	1.42	71	June 20	H. J. Jackson	2.17	450
1898.				1914.			
June 26	D. C. Humphreys....	2.02	403	Dec. 1	J. C. Mathers.....	1.84	152
Aug. 3do	2.20	665	1015.			
Aug. 6do	4.43	5,120	Feb. 6	W. Kessler	4.76	6,040
Dec. 22do	3.45	2,820	1016.			
1899.				Mar. 23	A. H. Horton.....	4.40	4,700
June 22do	1.04	456	Mar. 24do	4.05	5,000
Aug. 12do	1.36	104	Aug. 15	B. E. Jones.....	2.65	1,000
1900.				1017.			
Mar. 31do	4.07	5,130	May 29do	5.55	8,060
June 20do	1.90	504	30do	4.06	5,470
July 24do	1.97	493	31do	3.94	3,500
Aug. 20do	1.38	148	1918.			
Dec. 21do	2.24	834	Feb. 15	B. L. Hopkins.....	6.05	11,800
1901.				15do	7.15	12,400
Mar. 27do	3.45	3,140	10do	8.40	16,400
July 30do	1.73	276	18do	4.64	5,330
1902.				Apr. 15do	6.16	8,840
July 17do	1.03	211	May 15do	3.97	2,750
Aug. 12do	1.46	150	June 23do	3.64	2,000
12do	1.46	154	1020.			
1903.				May 21	Peterson and Bigwood	3.36	1,660
Sept. 21	Paul and Sawyer.....	2.03	373	June 23	B. L. Bigwood.....	3.77	2,400
Nov. 16	W. C. Sawyer.....	1.73	130	Nov. 21	Bigwood and Lam-		
1004.					oureux	2.80	031
June 15	F. H. Brundage.....	2.20	400	1922.			
Aug. 0	N. C. Grover.....	1.72	146	Feb. 24	Dirzulaitis and Big-		
Sept. 20	R. H. Bolster.....	1.63	114		wood	5.20	6,740
Oct. 1do	1.44	51	1923.			
20do	1.51	76	Apr. 4	J. J. Dirzulaitis.....	2.98	1,190
1005.				1924.			
Mar. 22	A. H. Horton.....	7.90	16,500	Oct. 18do	2.42	330
22do	7.17	13,700	Oct. 18do	2.43	338
23do	5.24	7,240	1025.			
23do	5.09	8,840	June 30do	2.64	602
Sept. 16	R. H. Bolster.....	2.01	350	Oct. 2do	1.00	101
1906.							
June 13	Robert Follansbee ..	2.30	002				

1.....	220	116	11.....	280	301	22.....	250	116
2.....	301	105	12.....	280	250	23.....	250	110
3.....	250	145	13.....	202	220	24.....	280	110
4.....	220	125	14.....	350	182	25.....	220	05
5.....	192	125	15.....	280	125	26.....	155	05
6.....	155	155	16.....	280	145	27.....	250	90
7.....	125	157	17.....	280	137	28.....	315	82
8.....	182	157	18.....	250	125	29.....	280	82
9.....	238	145	19.....	250	125	30.....	430	250
10.....	220	155	20.....	250	125	31.....	350	250

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1805-96.												
1.....	82	101	280	1,000	755	1,070	12,900	2,000	573	1,000	5,420	238
2.....	70	107	262	2,000	755	1,550	11,600	2,010	435	755	3,610	220
3.....	70	110	220	1,300	5,800	1,380	8,230	2,000	430	700	5,150	102
4.....	70	125	204	755	10,000	1,070	5,150	4,520	414	500	3,150	220
5.....	70	126	204	1,000	5,090	870	3,510	5,500	300	870	2,000	102
6.....	82	137	155	930	3,150	870	2,290	3,380	510	810	1,380	220
7.....	75	145	137	430	5,240	810	2,090	2,390	555	1,140	1,070	220
8.....	88	157	157	454	4,520	930	1,810	1,810	700	2,000	870	220
9.....	88	145	182	470	3,150	1,220	1,000	1,640	700	12,900	755	238
10.....	70	153	182	000	2,710	1,380	2,000	1,300	700	10,000	500	244
11.....	70	153	200	555	2,190	1,220	2,090	1,000	1,380	4,880	500	220
12.....	82	153	209	555	1,640	1,720	2,100	870	1,070	2,820	600	204
13.....	82	145	192	555	700	1,900	2,000	755	700	2,000	510	192
14.....	78	137	165	810	9,100	1,450	1,900	755	000	1,640	430	182
15.....	78	145	137	700	8,230	1,540	1,000	1,300	510	1,220	470	102
16.....	78	133	137	280	4,620	1,810	1,810	1,140	555	1,070	600	470
17.....	82	137	165	280	3,150	12,300	1,460	810	582	030	510	315
18.....	82	137	157	294	2,000	3,520	1,220	600	600	030	390	238
19.....	82	137	155	301	1,040	5,420	1,140	555	555	870	390	204
20.....	82	137	165	301	1,220	10,000	1,070	555	582	810	350	315
21.....	82	145	192	366	930	7,850	930	810	555	650	315	280
22.....	82	101	238	300	930	5,420	870	930	480	000	250	280
23.....	82	170	3,150	430	930	5,150	870	1,070	470	573	301	244
24.....	90	182	1,300	1,300	755	4,620	030	1,070	430	1,220	454	274
25.....	101	182	930	4,520	1,000	4,020	2,000	1,220	470	1,300	470	220
26.....	05	182	700	3,850	870	3,510	2,000	1,300	2,290	7,040	478	182
27.....	95	165	500	2,390	810	8,510	1,000	1,140	3,610	3,010	454	182
28.....	95	182	755	1,040	700	8,380	1,550	1,000	2,200	2,000	400	102
29.....	95	209	1,380	1,220	810	2,100	1,330	030	2,000	1,550	280	232
30.....	95	238	030	930	25,600	1,300	810	1,550	1,140	280	6,520
31.....	05	810	810	19,500	070	3,010	262
1800-97.												
1.....	11,500	202	5,240	510	700	2,000	1,460	080	1,140	537	870	232
2.....	4,100	301	3,610	470	700	1,720	1,460	2,820	1,300	2,710	755	209
3.....	2,290	280	2,710	494	700	1,550	1,300	5,240	1,000	4,880	1,000	209
4.....	1,550	350	2,000	510	555	1,400	1,300	5,150	870	2,710	1,000	192
5.....	1,140	15,400	1,550	600	528	2,190	1,550	3,850	1,070	1,040	870	155
6.....	030	21,800	1,300	1,040	4,100	2,190	0,240	3,150	1,000	2,710	810	165
7.....	700	16,200	1,220	1,550	11,600	2,710	4,620	2,930	930	2,510	580	182
8.....	600	3,150	1,070	1,000	7,360	3,610	3,150	2,390	810	2,390	550	165
9.....	510	2,190	1,000	870	4,620	2,930	2,930	2,000	755	1,810	555	157
10.....	470	1,640	1,300	870	3,150	3,150	5,240	1,720	810	1,460	510	165
11.....	430	1,220	2,090	870	2,820	6,240	5,420	1,640	755	1,000	510	161
12.....	300	1,070	1,810	700	4,360	5,420	4,100	2,100	670	1,380	480	157
13.....	700	930	1,550	555	7,940	5,590	2,930	17,500	700	1,220	438	141
14.....	650	810	1,220	510	7,080	5,150	2,290	32,300	755	930	300	141
15.....	030	755	1,220	470	6,800	10,000	2,000	9,700	700	810	374	137
16.....	500	550	1,900	700	7,080	7,080	1,900	6,240	1,000	070	374	133
17.....	510	600	1,900	700	7,080	4,350	1,810	4,100	1,300	582	350	125
18.....	470	555	1,640	700	5,150	3,610	1,810	3,150	1,140	510	336	125
19.....	430	537	1,300	1,220	5,420	9,700	1,550	2,890	1,000	500	815	119
20.....	390	510	1,140	1,300	5,150	12,300	1,460	2,000	1,720	050	294	119
21.....	336	510	1,000	1,140	5,150	10,300	1,380	1,720	1,810	870	280	125
22.....	315	591	870	1,140	24,100	4,620	1,220	1,550	1,550	6,240	262	125
23.....	301	700	755	1,220	51,500	4,100	1,070	1,380	1,300	5,690	202	110
24.....	280	700	700	1,220	20,700	3,150	1,000	1,220	1,070	4,100	350	116
25.....	280	700	430	1,000	9,100	2,820	870	1,070	1,000	4,100	301	119
26.....	201	680	280	1,070	5,690	2,390	870	1,070	1,070	2,500	329	125

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1895-1917—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1897-98.												
1.....	187	125	510	810	1,550	1,220	7,940	1,040	1,800	470	700	890
2.....	119	170	430	870	930	1,220	6,420	1,380	1,070	406	1,800	800
3.....	125	167	890	1,400	755	1,140	4,360	1,220	030	358	755	238
4.....	125	209	390	700	930	1,070	3,380	1,140	810	836	700	316
5.....	145	192	810	620	1,550	1,070	2,930	1,070	700	510	10,300	208
6.....	141	220	2,300	650	930	1,140	2,930	2,000	050	870	7,940	316
7.....	129	220	2,190	080	930	1,140	2,710	16,400	578	600	7,940	350
8.....	125	250	1,380	070	1,070	1,140	2,500	14,000	528	504	8,380	510
9.....	125	232	1,000	090	930	1,220	2,500	7,300	494	470	2,090	430
10.....	119	220	756	8,150	1,070	1,300	2,300	4,620	470	406	14,000	890
11.....	113	220	555	12,900	1,040	1,220	4,100	2,290	430	865	42,900	350
12.....	125	220	655	6,520	2,100	1,300	13,300	2,820	870	230	12,900	315
13.....	129	450	404	4,620	3,150	1,380	7,650	2,190	630	238	7,360	280
14.....	133	288	470	4,100	2,820	1,460	6,000	1,900	528	301	4,880	250
15.....	137	815	486	8,150	2,290	1,400	5,090	1,040	1,140	280	3,150	250
16.....	137	294	870	15,700	2,000	1,380	5,420	1,720	1,000	343	2,190	238
17.....	137	308	2,000	7,940	1,040	1,300	6,800	1,810	980	414	1,640	238
18.....	116	208	1,400	4,520	1,380	7,360	4,880	2,010	930	430	1,300	209
19.....	176	250	1,220	3,150	1,550	9,400	3,610	2,290	870	1,000	1,970	209
20.....	165	252	1,140	2,390	3,150	7,360	2,710	2,090	1,140	870	1,140	209
21.....	176	250	1,380	2,300	4,100	4,620	2,290	1,810	1,220	700	1,300	209
22.....	182	262	4,880	2,610	6,690	3,380	1,900	1,550	1,000	537	1,070	220
23.....	165	262	3,610	3,380	3,850	2,710	1,720	6,800	930	402	870	260
24.....	165	220	2,290	5,090	2,820	2,390	1,560	8,230	650	555	700	280
25.....	165	220	1,640	4,100	2,290	7,650	2,610	5,690	537	755	755	350
26.....	187	204	1,070	4,100	1,990	7,650	2,710	3,610	478	555	650	860
27.....	175	102	870	4,100	1,040	5,150	2,820	2,710	446	582	665	322
28.....	170	192	1,000	3,150	1,380	3,610	2,500	2,000	464	2,930	555	301
29.....	176	336	755	2,500	4,620	2,100	1,720	600	2,710	510	280
30.....	157	573	550	2,190	16,400	1,900	1,460	528	1,550	470	268
31.....	145	660	1,720	10,900	1,400	1,000	480
1898-99.												
1.....	232	1,220	1,300	1,550	1,300	9,100	4,360	930	1,460	330	157	95
2.....	232	1,220	1,300	1,460	1,720	6,800	3,380	930	2,190	315	137	125
3.....	204	1,140	1,000	1,460	930	10,000	2,610	1,400	2,290	301	137	119
4.....	204	930	870	1,460	756	35,500	2,100	1,220	1,720	204	104	110
5.....	209	930	2,390	2,930	5,690	46,300	1,720	1,140	1,300	262	157	145
6.....	220	870	4,100	7,940	10,300	27,600	1,640	1,070	1,000	226	110	138
7.....	398	870	2,610	18,600	9,100	10,900	1,460	1,300	870	299	75	116
8.....	336	930	2,000	9,700	5,420	6,520	1,900	7,940	755	226	110	153
9.....	280	870	1,550	5,000	3,380	4,620	0,240	14,000	1,000	238	110	192
10.....	280	810	1,380	3,350	2,090	3,850	4,360	10,600	755	232	107	153
11.....	280	930	1,900	2,930	1,380	3,610	3,150	5,690	755	244	107	165
12.....	204	3,850	870	2,290	1,300	3,610	2,710	4,880	1,380	220	113	105
13.....	280	2,710	1,970	2,000	3,150	2,390	5,690	2,500	209	119	153
14.....	280	2,000	1,140	1,900	2,820	2,190	0,240	1,720	220	101	133
15.....	280	1,720	1,300	3,850	1,250	3,150	2,000	5,150	1,040	209	810	119
16.....	250	1,380	1,070	2,500	4,100	1,810	3,610	1,400	108	510	125
17.....	250	1,220	630	4,100	1,220	4,360	1,640	2,820	1,070	176	294	119
18.....	250	1,380	755	3,380	1,380	3,610	1,550	2,190	870	176	220	101
19.....	755	5,590	766	2,930	1,550	6,800	1,460	2,000	755	192	119	125
20.....	1,220	10,800	930	2,390	2,090	10,900	1,380	1,720	700	209	125	153
21.....	1,810	7,360	1,460	1,990	5,150	0,800	1,220	1,720	650	192	110	157
22.....	23,000	4,620	2,820	1,720	12,300	4,620	1,070	1,380	537	238	133	133
23.....	11,600	3,610	8,810	1,720	11,600	3,010	1,000	1,220	422	220	113	250
24.....	4,880	3,850	8,810	1,550	7,360	3,380	930	1,070	600	187	116	280
25.....	2,820	3,150	5,150	1,640	4,880	2,930	870	030	470	192	95	220
26.....	2,000	2,890	3,380	2,500	3,610	2,510	1,220	870	446	155	95	209
27.....	1,900	2,090	2,500	2,190	18,600	2,290	1,720	755	406	187	90	102
28.....	1,220	1,720	2,090	2,000	15,000	2,190	1,380	700	430	165	82	182
29.....	1,220	1,380	1,720	1,810	10,900	1,140	670	446	209	116	105
30.....	1,070	1,380	1,460	1,380	9,700	1,070	650	390	176	101	105
31.....	1,220	1,380	1,460	5,690	810	165	101

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the years ending Sept. 30, 1895-1917—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1899-1900												
1.....	157	110	170		050	4,100	3,850	810	755	1,400	1,000	192
2.....	133	157	165		390	12,300	2,930	755	060	1,000	050	182
8.....	133	209	145		470	7,080	2,500	680	600	1,000	510	157
4.....	133	200	145	390	470	4,100	2,200	650	700	980	430	145
5.....	125	220	145		470	3,150	2,000	600	630	582	350	125
6.....	125	268	145		600	3,150	1,900	555	555	510	280	110
7.....	125	232	146	390	930	6,420	1,720	528	404	430	133	110
8.....	125	200	133	300	1,220	6,520	1,550	510	430	404	220	05
0.....	104	182	125	300	3,010	4,020	1,900	470	430	454	204	00
10.....	110	157	125	555	4,020	3,380	1,000	600	414	330	192	88
11.....	110	157	125	555	2,820	2,820	1,640	610	360	301	176	82
12.....	107	157	157	670	2,090	2,100	1,300	470	315	250	157	78
13.....	107	145	700	4,620	13,000	1,900	1,140	480	268	238	146	70
14.....	110	137	930	2,190	16,000	1,720	1,070	440	280	220	125	70
16.....	113	137	1,000	1,650	6,800	1,810	870	300	555	102	125	08
10.....	110	137	700	1,140	4,100	1,040	810	374	310	176	126	414
17.....	110	125	510	030	2,030	1,640	700	350	2,200	145	110	260
18.....	86	125	315	030	2,000	1,220	930	329	7,360	145	116	166
10.....	85	110	301	1,300	1,300	1,720	1,070	301	6,150	145	126	146
20.....	06	110	322	0,700	1,070	10,800	2,710	350	2,930	133	110	125
21.....	95	133	390	8,520	1,140	17,100	3,150	1,720	1,000	133	126	119
22.....	88	119	315	4,880	4,100	8,230	3,150	1,070	1,300	157	125	137
23.....	75	110	301	2,820	7,040	5,420	3,380	810	1,000	600	106	105
24.....	101	110	336	2,000	5,160	2,710	2,820	700	755	700	126	238
25.....	00	125	801	1,460	3,610	6,240	2,100	700	050	755	145	105
20.....	82	125	030	1,140	2,930	5,000	1,720	1,070	578	1,000	125	182
27.....	82	113	528	1,000	2,200	0,240	1,400	1,000	566	4,100	106	145
28.....	82	126	350	755	1,640	4,360	700	700	470	3,850	105	125
20.....	82	182	350	582	4,100	1,000	700	470	1,720	102	146
30.....	75	198	350	1,000	3,610	930	930	510	1,070	220	125
31.....	101	350	655	4,360	030	1,070	105
1900-1.												
1.....	125	145	1,000	870	810	600	1,550	1,810	3,610	2,930	320	1,300
2.....	105	209	1,460	1,140	700	600	1,040	1,640	2,820	2,800	204	1,810
3.....	146	220	1,220	1,140	756	600	0,240	1,560	2,100	1,000	208	1,140
4.....	137	2,200	2,930	870	1,220	600	12,000	1,300	1,810	2,610	238	870
5.....	126	1,460	12,000	755	1,300	1,640	7,650	1,070	1,550	3,150	244	680
0.....	110	765	7,360	680	1,000	5,150	7,360	1,000	1,300	2,610	280	537
7.....	116	565	4,100	755	765	2,710	11,600	870	1,720	4,100	030	494
8.....	110	464	3,150	765	930	2,000	7,040	870	1,900	2,820	1,070	446
9.....	119	430	2,710	080	1,070	1,900	5,420	870	2,610	1,810	755	430
10.....	116	350	2,710	020	2,000	1,900	4,100	1,070	1,900	1,300	540	800
11.....	110	204	2,190	755	2,390	9,100	2,710	2,820	1,400	1,000	430	800
12.....	110	315	1,720	21,100	2,000	12,300	2,610	2,100	1,070	810	470	374
13.....	104	315	1,460	11,000	1,640	6,690	2,500	1,900	030	700	870	1,220
14.....	145	390	1,300	6,090	1,220	3,860	2,820	1,720	1,000	630	655	810
16.....	280	350	1,140	3,610	1,000	3,150	11,000	1,400	1,220	070	2,000	600
16.....	280	336	1,000	2,710	1,000	2,610	7,940	1,300	0,240	930	1,380	494
17.....	220	316	810	2,000	1,070	2,000	5,600	1,070	20,000	1,000	030	430
18.....	102	208	765	1,720	1,070	1,640	5,150	1,000	0,700	030	810	000
19.....	105	250	755	1,140	1,140	1,380	4,360	1,070	0,800	1,000	700	2,000
20.....	145	232	700	1,070	755	1,300	13,300	1,070	4,300	870	582	1,220
21.....	145	220	755	765	1,300	1,300	20,400	1,070	3,160	810	1,140	810
22.....	137	294	700	870	810	2,100	11,300	17,800	2,300	700	1,140	050
23.....	1,000	1,000	582	1,140	930	2,500	7,050	17,800	2,710	582	030	640
24.....	3,380	2,710	555	1,140	1,000	2,000	6,520	8,810	12,300	494	756	446
25.....	1,220	3,380	630	1,380	1,070	1,720	5,150	4,620	6,240	430	765	414
26.....	1,000	56,800	755	1,720	870	1,640	4,620	3,150	3,010	454	700	800
27.....	756	21,100	080	1,640	766	2,710	3,850	14,300	3,150	430	2,500	329
28.....	510	6,800	630	1,040	600	2,500	3,150	10,300	3,860	316	2,000	315
20.....	315	4,100	620	1,640	2,500	2,610	10,000	2,030	322	1,140	280
30.....	280	2,610	030	1,140	1,900	2,090	7,040	4,620	301	810	280
31.....	280	1,070	1,640	5,150	280	756

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1895-1917—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1001-2.												
1.....	315	145	238	6,240	7,360	30,700	5,150	700	310	870	105	113
2.....	301	119	238	4,100	6,240	12,900	3,010	700	680	810	165	104
3.....	244	145	256	3,150	6,800	8,520	2,930	756	000	870	280	101
4.....	238	145	650	2,500	4,100	5,420	2,500	766	555	930	220	110
5.....	232	105	2,610	1,000	3,850	3,850	4,100	766	610	810	102	104
6.....	220	165	1,640	1,220	1,900	2,820	3,850	755	440	510	182	101
7.....	220	149	1,070	1,070	1,900	2,820	5,160	756	430	510	166	110
8.....	220	145	810	1,140	1,720	2,820	7,940	755	390	556	165	101
9.....	220	153	700	1,300	1,720	4,300	7,040	700	350	600	165	00
10.....	200	140	700	1,140	2,190	9,100	6,240	650	330	660	105	82
11.....	192	145	1,220	1,070	810	7,040	6,800	600	315	510	105	75
12.....	182	145	2,290	870	870	0,700	5,900	555	301	464	165	75
13.....	192	220	1,550	755	810	15,000	5,420	610	316	306	149	75
14.....	200	176	1,220	650	370	15,700	2,190	494	336	350	133	82
15.....	220	166	30,700	510	810	7,360	3,160	404	350	350	125	82
16.....	220	165	14,300	650	650	7,940	2,710	430	374	320	125	90
17.....	232	165	6,240	620	700	16,000	2,500	430	382	315	125	82
18.....	232	105	3,380	600	650	0,700	2,000	446	300	315	125	82
19.....	238	165	2,390	510	650	5,600	1,810	300	308	308	122	82
20.....	244	165	1,810	470	600	3,850	1,720	390	480	308	110	70
21.....	250	157	1,380	682	650	2,930	1,040	573	470	280	110	70
22.....	250	153	810	510	650	2,390	1,460	555	510	268	113	70
23.....	250	145	810	670	1,000	1,000	1,380	810	600	220	110	70
24.....	266	166	810	555	2,000	1,810	1,300	650	060	192	110	70
25.....	256	220	870	470	15,700	1,310	1,220	030	660	165	05	75
26.....	262	350	1,000	600	17,800	2,000	1,070	1,640	682	105	06	70
27.....	238	422	1,000	10,300	10,900	1,810	1,000	2,930	4,100	105	88	82
28.....	209	350	1,000	12,900	29,600	1,640	810	2,090	1,640	165	82	164
29.....	182	250	23,000	6,240	10,000	810	1,550	1,550	165	06	120
30.....	163	202	28,000	0,520	15,700	755	1,220	030	165	82	140
31.....	105	11,000	11,600	9,400	1,000	165	104
1902-3.												
1.....	140	86	1,160	576	6,520	23,800	2,980	2,620	866	2,500	280	207
2.....	101	80	4,100	708	4,300	9,400	2,330	2,040	1,020	1,520	544	164
3.....	246	80	3,220	23,100	5,420	5,420	2,040	1,620	1,100	1,080	788	154
4.....	315	101	4,620	13,600	14,300	3,850	2,040	1,330	855	814	635	154
5.....	240	80	2,800	8,520	10,000	2,860	2,500	1,240	600	508	866	154
6.....	140	79	2,500	5,150	7,650	2,260	2,500	1,080	608	855	301	132
7.....	120	63	2,040	3,000	5,160	1,920	2,140	928	2,380	1,080	260	113
8.....	80	63	1,820	2,860	3,220	2,020	3,600	928	8,810	1,330	220	05
9.....	76	51	1,420	2,040	2,860	7,080	8,520	855	4,620	028	200	206
10.....	70	53	1,330	1,720	2,260	7,040	6,800	788	2,860	606	188	132
11.....	76	68	1,240	928	1,720	5,420	4,620	720	2,740	490	154	113
12.....	80	58	788	1,420	4,360	6,240	3,100	659	4,100	443	58	113
13.....	164	58	1,920	1,620	6,240	5,090	2,620	598	2,800	1,240	576	95
14.....	415	58	7,300	1,720	4,360	4,620	2,500	544	1,020	2,260	424	96
15.....	306	53	6,090	928	3,000	3,000	2,860	544	1,520	1,020	267	79
16.....	280	63	6,420	788	20,700	2,740	2,980	490	1,240	1,160	154	95
17.....	205	53	0,800	1,080	24,900	2,140	2,800	462	1,160	788	154	145
18.....	188	56	7,080	1,080	0,400	1,720	2,380	443	865	598	120	207
19.....	188	56	4,880	1,160	5,690	1,520	2,380	372	720	400	113	301
20.....	104	58	2,860	834	3,850	1,240	2,040	390	598	300	154	287
21.....	120	68	2,380	855	2,800	1,160	2,040	356	612	356	132	207
22.....	80	53	1,720	1,100	2,620	2,620	2,380	350	415	188	120	246
23.....	86	86	2,500	1,330	2,140	28,000	2,260	331	306	280	132	217
24.....	36	101	2,140	1,240	2,140	31,100	2,040	315	396	240	132	188
25.....	80	117	1,720	1,080	2,140	10,600	1,920	331	396	217	120	140
26.....	80	2,260	1,330	1,000	2,040	6,000	3,850	390	544	188	120	132
27.....	86	3,100	071	971	1,020	4,100	10,000	396	471	160	65	120
28.....	86	2,740	598	2,620	20,700	2,980	6,520	2,380	059	104	58	120
29.....	86	2,620	598	11,300	2,380	4,300	1,520	2,260	128	58	120
30.....	80	1,240	788	12,000	1,320	3,220	1,080	4,100	164	234	120

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the years ending Sept. 30, 1895-1917—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1003-4.												
1.....	120	140	101	234	659	5,090	2,380	3,860	1,720	1,160	164	80
2.....	140	120	120	240	747	8,620	4,620	3,100	3,100	866	140	101
3.....	120	120	140	566	400	5,900	4,360	2,620	6,150	069	101	80
4.....	120	120	132	1,080	301	6,150	3,360	3,100	7,300	644	101	80
5.....	101	120	120	1,020	306	2,740	2,500	3,000	3,600	644	140	86
6.....	101	140	132	1,000	246	3,220	1,920	1,020	2,040	644	164	86
7.....	120	140	120	622	490	2,620	1,720	1,720	1,720	400	188	86
8.....	120	140	101	490	7,300	10,000	1,620	1,520	1,420	400	188	86
9.....	120	140	120	612	5,420	7,650	1,620	1,330	1,240	490	140	86
10.....	120	120	120	400	3,100	5,150	1,720	1,240	1,080	443	140	86
11.....	120	120	132	306	2,140	3,350	1,720	1,060	1,000	396	140	86
12.....	140	120	140	339	1,330	2,300	1,720	1,080	856	490	164	86
13.....	246	120	101	315	1,080	2,860	1,020	028	788	315	120	101
14.....	183	104	132	204	788	2,380	1,020	788	659	315	104	101
15.....	188	120	132	246	622	2,260	1,520	788	544	280	140	86
16.....	188	120	101	443	855	2,620	1,330	856	400	240	120	70
17.....	140	140	132	416	1,620	1,020	1,330	928	544	217	120	70
18.....	164	104	140	380	306	1,720	1,240	1,080	490	188	140	86
19.....	104	188	120	200	316	1,520	1,080	15,700	443	188	140	70
20.....	104	188	101	217	720	1,520	1,080	9,100	443	188	140	120
21.....	188	246	140	246	788	1,420	1,000	6,240	390	188	164	188
22.....	188	217	188	490	7,080	1,420	928	5,150	306	140	188	70
23.....	104	217	234	16,400	0,400	3,000	788	3,220	644	140	140	70
24.....	164	188	234	7,040	5,690	7,660	720	2,740	443	104	188	86
25.....	164	188	229	4,100	5,150	5,160	659	2,740	490	140	188	70
26.....	104	178	316	2,260	3,000	3,000	788	2,600	300	188	140	70
27.....	104	164	306	1,820	2,200	2,080	2,740	2,140	400	188	120	70
28.....	140	164	372	1,020	1,620	2,740	8,810	1,820	720	217	101	101
29.....	140	120	315	2,040	2,980	2,140	6,800	1,420	855	188	101	70
30.....	120	70	316	1,520	1,620	6,150	1,240	1,420	104	86	40
31.....	140	490	1,080	1,420	1,330	104	86
1004-6.												
1.....	46	101	101	490	720	1,000	1,020	1,720	1,000	644	660	315
2.....	46	80	101	644	720	2,140	1,620	1,520	856	1,720	644	306
3.....	40	58	101	059	720	2,600	1,520	1,330	720	1,420	400	400
4.....	40	70	101	490	720	2,620	1,330	1,100	508	1,080	300	028
5.....	46	80	120	460	608	4,620	1,160	1,000	400	2,380	400	720
6.....	40	70	140	430	490	4,620	2,380	865	490	1,720	598	544
7.....	58	46	240	306	390	5,150	3,600	1,520	300	1,330	720	443
8.....	70	70	280	370	246	7,940	3,220	2,140	396	1,520	720	315
9.....	101	68	217	340	315	1,000	2,860	2,040	315	1,080	644	280
10.....	86	46	164	315	315	25,600	2,380	1,520	315	855	443	246
11.....	101	40	140	443	240	18,600	2,140	1,420	315	720	855	246
12.....	70	70	120	1,240	315	10,900	2,140	28,400	316	608	1,000	246
13.....	70	70	120	6,420	490	8,230	1,020	21,100	316	8,230	866	300
14.....	70	70	140	5,690	659	6,240	1,720	10,600	280	12,900	720	315
15.....	68	101	104	3,350	856	6,240	1,520	7,940	280	8,230	788	315
16.....	68	101	164	2,380	1,000	5,960	1,330	7,080	356	4,360	928	280
17.....	46	101	101	1,020	788	6,090	1,240	6,240	315	2,380	1,080	246
18.....	70	101	101	1,330	598	6,800	1,080	5,150	246	1,720	1,000	217
19.....	70	101	101	855	608	0,800	1,000	3,860	240	1,240	865	188
20.....	70	70	140	856	490	7,940	855	2,800	246	928	059	188
21.....	70	80	120	855	396	10,300	855	2,040	246	856	544	164
22.....	70	101	120	856	396	15,000	028	1,620	1,330	650	490	140
23.....	58	101	70	788	855	0,100	1,080	1,420	1,920	3,600	390	140
24.....	46	101	101	720	856	5,690	1,000	1,160	2,040	6,800	856	140
25.....	70	101	164	720	1,160	4,360	855	1,000	5,060	3,600	316	140
26.....	58	101	720	720	1,520	4,100	855	855	8,600	2,200	396	101
27.....	46	101	2,380	720	1,620	3,600	855	855	2,860	1,520	315	140
28.....	40	101	2,140	720	1,420	3,100	1,000	788	2,040	1,080	698	120
29.....	40	101	1,920	720	2,620	2,140	720	1,330	1,000	400	101
30.....	46	86	1,420	720	2,380	1,920	855	855	788	443	101

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1895-1917—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1905-6.												
1.....	101	396	598	1,520	2,140	598	7,080	1,520	1,000
2.....	70	230	720	1,330	1,720	544	6,240	1,330	1,000
3.....	140	280	4,620	5,150	1,330	544	4,020	1,240	928
4.....	101	246	11,300	14,600	1,080	4,620	3,350	1,720	928
5.....	101	246	5,420	11,300	028	0,240	2,860	2,500	028
6.....	101	246	3,100	0,240	1,000	3,000	2,860	2,380	928
7.....	101	246	2,040	3,850	028	2,380	2,140	2,140	028
8.....	101	246	1,020	2,860	788	1,920	4,100	2,140	1,720
9.....	101	315	1,240	2,380	720	1,820	3,600	2,140	1,720
10.....	101	356	1,080	1,820	788	1,720	1,920	1,020	1,000
11.....	140	315	855	1,080	720	1,000	5,600	1,720	028
12.....	188	280	788	1,720	659	1,000	4,360	1,520	028
13.....	315	246	720	1,520	598	028	3,350	1,160	855
14.....	240	246	508	4,360	598	1,000	2,020	1,000	855
15.....	356	188	720	4,360	508	6,520	7,050	855	855
16.....	246	188	598	4,100	508	13,300	7,040	1,000	855
17.....	188	188	544	3,000	396	8,230	6,520	1,160	855
18.....	188	188	544	3,100	315	4,620	3,350	1,380	928
19.....	164	217	490	2,740	315	3,350	2,080	1,330	1,000
20.....	140	246	720	2,380	396	4,360	2,620	1,240	1,240
21.....	140	390	1,000	2,040	443	4,620	2,140	1,160	2,800
22.....	140	598	0,700	1,720	544	4,100	1,020	1,160	2,260
23.....	140	544	6,240	21,100	1,420	4,300	1,820	1,080	1,720
24.....	140	490	4,620	19,300	1,160	4,100	1,520	1,080	1,240
25.....	164	443	3,600	8,230	1,000	4,100	1,330	1,000	855
26.....	240	300	2,980	5,420	855	5,420	1,240	855	855
27.....	720	396	2,140	3,000	788	7,080	1,240	855	508
28.....	1,620	350	1,720	3,850	720	12,900	3,000	855	490
29.....	788	356	1,520	3,850	11,600	2,260	928	490
30.....	598	400	1,520	3,100	9,700	1,820	028	396
31.....	598	1,520	2,620	10,000	1,000
1007.												
1.....	7,360	1,000	855	400
2.....	8,230	855	508	300
3.....	6,800	788	490	300
4.....	4,360	720	396	400
5.....	3,350	855	356	400
6.....	2,860	720	356	443
7.....	2,740	598	315	443
8.....	2,260	544	788	390
9.....	14,300	598	598	396
10.....	5,060	0,800	544	490	315
11.....	4,020	6,240	2,620	396	598
12.....	3,600	7,650	1,720	598	1,920
13.....	2,860	11,300	1,420	598	2,020
14.....	2,260	41,200	1,920	400	1,420
15.....	1,920	14,300	1,420	356	1,000
16.....	1,720	7,300	1,100	315	720
17.....	1,520	5,690	1,080	246	508
18.....	1,330	3,350	6,240	280	1,000
19.....	1,160	2,500	6,240	246	855
20.....	1,000	1,920	3,100	240	855
21.....	1,000	1,720	2,140	217	720
22.....	855	1,330	1,330	164	508
23.....	788	1,330	1,080	246	659
24.....	720	4,100	855	315	2,620
25.....	720	2,860	855	4,880	3,350
26.....	720	2,860	659	3,600	1,720
27.....	855	2,380	544	2,140	1,100
28.....	855	1,720	490	1,330	855
29.....	855	1,330	490	1,000	720
30.....	788	1,160	598	720	650
31.....	928	1,160	598

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1895-1917—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1007-8.												
1.....	720	300	1,520	7,660	1,330	1,520	26,000	2,380	3,100	306	355	443
2.....	608	670	1,330	6,420	1,330	5,000	15,700	3,100	2,140	315	659	390
3.....	490	0,800	1,240	5,150	1,720	11,600	10,300	2,740	1,520	400	544	315
4.....	443	4,880	1,080	3,100	2,860	9,700	0,800	2,860	1,520	1,620	443	280
6.....	390	2,860	855	2,860	1,330	8,520	4,880	2,020	5,160	4,360	306	246
6.....	490	2,140	720	2,020	1,000	16,000	3,600	2,620	4,100	2,860	306	240
7.....	865	1,720	720	2,200	1,080	25,300	3,100	19,300	2,020	3,100	246	240
8.....	855	1,020	698	1,720	1,160	22,200	2,620	21,500	1,920	2,380	396	217
9.....	7,940	1,920	855	1,020	1,160	12,300	2,380	9,100	1,620	1,420	1,000	217
10.....	4,100	4,100	2,860	1,620	1,160	10,800	3,100	6,520	1,330	1,000	1,520	188
11.....	2,330	0,400	15,700	1,420	1,000	6,800	5,150	10,300	1,100	788	1,620	188
12.....	1,720	6,060	8,520	21,800	1,520	5,150	7,940	0,240	1,000	598	1,000	164
13.....	1,100	3,860	4,880	21,500	4,100	4,620	5,690	4,360	855	490	720	164
14.....	1,000	2,740	3,600	10,900	13,000	4,020	4,360	3,350	720	443	644	104
15.....	788	1,920	4,100	6,240	35,500	4,620	3,350	2,620	855	396	490	104
16.....	720	1,520	3,850	4,620	39,600	4,620	4,100	2,140	855	306	306	140
17.....	508	1,330	3,100	3,850	10,000	5,150	6,240	1,720	856	300	396	140
18.....	490	1,100	2,380	2,800	0,800	8,620	5,150	1,720	720	316	356	140
19.....	400	2,200	2,140	2,620	4,620	8,520	4,880	2,140	598	315	316	140
20.....	396	4,100	1,720	2,140	4,100	10,900	6,150	3,850	400	315	280	140
21.....	306	3,600	1,520	1,920	2,860	7,300	4,100	6,600	598	188	246	120
22.....	306	2,800	1,330	2,140	2,380	5,420	3,600	4,880	720	306	240	140
23.....	315	3,100	2,200	2,800	2,140	4,300	2,020	8,230	400	306	240	140
24.....	315	7,300	10,600	3,350	1,020	4,620	2,140	4,880	490	544	217	120
26.....	315	8,810	10,600	2,380	1,720	4,100	2,040	4,360	1,000	508	246	120
26.....	315	5,000	6,240	1,520	1,720	3,350	4,100	5,090	720	1,240	1,330	120
27.....	316	4,100	4,360	3,100	1,720	2,800	5,420	3,850	598	3,600	2,140	101
28.....	280	2,860	3,350	4,100	1,520	2,140	3,850	2,860	598	4,620	1,620	101
29.....	315	2,380	3,350	3,000	1,420	2,500	2,860	2,620	490	3,600	1,000	101
30.....	315	1,020	7,300	3,100	3,100	2,260	4,100	396	1,720	720	101
31.....	350	10,300	2,620	13,300	5,420	1,160	544
1008-9.												
1.....	140	608	240	6,160	1,160	3,100	2,860	6,800	1,420	1,000	2,380	140
2.....	140	443	217	3,100	788	2,620	2,380	6,800	1,240	1,330	1,160	101
3.....	140	315	188	2,140	788	2,260	2,140	5,150	1,100	3,100	788	101
4.....	140	280	188	1,720	1,160	4,880	2,140	3,850	1,100	1,520	508	188
5.....	120	217	188	2,860	1,160	4,620	2,040	3,100	1,520	1,000	490	164
6.....	140	188	164	11,300	1,160	3,600	3,100	2,590	1,720	855	396	140
7.....	140	188	240	7,360	1,160	3,000	3,100	1,020	1,330	3,000	396	140
8.....	140	188	598	4,620	1,100	6,240	2,620	1,820	1,080	2,140	315	188
9.....	140	188	059	2,860	1,000	7,080	2,140	1,720	1,240	1,330	315	188
10.....	140	140	490	2,140	6,800	6,800	1,720	2,500	1,520	855	188	188
11.....	104	188	400	1,720	12,300	8,230	1,520	5,090	2,140	720	246	217
12.....	140	164	1,920	1,620	7,080	5,690	1,330	4,620	1,520	598	246	188
13.....	140	104	2,020	1,330	4,360	4,100	1,160	3,220	1,100	400	217	188
14.....	140	164	1,720	1,240	4,300	4,100	9,700	2,380	1,000	396	188	188
15.....	140	188	1,100	1,520	5,150	3,600	15,900	1,920	855	396	246	188
16.....	140	140	855	8,520	5,600	2,860	7,080	1,720	720	396	390	246
17.....	140	140	720	10,300	7,940	2,380	4,620	1,420	720	356	598	544
18.....	140	140	720	6,520	5,690	2,040	3,350	1,160	1,000	315	720	315
19.....	101	217	1,000	4,360	4,100	1,720	2,620	1,080	1,160	315	544	246
20.....	140	315	1,520	3,600	3,600	1,520	2,140	928	1,330	246	396	240
21.....	120	306	1,330	3,600	4,100	1,520	1,920	855	1,000	246	356	188
22.....	120	443	1,000	4,100	4,620	1,720	2,140	5,150	855	217	246	188
23.....	120	396	855	4,100	4,880	2,040	4,880	5,420	659	246	246	164
24.....	140	390	720	4,620	4,880	2,860	8,520	3,350	659	246	188	188
25.....	140	315	720	4,880	6,240	3,850	6,520	2,380	788	188	217	164
26.....	188	315	1,720	4,100	5,960	10,000	4,620	4,100	720	188	188	164
27.....	217	280	1,520	3,100	4,620	6,520	3,600	7,080	598	188	188	188
28.....	246	240	1,330	2,380	3,850	5,150	2,860	5,420	490	188	188	164
29.....	240	246	1,330	1,920	3,350	2,860	3,600	720	188	164	140
30.....	659	240	1,920	1,920	4,620	3,100	2,380	1,000	246	140	140
31.....	720	5,150	1,520	3,850	1,720	246	140

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the years ending Sept. 30, 1895-1917—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1009-10.												
1.....	140	315	315	490	1,000	5,420	550	2,380	855	884	315	246
2.....	140	280	280	400	855	11,300	598	1,820	828	747	301	246
3.....	140	246	246	738	1,000	0,240	2,400	1,480	788	696	280	544
4.....	120	246	240	5,560	1,330	5,690	884	1,300	747	650	246	1,240
5.....	101	217	246	4,100	1,720	4,300	1,240	1,100	028	788	331	1,820
6.....	101	188	246	2,620	1,520	3,600	1,150	971	3,860	884	380	1,720
7.....	101	188	246	5,150	1,150	3,350	1,160	855	6,820	1,080	315	1,330
8.....	101	188	315	8,380	855	3,100	071	028	3,480	1,030	301	028
9.....	86	217	396	4,620	855	2,020	884	1,190	2,380	1,240	246	650
10.....	80	246	490	3,350	1,150	2,140	828	1,240	2,420	1,100	246	512
11.....	316	720	246	2,860	1,160	1,920	720	1,100	6,520	928	217	443
12.....	855	1,720	390	1,330	720	1,720	788	1,160	8,950	747	200	380
13.....	1,520	1,240	306	1,150	855	1,720	1,880	1,620	10,900	1,620	217	396
14.....	1,080	855	0,700	1,330	865	1,020	2,140	2,200	13,900	2,620	178	612
15.....	060	720	6,240	1,160	1,000	2,100	1,970	1,920	8,810	2,100	240	490
16.....	400	598	3,600	1,000	1,330	1,020	1,020	1,620	14,000	1,920	280	380
17.....	396	400	2,250	855	4,100	1,680	1,520	1,300	34,500	1,500	246	315
18.....	306	443	1,620	865	14,300	1,420	1,880	1,240	11,600	2,000	178	250
19.....	300	306	1,100	3,100	10,000	1,300	2,140	1,100	6,240	4,880	188	246
20.....	315	300	1,520	6,420	6,420	1,130	2,200	1,100	7,710	2,570	188	200
21.....	246	306	1,000	4,100	3,850	1,000	2,380	1,100	5,150	1,620	178	188
22.....	246	316	1,000	14,500	3,850	1,000	2,380	1,020	3,980	1,100	217	188
23.....	217	316	720	7,080	5,000	028	2,200	1,540	3,000	028	178	178
24.....	598	315	698	4,360	4,020	028	2,500	1,480	4,400	788	606	178
25.....	608	306	544	3,100	3,600	865	3,480	1,480	2,860	593	443	154
26.....	490	390	400	2,140	2,800	855	3,150	1,480	1,920	400	390	260
27.....	698	390	490	1,020	2,850	855	2,860	1,330	1,420	490	301	1,420
28.....	490	350	443	1,720	3,100	855	3,220	1,190	1,330	490	240	788
29.....	396	316	396	1,720	600	3,220	071	1,100	415	200	1,330
30.....	396	315	390	1,330	720	2,810	884	1,030	390	178	747
31.....	316	598	1,160	720	856	380	188
1010-11.												
1.....	612	250	720	6,000	0,100	2,140	3,850	1,720	828	250	101	3,220
2.....	380	200	670	11,600	7,080	1,020	3,480	1,720	696	234	86	2,140
3.....	366	246	443	23,000	5,420	1,720	4,100	1,480	598	380	86	1,030
4.....	315	240	306	23,100	4,620	1,480	7,300	1,300	512	380	95	747
5.....	240	200	443	10,300	3,980	1,240	17,800	1,130	612	301	380	622
6.....	200	188	443	4,020	3,220	6,600	16,400	1,000	855	280	154	512
7.....	240	188	300	3,350	2,360	8,620	9,100	855	622	240	140	380
8.....	240	188	331	2,380	2,020	6,590	6,800	855	659	415	188	315
9.....	200	188	315	1,820	4,100	4,360	0,100	855	071	315	331	246
10.....	022	188	415	1,420	5,600	6,800	8,230	855	028	280	315	028
11.....	622	178	306	1,160	4,360	9,400	5,500	828	650	471	188	828
12.....	471	160	356	1,370	3,220	7,350	4,360	788	544	306	188	720
13.....	356	104	350	1,620	2,740	5,150	3,600	720	490	315	415	1,560
14.....	301	154	380	8,520	2,140	5,500	3,350	022	443	301	443	928
15.....	234	164	200	5,960	1,920	5,150	9,700	675	576	260	300	659
16.....	217	164	315	4,020	1,620	4,100	8,520	544	471	234	356	855
17.....	217	104	315	3,600	1,370	3,600	5,900	512	443	234	280	3,850
18.....	188	188	260	2,740	1,300	3,100	4,360	380	396	188	217	2,620
19.....	188	188	356	2,140	1,160	2,080	3,480	471	396	164	188	1,520
20.....	200	164	315	1,720	1,580	6,240	4,880	471	315	150	140	1,030
21.....	178	164	443	1,020	1,720	0,800	5,600	396	356	140	140	788
22.....	178	150	490	3,100	1,420	4,880	5,420	396	356	140	120	622
23.....	188	150	306	8,520	1,240	3,720	4,300	443	490	140	101	576
24.....	246	178	544	5,960	1,130	3,220	3,850	400	396	150	109	400
25.....	234	150	1,370	3,980	1,240	2,620	3,220	396	315	140	109	443
26.....	245	104	1,420	3,220	1,330	2,140	2,860	331	315	109	109	331
27.....	260	331	1,240	3,850	1,520	2,740	2,380	350	356	101	132	280
28.....	246	300	1,030	5,690	1,720	5,150	2,040	315	443	101	101	315
29.....	217	508	1,160	6,800	4,100	1,720	512	396	120	120	315
30.....	246	788	6,940	35,100	3,850	1,720	855	316	120	132	380
31.....	246	8,660	19,300	4,100	1,000	101	217

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the years ending Sept. 30, 1895-1917—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1011-12.												
1.....	315	696	1,480	10,000	3,720	4,100	5,960	2,740	501	1,330	400	183
2.....	698	698	1,330	0,520	2,980	2,860	7,940	2,260	481	942	396	178
3.....	598	544	1,230	4,620	2,380	2,600	9,700	1,920	443	696	347	154
4.....	1,130	490	1,160	3,480	1,660	2,040	6,800	1,600	415	587	416	145
5.....	1,160	415	028	2,620	1,480	1,720	4,880	1,300	380	1,060	380	120
6.....	928	720	842	1,620	1,190	1,560	3,000	1,300	364	1,720	301	117
7.....	747	10,300	801	1,100	1,330	1,400	3,100	1,030	331	1,050	287	117
8.....	508	6,800	774		1,080	1,640	2,800	2,980	287	913	253	128
9.....	3,850	3,720	774		1,300	3,220	2,380	2,620	240	683	229	109
10.....	2,380	2,740	720		928	3,980	2,140	2,260	246	566	211	101
11.....	1,680	2,040	720		942	3,100	1,920	1,920	188	587	188	109
12.....	1,680	1,680	720		788	2,800	1,640	10,600	188	842	205	95
13.....	1,680	2,140	708	700	801	6,520	1,480	17,600	205	708	200	70
14.....	1,420	2,140	720		720	7,940	1,310	9,700	188	633	217	73
15.....	1,190	1,920	734		683	10,900	1,230	5,960	211	671	188	76
16.....	4,100	1,700	814		059	31,100	1,140	10,000	194	696	160	73
17.....	3,720	1,480	1,190		671	11,000	1,230	18,200	194	471	140	08
18.....	18,200	1,480	1,820		720	6,800	2,140	8,230	217	443	160	79
19.....	10,300	3,600	1,720	1,330	950	6,150	3,480	4,880	331	471	174	200
20.....	6,150	3,850	1,520	2,380	1,370	4,620	2,980	3,600	416	396	164	164
21.....	2,860	2,620	1,310	2,140	3,360	6,690	2,260	2,620	405	000	140	109
22.....	1,920	2,140	1,300	2,040	16,800	6,960	1,920	2,040	350	644	104	95
23.....	2,380	1,620	6,420	1,720	9,100	5,150	1,820	1,620	490	481	169	178
24.....	3,100	1,500	4,620	1,720	6,150	7,940	1,920	1,370	300	1,700	140	2,980
25.....	2,140	1,620	6,690	2,040	3,600	12,600	1,820	1,180	655	655	136	2,200
26.....	1,720	1,680	4,880	1,920	3,350	8,310	1,680	1,020	644	6,800	146	1,400
27.....	1,420	1,620	5,690	1,820	17,800	5,960	1,820	913	544	2,620	140	942
28.....	1,190	1,480	6,800	1,620	10,300	4,300	2,600	788	2,040	1,420	117	508
29.....	971	1,440	4,620	2,140	6,900	17,500	2,380	720	1,920	913	105	471
30.....	828	1,620	3,480	6,420	16,800	2,740	606	1,680	671	113	405
31.....	720	3,720	6,160	8,810	687	533	130
1912-13.												
1.....	347	223	183	9,700	3,360	4,880	3,360	1,190	5,160	533	416	246
2.....	301	217	217	4,620	3,480	3,850	2,620	1,030	3,480	022	350	246
3.....	246	200	246	2,740	3,220	2,860	2,140	071	2,020	434	512	178
4.....	240	194	360	4,020	7,360	2,140	1,820	913	5,090	1,000	304	183
5.....	205	188	490	4,100	7,080	1,920	1,620	855	5,150	1,420	301	150
6.....	183	188	1,100	3,480	4,020	1,720	1,400	801	3,100	1,230	253	178
7.....	188	315	2,620	7,360	3,220	1,400	1,100	1,100	2,140	1,330	223	140
8.....	178	5,690	2,140	14,300	2,140	1,130	1,060	1,180	3,100	986	188	160
9.....	160	3,980	1,520	13,600	1,080	870	986	1,020	2,980	696	178	140
10.....	104	2,140	1,130	10,300	1,540	1,030	913	884	2,260	598	104	140
11.....	169	1,440	384	4,620	1,540	3,100	913	774	1,700	8,230	217	124
12.....	150	1,060	747	3,000	1,820	5,690	2,600	671	1,300	3,360	260	124
13.....	146	828	610	3,850	1,460	4,100	19,300	610	1,140	2,140	308	140
14.....	154	606	415	3,850	1,160	10,000	11,900	698	971	1,420	2,860	113
15.....	200	598	331	2,980	1,030	17,100	17,100	676	828	1,080	2,740	117
16.....	188	533	452	2,380	1,000	12,300	12,900	544	696	928	1,200	136
17.....	178	481	555	1,920	1,130	7,080	7,940	622	622	834	788	120
18.....	150	415	501	1,820	071	4,360	5,150	747	555	734	555	136
19.....	183	388	471	1,600	842	3,220	3,850	1,260	490	610	676	146
20.....	234	356	452	1,480	774	2,620	3,100	1,370	443	855	490	132
21.....	434	331	315	1,350	801	2,140	2,500	3,350	396	1,130	481	183
22.....	331	308	234	1,260	884	1,920	1,920	2,260	380	783	471	294
23.....	331	280	178	1,240	1,080	1,620	1,700	2,250	366	622	576	331
24.....	388	274	217	1,370	1,400	1,330	1,500	6,800	364	587	1,400	443
25.....	490	260	280	2,860	1,310	1,230	1,330	7,360	364	512	1,240	347
26.....	443	267	253	3,600	1,140	1,500	1,160	4,360	659	747	788	260
27.....	405	246	280	3,600	1,370	42,500	1,230	4,880	013	622	576	217
28.....	331	246	452	4,620	3,980	34,700	1,330	15,700	828	490	452	188
29.....	301	234	347	3,850	10,300	1,370	9,400	622	434	388	183
30.....	287	194	1,480	3,100	6,240	1,310	5,420	512	380	323	164
31.....	246	12,900	2,740	4,360	6,240	396	280

Daily discharge, in second-feet, of Greenbrier River at Alderson, W. Va., for the years
ending Sept. 30, 1895-1917—Continued.

Day.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.
1913-14.												
1.....	183	828	2,620	1,820	12,300	2,860	4,880	2,380	237	280	200	512
2.....	164	690	6,520	1,620	0,800	2,500	5,600	2,040	294	240	200	406
3.....	178	698	6,150	1,620	4,100	1,920	8,520	1,720	274	234	200	315
4.....	109	533	3,480	1,440	3,220	1,720	6,240	1,500	246	223	188	287
5.....	211	481	2,500	1,370	2,620	2,140	4,360	1,480	301	188	104	240
6.....	104	434	2,040	1,300	2,020	2,140	3,350	2,040	287	183	140	223
7.....	287	380	2,040	1,230	6,300	1,920	2,740	3,220	260	164	150	200
8.....	246	415	3,860	1,230	8,520	1,720	2,380	2,860	246	188	200	211
9.....	234	884	3,480	1,920	5,420	1,680	0,520	2,620	280	234	194	188
10.....	217	3,980	2,620	4,100	3,850	1,480	6,690	2,140	287	246	260	169
11.....	331	3,360	2,260	4,100	3,100	2,380	3,850	1,820	240	223	240	217
12.....	071	2,380	1,820	3,480	2,380	1,920	3,100	1,560	217	178	331	350
13.....	676	2,140	1,480	2,860	1,020	3,480	2,500	1,370	200	200	380	471
14.....	490	3,480	1,230	2,380	1,240	2,860	1,920	1,210	183	1,000	347	452
15.....	390	0,700	1,160	2,140	1,190	3,350	2,140	1,080	194	720	315	380
16.....	304	11,000	1,030	1,820	1,230	6,240	0,800	971	169	544	331	260
17.....	323	14,300	942	1,480	1,130	11,300	7,030	842	164	633	301	200
18.....	308	8,230	842	1,300	1,190	13,300	6,150	774	188	610	240	160
19.....	280	4,880	814	1,300	3,100	0,100	3,030	720	160	443	211	06
20.....	364	3,360	747	1,240	14,000	6,960	4,380	622	169	347	194	101
21.....	1,720	2,600	708	6,090	8,810	4,100	7,360	587	160	301	169	101
22.....	2,140	1,820	600	0,700	5,420	3,480	6,420	544	164	301	160	101
23.....	1,480	1,480	720	5,160	5,420	3,220	4,100	512	140	200	140	106
24.....	1,240	1,190	708	3,850	4,880	3,220	3,220	400	140	211	124	101
25.....	5,160	071	801	0,100	3,600	3,350	2,020	452	160	183	132	101
26.....	7,030	855	2,860	8,230	3,100	5,600	2,740	424	183	183	164	120
27.....	4,880	760	0,240	6,420	2,080	10,900	12,600	406	234	211	274	124
28.....	2,860	828	3,720	3,980	2,860	9,400	6,240	304	659	178	396	100
29.....	1,920	2,040	2,860	3,350	9,100	4,360	304	415	100	1,130	113
30.....	1,390	2,380	2,020	3,480	6,800	3,100	323	315	211	660	100
31.....	1,030	1,020	4,100	5,420	308	263	659
1914-16.												
1.....	64	218	160	4,490	4,490	2,500	804	1,250	204	185	137	463
2.....	66	185	547	3,710	27,800	2,090	777	1,080	516	169	133	394
3.....	66	160	1,250	2,810	24,800	1,700	726	1,000	626	186	126	346
4.....	69	168	1,420	1,510	15,200	1,420	686	1,100	7,500	204	530	304
5.....	72	164	1,990	1,080	6,120	1,420	649	1,160	5,300	268	910	726
6.....	90	153	5,300	1,510	5,850	1,340	637	1,000	2,810	330	637	1,000
7.....	92	142	3,030	26,300	7,600	1,420	592	832	1,990	336	586	818
8.....	92	126	2,100	16,700	0,680	1,510	680	777	1,420	355	394	637
9.....	97	133	1,790	11,600	5,580	1,340	930	699	1,100	374	811	526
10.....	92	115	1,700	9,800	3,960	1,340	1,080	640	902	302	302	432
11.....	100	118	1,700	7,500	3,250	1,600	1,000	681	712	239	253	336
12.....	118	111	1,890	6,400	2,600	1,790	1,100	526	502	211	246	277
13.....	111	115	1,890	6,400	2,190	1,790	1,080	626	620	197	260	260
14.....	122	115	1,800	6,580	1,600	1,610	1,250	484	615	190	364	263
15.....	239	122	1,890	3,480	3,480	1,420	1,080	484	1,600	185	790	294
16.....	403	122	1,800	4,226	7,780	1,420	930	403	1,700	197	604	232
17.....	1,080	218	1,890	6,300	5,300	1,600	874	442	1,510	153	818	190
18.....	860	311	1,790	11,600	3,480	1,510	777	413	2,000	137	637	174
19.....	637	253	1,090	14,900	2,600	1,420	725	345	1,990	122	526	168
20.....	581	280	4,490	13,400	1,990	1,100	673	355	1,340	107	818	164
21.....	432	253	8,600	0,500	1,700	1,160	060	355	874	260	604	345
22.....	374	186	8,900	6,080	1,510	1,080	712	336	000	204	463	660
23.....	311	185	8,050	5,850	1,340	930	764	403	484	185	526	930
24.....	239	158	7,220	5,300	3,250	800	558	463	462	355	570	738
25.....	239	118	6,400	5,850	12,800	818	520	442	384	208	442	526
26.....	268	137	5,580	6,120	6,950	790	604	474	294	204	345	384
27.....	286	163	4,760	5,850	4,220	832	531	484	246	174	302	374
28.....	345	142	6,300	5,030	3,030	874	592	463	211	211	277	336
29.....	320	164	5,850	3,710	860	712	452	190	100	294	294
30.....	294	190	6,120	3,030	888	1,250	452	179	169	423	268
31.....	225	6,120	2,810	902	239	158	547




Figure 2.—Map showing the physiographic provinces in Greenbrier County and surrounding territory as modified after N. M. Fenneman.

FIGURE 3

MAP


OF

GREENBRIER COUNTY

SHOWING

PHYSIOGRAPHIC PROVINCES

PROVINCES



The reasons for shifting this boundary to the
summarized as follows:

1. The presence of a strong anteline (Williamsburg).
2. The presence of an erosion scarp.
3. The change from a trellis to a dendritic drainage pattern.
4. The change from mountains having a general northeast trend to those with no regional trend.

The Valley and Ridge Province, generally includes a series of parallel ridges composed of resistant strata alternating with parallel valleys developed on non-resistant strata. In this county the folds are not so regular nor so severe as in the Ridge and Valley Province as a whole, and as a result the ridge and valley topography is not so well developed as in other parts of the province. Greenbrier River, which has developed the largest valley of the area, in general, parallels the strike of the rocks, but has entrenched itself, with many tortuous meanders, through and over rocks that can well be called resistant to erosion. The trellis or rectilinear pattern of stream drainage is well developed in that part of the county east of the boundary shown in Figure 3. This fact is partially obscured west of the Greenbrier River by subsurface drainage on the Greenbrier Limestone, but is readily apparent in the vicinity of Brushy Ridge.

The western part of the county lies within the Appalachian Plateau Province and presents a different drainage pattern as well as different land forms. Here the mountains attain their highest elevation, (over 4,000 feet A. T.) and their forms are the results of dissection by streams, that have cut deep V-shaped gorges into the elevated plateau. It should be noted that the slope of the ridges and mountain summits is to the northwest, and that the regional dip is also in that direction. In this part of the county the streams have been only slightly controlled by the structure of the rocks and a dendritic (more

Figures 4 and 5 show the difficulty of recognizing any of the older erosion surfaces described in other publications on the physiography of the Appalachian region. In 1925 Wright⁴ described and contoured the "Upland (Schooley) Peneplane" for part of Virginia and West Virginia, including Greenbrier County. In that report the "surface" represented by the northwest sloping ridges and mountain summits of the plateau region are correlated with the "surface" represented by ridge tops east of the Greenbrier River. An examination of Figure 5 offers serious objection to such a correlation. In making this correlation Wright postulated that the greatest uplift occurred a short distance northwest of the present erosion scarp and several local domes are mapped along this line.




Figure 4.—Seven profiles drawn on glass, along the five-minute latitude lines across Greenbrier County. Originals drawn by John P. Nolting. Vertical exaggeration x 26.




Figure 5.—Same as Figure 4, with superimposed projected profiles. Originals drawn by John P. Nolting. Vertical exaggeration x 26.

Warping of the magnitude postulated by Wright, and can be shown by anomalies in structure and it will be noted on Map II (in Atlas) that there is a structural dome in the vicinity of Grassy and Cold Knobs. A distinct synclinal saddle separates this structural dome from the northern end of the Williamsburg Anticline and it appears likely that the dome has a different structural origin than the anticline to the east. Cold Knob (4345' L) Grassy Knob (4372' L) and Job Knob (4338' L) are at nearly the same present elevation, while, structurally, Cold Knob (near the top of the dome) is 1000 feet higher than Job Knob and 300 feet higher than Grassy Knob. About 400 to 500 feet of the difference in the structural elevation between *Cold Knob and Job Knob* may be attributed to the formation of this dome. It appears, therefore, that this structural dome must have been in existence before the formation of the oldest peneplain surface represented in Greenbrier County.

There are no structural irregularities that can be correlated with the warping of the "Upland Peneplane" as shown on Wright's⁶ map. It appears that Wright has contoured the average elevations of the ridge tops in Greenbrier County, but whether or not these average elevations represent one peneplain is seriously questioned.

The absence of a relatively flat and clearly defined Schooley Peneplain in Greenbrier County may be accounted for by one or more of the following hypotheses:

1. During post-Schooley uplift, the surface was subjected to complex warping. This theory has been discussed above.
2. The peneplain surface has been destroyed by post-Schooley erosion. In support of this theory it is noted that the total available relief, in Greenbrier County, of over 2500 feet, which is greater than the elevation of the Schooley in many places, would cause erosion to proceed at a rapid pace. This hypothesis would require two assumptions; (a) complete or nearly complete Schooley planation of the region; (b) that the region in question be at or near the point of greatest uplift. Both of these assumptions are plausible but improved.

In central Greenbrier County there is an area of relatively low relief, developed mainly upon Greenbrier Limestones with the shales and sandstones of the Maecrady and upper Pocono also affected. This area, some six or eight miles wide, crosses the county in a northeast-southwest direction. The surface is best seen around Lewisburg and Frankford where it has an elevation of 2250 to 2350 feet and is approximately 600 feet above Greenbrier River.

In Pocahontas County, Price¹⁴ has described an intermediate erosion surface at an elevation of about 2500 feet or roughly 400 feet above Greenbrier River. Despite the difference of elevation, it is believed that the area of low relief in central Pocahontas County and that in central Greenbrier are of the same age. In Monroe County, Reger¹⁵ has described the same surface as occurring around Union, Pickaway, Monitor, Sinks Grove, and Johnson Crossroads, at an elevation of 2000 to 2200 feet.

In the reports cited above this erosion surface was correlated with the Harrisburg Peneplain of Dauphin County, Pennsylvania, but it now appears that it is more likely to correlate with the Allegheny Peneplain of Ashley.¹⁶

STREAM TERRACES.

Stream terraces are found in many localities along the major streams of Greenbrier County. Prominent local terraces were noted at Harpers, Judyton P. O. (Keister Station), Anthony, and at several other points. It is significant that most if not all of the terraces along the Greenbrier River are on the up-stream side of meanders. This fact suggests that the terraces originated in the normal migration of the meanders of the river and that they can not be correlated with cycles or partial cycles of erosion.

¹⁴Price, P. H., Pocahontas County Rept., W. Va. Geol. Sur., pp. 24-25, 1929.

¹⁵Reger, D. B., Mercer, Monroe and Summers Counties Rept., W. Va. Geol. Sur., pp. 62-63, 1926.

¹⁶Ashley, Geo. H., Scenery of Pennsylvania, Pa. Geol. Sur., Bull. G-6, pp. 23 ff., 1933. See also, Fridley, H. M., and Nolting, J., Peneplains of the Appalachian Plateau, Jour. Geol., Vol. 39, pp. 749-756, 1931.

3. Wasting of the Schooley surface with greater reduction on the softer and/or the more steeply dipping rocks has resulted in an uneven and poorly defined level. Wasting and reduction of a peneplain, with or without loss of its horizontality, has been advocated by Hayes⁷, Fenneman⁸, Wright⁹, and Ashley¹⁰. Each writer suggests that the reduction of the Schooley Peneplain in some places may be measurable in hundreds of feet. This hypothesis, like the second, can not be entirely excluded because wasting is practically certain to have had some effect. To stand alone, however, it requires the assumption that the area in question was completely or nearly completely leveled in Schooley time.

4. The region under discussion was near the headwaters of pre-Schooley streams and was never completely reduced, i. e., a monadnock area or a divide with considerable relief. Most of the present physiographers agree that the present main drainage systems antedate the Schooley Peneplain¹¹. If this is true it is necessary that the Schooley streams have some gradient and major divides were probably somewhere near their present location. Fenneman¹² recognizes Wright's¹³ delineation of the Schooley surface and apparently accepts the warping hypothesis. On page 260 of the same book, however, Fenneman states:

"Remarkable as the Schooley peneplain was, its perfection must not be overestimated. Streams were not left without gradient nor divides without slope."

⁷Hayes, C. W., *Physiography of the Chattanooga District*, U. S. Geol. Sur. Ann. Rept., pt. 2, p. 26, 1899.

⁸Fenneman, N. M., *Jour. Geol.*, Vol. 16, pp. 746-754, 1908; *Bull. Geol. Soc. Amer.*, Vol. 47, pp. 173-186, 1936; *Physiography of Eastern United States*, McGraw-Hill, pp. 199-200, 1938.

⁹Wright, F. J., *The Older Appalachians of the South*, *Jour. Sci. Lab. Denison Univ.*, Vol. 26, p. 156, 1931.

¹⁰Ashley, G. H., *Bull. Geol. Soc. Amer.* Vol. 46, p. 1403, 1935.

¹¹See, Ver Steeg, Karl, *Wind Gaps of the Northern Appalachians*, *Ann. N. Y. Acad. Sci.*, Vol. 32, pp. 87-220, 1930; also Johnson, Douglas, *Stream Sculpture on the Atlantic Slope*, Columbia Univ. Press, 1931; also Meyerhoff, H. A. and Olmstead, E. W., *The Origin of the Appalachian Drainage*, *Amer. Jour. Sci.*, 5th ser. Vol. 32, No. 187, pp. 21-42, July 1936; also Fenneman, op. cit., pp. 260.

FLOOD-PLAINS.

Probably the most interesting physiographic feature in Greenbrier County is the existence of comparatively broad, local flood-plains along many of the streams. From west to east across the county, streams that have formed broad flood-plains are as follows:

1. Meadow River and tributaries (2425).
2. Muddy Creek (1625).
3. Sinking Creek (2190).
4. Roaring Creek and Little Roaring Creek (2275).
5. Howard Creek (1800).
6. Anthony Creek (1925-1950).

The figures in parenthesis are the approximate elevations of the major flood-plain along the stream indicated. The fact that no two flood-plains are near the same elevation indicates that each is due to local conditions.

Meadow River has developed a large flood-plain that is the result of planation of non-resistant rocks behind a barrier of rock resistant to erosion. The river and its tributaries have base-leveled 15 to 20 square miles of area, cutting across different beds of the Mauch Chunk Series. The chief barrier appears to be that of the Pottsville sandstones which dip below stream level about one and a half miles northwest of Rainelle. The Princeton Sandstone goes below drainage between Rupert and Charcoal and was undoubtedly a contributing factor.

The flood-plains of Muddy Creek, Sinking Creek, and Roaring Creek have been developed at or near the contact of the Greenbrier Limestone and the overlying Mauch Chunk shales. Apparently the limestone is sufficiently resistant to surface erosion to act as a barrier, holding up the stream and thereby causing planation of the non-resistant shales. These three flood-plains suggest a method that may have operated in the past to expose a part of the vast area of Greenbrier Limestone outcropping in the county.

Howard Creek has developed a rather large flood-plain on the shales of the lower Portage, Genesee, and Marcellus. The chief barriers to the local planation along this stream are the

Along Anthony Creek the conditions favorable to local base-leveling are found in two localities and as a result two comparatively broad flood-plains have been developed. In each case the shales and thin sandstones upon which the flood-plains have been developed are of Portage, Genesee, and Mareellus age. The flood-plain west of Alvon narrows abruptly as the Chemung-Portage contact is crossed and disappears just west of Blue Bend Forest Park. It is apparent that the Chemung sandstones form the erosion barrier. Whether the flood-plain east of Alvon is genetically separate from the one just described is open to question. However, the presence of rapids in the gorge, just west of Alvon, and the fact that the eastern flood-plain is at a somewhat higher elevation than the one west of Alvon, indicate that the Lower Devonian and Silurian rocks in the gorge have been effective barriers to erosion. It is probable that the flood-plain east of Alvon would not have been so extensive if the Chemung sandstones had not, in effect, decreased the stream gradient west of Alvon.

PRESENT TOPOGRAPHIC FEATURES.

All of the mountains in Greenbrier County that have an elevation of 4000 feet or over are in the northwest part of the county and in each case the mountain is capped by Pottsville sandstone. The major drainage channel in the western part of the county have elevations from 2400 to 3000 feet, making the net height of the mountains 1000 to 1500 feet. The highest point in the county is Grassy Knob with an elevation of 4372 A. T. Other points above 4300 are Cold Knob (4345) and Job Knob (4338). The lowest point in the county is where the Greenbrier River leaves the county just west of Alderson with an elevation of approximately 1520 feet.


East of the Greenbrier River the mountains rarely reach an elevation of 3500 feet and the majority have an elevation between 2750 and 3500 feet. The major streams are at elevations of 1800 to 2300 making the net height of the mountains 700 to 1500 feet. From these figures it may be seen that the topographic relief of the eastern part of the county is practically as great as that of the western part.

In the main, the relatively flat area is underlain by the Greenbrier Limestone and is characterized by the presence of hundreds of sink-holes.

DRAINAGE BASINS.

A general view of the drainage system of Greenbrier County can be seen on Figure 6, and a detailed study can be made of the streams from Maps I and II, which are found in the Atlas accompanying this report. East of the plateau region the major streams, in general, parallel the mountain ranges, while the minor streams have cut across them at right angles, a condition that prevails throughout the Allegheny Ridges region. In the Plateau region the streams have followed the lines of least resistance or down the regional dip.

DRAINAGE



The following table by Professor Geo. W. Grow gives a list of all the principal streams of Greenbrier County, the length of the streams as well as the air-line distance from source to mouth, also the total fall of the streams and rate of fall per mile. In the last column is given the ratio of the meander distance or total distance (T. D.) to the air-line distance (A. L. D.):

Table of Stream Data.

STREAMS.	Total Distance Miles.	Total Fall Feet.	Rate of Fall per Mile. Feet.	Air-Line Dis- tance. Miles.	Ratio, Total Distance to A. L. D.
Greenbrier River, source to East Fork to mouth.....	164.8	2500	15.17	98.64	1.67
Greenbrier River, source of West Fork to mouth.....	162.9	2250	13.81	97.14	1.68
Greenbrier River, Pocahontas Co. line to Summers Co. line....	55.61	432	7.77	31.86	1.75
Greenbrier River, Summers-Greenbrier Co. line to mouth....	26.76	130	4.86	13.21	2.03
Greenbrier River, junction of East and West Forks at Durbin to mouth.....	144.0	1325	9.20	83.49	1.72
Muddy Creek.....	19.20	750	12.96	12.15	1.58
Mill Creek.....	7.90	955	120.89	6.61	1.19
Kitchen Creek.....	8.30	900	108.43	5.75	1.44
Saw Mill Hollow.....	2.40	525	218.75	1.70	1.41
Lorenze Creek.....	4.40	700	159.09	3.73	1.18
Snake Run.....	5.70	1250	219.30	4.90	1.16
Alum Run.....	4.10	640	156.10	3.15	1.30
Second Creek.....	23.80	990	41.58	14.89	1.60
Howard Creek.....	14.30	555	38.81	10.65	1.34
Monroe Draft.....	4.65	935	203.23	4.04	1.15
Harts Run.....	6.86	550	80.17	5.51	1.25
Roekliek Run.....	2.15	760	353.49	1.92	1.12
Dry Creek.....	9.45	810	85.71	7.60	1.24
Broad Run.....	2.20	605	275.00	1.97	1.12
Tuekahoe Run.....	4.05	515	127.16	3.55	1.14
Quarry Hollow.....	2.05	715	348.78	1.97	1.04
Spring Run.....	4.80	655	136.46	4.39	1.09
Jericho Draft.....	4.40	480	109.09	3.99	1.10
Sulphur Lick Run.....	3.20	575	179.69	2.91	1.10
Pond Liek Run.....	2.85	655	129.82	2.47	1.15

STREAMS.	Total Distance Miles.	Total Fall Feet.	Rate of Fall per Mile. Feet.	Air-Line Dis- tance. Miles.	Ratio, Total Distance to A. L. D.
Boulder Run.....	2.37	980	413.50	2.27	1.04
Anthony Creek.....	27.99	1425	50.75	22.36	1.25
Lanrel Creek.....	2.96	1050	354.73	2.76	1.07
Blg Draft.....	2.76	360	130.43	2.27	1.22
Rocky Run.....	2.86	1130	395.10	2.32	1.23
Little Creek.....	8.38	725	86.52	7.99	1.05
Dawson Run.....	1.58	705	446.20	1.53	1.03
Pantherlick Run.....	1.53	505	330.07	1.50	1.02
Fleming Run.....	4.39	435	89.09	4.00	1.10
Whitmans Draft.....	4.39	780	177.68	3.70	1.19
Whites Draft.....	4.44	765	172.30	3.95	1.24
Wades Draft.....	3.95	925	234.18	3.76	1.05
Turkeypen Run.....	1.48	610	412.16	1.31	1.13
Wiley Run.....	2.37	760	320.68	2.27	1.04
Humphreys Draft.....	2.76	660	239.13	2.38	1.16
Barnes Lick Run.....	2.07	715	345.41	1.91	1.08
Stony Run.....	2.47	605	244.94	2.27	1.09
Sims Run	2.37	475	200.40	1.97	1.20
Meadow Creek.....	14.06	1415	100.64	11.10	1.27
Laurel Creek.....	4.44	656	147.75	4.27	1.04
North Fork of Anthony Crk.	12.33	1205	97.73	11.25	1.10
Onemile Run.....	1.68	445	264.88	1.58	1.06
Twomile Run.....	1.97	655	332.49	1.68	1.17
Fourmile Run.....	1.59	405	254.72	1.48	1.07
Hoffman Run.....	1.48	355	239.86	1.28	1.16
Coles Run.....	2.47	675	273.28	2.07	1.19
Pondlick Run.....	1.18	330	279.66	1.09	1.08
Sugar Run.....	2.47	765	309.72	2.22	1.11
Bear Branch.....	1.82	585	321.43	1.78	1.02
Laurel Run.....	7.69	1055	137.19	5.53	1.39
Boardhouse Run.....	1.68	750	446.43	1.53	1.10
Spring Creek.....	21.31	1675	78.60	11.30	1.89
Dry Run.....	2.71	1110	409.59	2.47	1.10
Robbins Run.....	7.05	1405	199.29	4.34	1.62
Boggs Run.....	1.87	1030	550.80	1.68	1.11
Rockcamp Run.....	2.17	570	262.67	2.08	1.04
Panther Camp Creek.....	3.45	1055	305.80	2.77	1.52
Board Lick Run.....	1.23	1150	934.96	1.14	1.08
Wolfpen Run.....	2.07	520	251.21	1.99	1.04
Boggs Run.....	1.68	370	220.24	1.66	1.04
Blg Run.....	1.97	650	329.95	1.89	1.04
Snodgrass Run.....	3.95	790	200.00	3.16	1.25
Slabeamp Run.....	5.18	800	154.44	3.46	1.50
Red Run.....	1.88	505	268.61	1.73	1.09
Kincaid Run.....	3.06	1125	367.65	2.86	1.07

STREAMS.	Total Distance Miles.	Total Fall Feet.	Rate of Fall per Mile. Feet.	Air-Line Dis- tance. Miles.	Ratio, Total Distance to A. L. D.
Milligan Creek (surface length only).....	6.61	385	58.25	5.30	1.25
Culverson Creek (surface length only).....	9.47	400	42.24	5.92	1.60
Burns Run.....	2.96	600	222.97	2.45	1.21
Spice Run.....	2.76	210	76.09	2.09	1.32
Indian Creek.....	3.45	810	234.78	3.02	1.14
Sinking Creek (surface length only).....	12.53	1450	157.22	5.87	2.13
Hughart Creek.....	4.93	800	162.27	4.57	1.08
Flynn Creek.....	4.19	1835	437.95	3.28	1.28
Roaring Creek (surface length only).....	5.23	1715	327.92	3.24	1.61
Little Roaring Creek.....	2.86	1400	489.51	2.09	1.37
Meadow River.....	52.58	1620	30.81	31.09	1.69
Anglins Creek.....	12.53	1455	116.12	9.76	1.28
Youngs Creek	5.57	860	154.40	3.77	1.48
North Prong	2.52	790	313.49	2.37	1.06
Spring Creek.....	1.68	580	345.24	1.63	1.03
Haynes Branch.....	1.59	690	433.96	1.53	1.04
Burdette Creek.....	3.45	970	281.16	3.41	1.01
Piney Creek.....	1.73	305	176.30	1.58	1.09
Toms Creek.....	2.86	470	164.34	2.82	1.01
Meadow Creek.....	8.24	340	41.26	7.11	1.16
Sewell Creek.....	10.16	540	53.15	8.13	1.25
Little Sewell Creek.....	4.73	240	50.74	4.14	1.14
Boggs Creek.....	5.13	285	55.56	4.74	1.08
Wolf Pen Creek.....	2.71	670	247.23	2.14	1.27
Little Creek.....	3.45	755	218.84	3.04	1.13
Laurel Creek.....	3.55	785	221.12	3.19	1.11
Mill Creek.....	5.62	1085	193.06	5.16	1.09
Big Clear Creek.....	14.30	1110	77.62	12.41	1.15
Brown Creek.....	5.23	1220	233.27	4.94	1.06
South Fork.....	8.88	1375	154.84	7.75	1.15
Smokehouse Branch.....	2.27	625	275.33	2.22	1.02
Old Field Branch.....	2.86	665	232.52	2.78	1.03
Job Knob Branch.....	3.95	965	244.30	3.28	1.20
Sam Creek.....	2.95	580	120.43	2.81	1.05
Elijah Branch.....	2.07	420	202.90	1.97	1.05
Road Branch.....	1.49	405	271.81	1.30	1.15
North Fork.....	3.06	305	99.67	2.86	1.07
Little Clear Creek.....	14.11	1655	117.29	10.53	1.34
Beaver Creek.....	4.44	845	190.32	3.43	1.29
Stony Run.....	2.38	1320	554.62	2.29	1.04
Rader Run.....	2.27	1380	607.93	2.17	1.05
Laurel Creek.....	3.06	585	191.18	3.02	1.01

STREAMS.	Total Distance Miles.	Total Fall Feet.	Rate of Fall per Mile. Feet.	Air-Line Dis- tance. Miles.	Ratio, Total Distance to A. L. D.
Otter Creek.....	6.31	355	56.26	2.38	2.65
Methodist Branch.....	2.56	35	13.67	2.41	1.06
Smoot Branch.....	1.97	35	17.77	1.85	1.06
Eagle Branch.....	2.22	70	31.53	1.63	1.36
Buffalo Creek.....	4.04	325	81.05	3.95	1.02
Morris Branch.....	4.78	245	51.26	3.30	1.45
Patterson Creek.....	2.47	140	56.68	2.20	1.12
(Gauley River)					
Hominy Creek.....	22.59	2220	98.27	14.91	1.52
Prlec Fork.....	2.81	445	158.36	2.56	1.10
Preaser Branch.....	2.56	605	236.33	2.27	1.13
(Cherry River)					
Laurel Creek.....	15.14	1550	102.38	12.05	1.26
McMillon Creek.....	4.39	1010	230.07	4.00	1.10
Mill Branch.....	1.97	580	294.42	1.87	1.05
Beech Run.....	2.66	740	278.20	2.52	1.06
Hogcamp Run.....	2.71	860	317.34	2.39	1.31
Manning Branch.....	2.71	855	315.50	2.53	1.07
Middle Branch.....	2.81	730	259.79	2.71	1.04
Cold Spring Branch.....	3.21	760	236.76	2.70	1.19
Linn Branch.....	1.97	770	390.86	1.62	1.22
Little Laurel Creek.....	6.71	950	141.58	5.30	1.27
Baber Branch.....	1.82	980	538.46	1.73	1.05
Improvement Branch.....	2.96	700	236.49	2.50	1.18
South Fork of Cherry River.....	16.28	1860	114.25	10.60	1.54
Shiras Run.....	1.63	830	509.20	1.34	1.22
Elklick Run.....	1.68	920	547.62	1.61	1.04
Rocky Run.....	5.08	1235	243.11	3.51	1.45
Little Rocky Run.....	1.97	1070	543.15	1.83	1.08
Becky Run.....	2.27	710	312.78	1.94	1.17
Cold Knob Fork.....	5.18	755	145.75	4.49	1.15
Blizzard Run.....	1.43	875	611.89	1.35	1.06
Little Blizzard Run.....	1.18	640	542.37	1.06	1.11
Blg Run.....	1.68	580	345.24	1.56	1.08
North Fork of Cherry River.....	17.26	2045	118.48	10.62	1.63
Coats Run.....	2.17	985	453.92	1.85	1.17
Little Lick Run.....	1.43	750	524.48	1.20	1.19
Windy Run.....	1.48	470	317.57	1.33	1.12
Armstrong Run.....	1.13	490	433.63	0.986	1.15
Hamrick Run.....	1.48	590	398.65	1.13	1.31
Rabbit Run.....	0.94	305	324.47	0.70	1.34
Carpenter Run.....	1.18	550	368.10	1.12	1.05
Deacon Run.....	0.99	500	505.05	0.95	1.04
Fallen Timber Run.....	1.23	640	520.33	1.01	1.22
Bear Run.....	2.27	480	211.45	2.03	1.12
Dogway Fork (of Cranberry					

The following table by Professor Geo. W. Grow gives a list of the principal streams of Greenbrier County with their drainage areas computed by planimeter from the topographic maps:

Areas of Drainage Basins.

STREAMS	Square Miles.
Greenbrier River, entire.....	1,634.65
Greenbrier River, in Greenbrier County.....	679.02
Greenbrier River, in Pocahontas County.....	687.06
Muddy Creek.....	75.91
Mill Creek.....	15.74
Kitchen Creek, Total area.....	26.50
Kitchen Creek, in Greenbrier County.....	25.77
Kitchen Creek, in Summers County.....	0.73
Saw Mill Hollow.....	4.25
Lorenze Creek.....	5.75
Snake Run.....	7.66
Alum Run.....	6.87
Second Creek, entire.....	116.34
Second Creek (Greenbrier County).....	13.86
Howard Creek (entire).....	91.01
Monroe Draft.....	6.67
Harts Run.....	11.84
Rock Lick Run.....	1.36
Dry Creek.....	22.88
Broad Run.....	2.50
Tuekahoe Run.....	9.33
Quarry Hollow.....	2.11
Spring Run.....	7.11
Jericho Draft.....	12.49
Sulphur Lick Run.....	4.95
Pond Lick Run.....	2.01
Slash Lick Run.....	3.47
Boulder Run.....	2.17
Anthony Creek (entire).....	147.87
Laurel Creek.....	2.54
Big Draft.....	3.67
Rocky Run.....	1.86
Little Creek.....	15.05
Dawson Run.....	0.75
Panther Lick Run.....	0.87
Fleming Run.....	9.14
Whitmans Draft.....	2.90
Whites Draft.....	7.12
Wades Draft.....	3.79
Turkeypen Run.....	0.79
Wiley Run.....	3.48
Humphreys Draft.....	2.31
Barnes Lick Run.....	0.82
Stony Run.....	1.11

Falling Springs or Renick Post-Office and Station is located sixteen miles north of Lewisburg, on the Greenbrier River, and is served by the Greenbrier Division of the Chesapeake and Ohio Railway and the Seneca Trail (State Route 24).

The town is supplied with a bank, an electric milling company, a limestone quarry, and several mercantile establishments that furnish supplies for the immediately surrounding area.

The population according to the 1930 Census report was 355.

WILLIAMSBURG.

Williamsburg, a strictly agricultural village, surrounded by good farms, is located near the center of the county in a limestone area. A hard-surfaced road connects the town with the Midland Trail, but there are no railroad facilities. The town is supplied with good schools and churches.

The population in 1930 was 148.

Villages.—Other small villages with approximate populations are as follows: Frankford, 140; Neola, 125; Anthony, 50; Fort Spring, 150; Clintonville, 50; Rupert, 300; Quinwood, 500; Leslie, 200; Bellburn, 150; Anjean, 300; Duo, 100; Clearco, 150.

TRANSPORTATION

WATERWAYS.

Since the coming of railways in the county, waterways have played a very minor role. Prior to that time, however, the larger streams and particularly Greenbrier River were used to float logs to band mills that were set up at strategic points. The Greenbrier River was well suited for that purpose as it carries a considerable volume of water and has a fairly low gradient, averaging 7.7 feet fall per mile across the county, a distance of 55.6 miles.



PLATE IV.—View from U. S. Route 60 near Clintonville showing Manch Chunk and Greenbrier topography. Shaped Weaver Knob can be faintly seen in the background.



PLATE III.—View of Howard Creek Valley from Kates Mountain, looking west toward Caldwell. Water-gap in background cut in rapidly dipping Focono and Chemung rocks. Photo. by Cummins.



PLATE II.—Front view of Greenbrier Hotel at White Sulphur Springs. Photo. by Cummins.



TE V.—View from U. S. Route 219, near Renick, showing Greenbrier Limestone topography with sink-holes

due to methods of transportation other than waterways.

RAILROADS.

Chesapeake and Ohio—Main Line.

The construction of the Chesapeake and Ohio Railway into West Virginia in 1873 (to White Sulphur in 1869) was as important in the development of Greenbrier County in comparison as the construction of this road was to the development of the State as a whole. The main line extends from Fortress Monroe, Virginia, westward across Virginia, West Virginia, and other States. The line is now equipped with double tracks (completion of double track in tunnels, 1932) and is doing a large business in coal, general freight, and passenger service.

This railroad enters Greenbrier County at the Allegheny Tunnel on Allegheny Mountain at the Virginia State line one mile east of Tuckahoe and follows the drainage of Dry Creek to White Sulphur Springs; thence along Howard Creek to its junction with the Greenbrier River at Caldwell, thence following the river, excepting the two tunnels near Fort Spring, to a point near Alderson where it enters Monroe County.

As the corporate history of the Chesapeake and Ohio Railroad has already been published in one of the Survey Reports¹ it is not deemed advisable to reproduce it here, but because of the importance of the construction of this road into West Virginia the reader is here referred to it.

Chesapeake and Ohio—Greenbrier Division.

The Greenbrier Division is a branch from the main line at Whitecomb, this county, and extends entirely across it northward, following the Greenbrier River to its northern termination at Winterburn Station (Thornwood P. O.), Pocahontas County. At Durbin it connects with the Western Maryland Railroad. The construction of this branch began in 1899 and was completed to Winterburn in 1905. Inasmuch as the main line served only the southern end of the county the completion

consolidated with the Sewell Valley Railroad Company on March 1, 1928. On December 30, 1931, the Sewell Valley Railroad Company and the Loop and Lookout Railroad Company, and the Greenbrier & Eastern Railroad Company were consolidated with the Nicholas, Fayette, and Greenbrier Railroad Company as a single corporation, the charter of the first three named roads being surrendered to the State of West Virginia.

"On January 6, 1932, joint operation of the consolidated properties was established by the two roads owning the property. The Nicholas, Fayette and Greenbrier Railroad is now owned jointly by The Chesapeake & Ohio Railway Company and The New York Central Railroad Company, each owning one-half interest. The Sewell Valley and the Loop and Lookout Railroads were previously owned by Mr. T. W. Raine and his associates, who built these roads. The Greenbrier & Eastern Railroad Company was previously owned by Coal Companies or their representatives, who were located on this line."

As described in the above letters, the Nicholas, Fayette, and Greenbrier Railroad leaves the main line of the Chesapeake and Ohio Railroad Company at the town of Meadow Creek, Summers County, and follows Meadow Creek to Springdale, Fayette County. From there it follows Sewell Creek north to Bellwood, thence northeast to Rainelle and East Rainelle, Greenbrier County. From here a branch follows along Meadow River, eastward to Rupert, thence northward along Big Clear Creek to Anjean, Duo and its termination at Cleareo. There is also a short branch from Rupert to Little Clear Creek and just below Anjean a branch logging road extends up Brown Creek. This line from Rainelle to Cleareo, is used for transportation of logs to the Meadow River Lumber Company and provides an outlet for the coal from the mines at Midland, Anjean, Duo, and Cleareo.

The branch of the Nicholas, Fayette, and Greenbrier Railroad that was formerly known as the Greenbrier and Eastern crosses Meadow River at East Rainelle and follows the north side of that stream to the mouth of Meadow Creek, thence northeast along Meadow Creek to Bellburn, Leslie, Crichton, Quinwood, and Marfrance. This branch is the outlet for the coal from the many commercial mines near the above-named towns.

The portion of this railroad along the southwest side of Meadow River from East Rainelle to Burdette Creek was constructed in 1911. At Burdette Creek the railroad crosses to

Nicholas, Fayette, and Greenbrier Railroad Company.

On December 30, 1931, the Sewell Valley Railroad Company, the Greenbrier and Eastern Railroad Company, and the Loop and Lookout Railroad Company were consolidated with the Nicholas, Fayette, and Greenbrier Railroad Company.

The following quotation from a letter from Mr. J. W. Raine, President of The Raine Lumber and Coal Company, summarizes the history of the Sewell Valley Railroad:

"Duo, W. Va., October 8, 1937.

"Father (T. W. Raine) began construction of the Sewell Valley Railroad at Meadow Creek the spring of 1908. It was completed to Rainelle in February 1910. In 1911 it was completed to the mouth of Burdette Creek. During 1916 it was extended to Nallen to serve the Wilderness Lumber Company. The branch from Rainelle to Rupert and Glencoe was begun in 1920 and completed 1922. The Big Clear Creek extension was begun 1927 and completed in 1929.

"The Sewell Valley Railroad was owned by The Meadow River Lumber Company from the beginning until July, 1921, when my father and brothers purchased it. They sold it to the Chesapeake and Ohio Railway Company in July 1927.

"Father (T. W. Raine) was not connected with the Greenbrier & Eastern. This was begun in 1920, I think, and was sold to the Chesapeake and Ohio about the same time that they bought Sewell Valley.

"The main traffic at the first was lumber and continued so until about 1920. From that time on coal business developed and now it is the principal traffic."

The following quotation from a letter from Mr. J. M. Raine, Assistant Superintendent of the Nicholas, Fayette, and Greenbrier Railroad Company, summarizes the corporate history of this company:

"Rainelle, W. Va., November 29, 1937.

"The original Nicholas, Fayette, and Greenbrier Railroad Company projected from Swiss, West Va., on the New York Central to Nallen, West Va., to the Loop & Lookout Railroad Company (Sewell Valley Railroad Co., Lessees). Construction started in February, 1929, and completed in October 1930—distance 28.2 miles, and the track was put into operation January 6, 1932.

"At the present time there is no traffic originating directly on this

constructed in what is now Greenbrier County was the James River and Kanawha Turnpike. An early writer who traveled over the route pronounced it "one of the principal chains destined by nature to bind together the eastern and western portions of this great republic." The need for such a route was brought to the attention of the Virginia Assembly by Washington in 1784 and was promptly passed in an act incorporating the James River Company, and in 1785 authorized the construction of the "State Road" (for wagons) which was completed to the navigable waters of the Kanawha by 1790 and opened to the Ohio by 1800 (For a more complete history of this road see Calahan's Semi-Centennial History of West Virginia, 1913).

The present U. S. Highway No. 60 enters Greenbrier County from Virginia on Allegheny Mountain at a point four miles east of White Sulphur Springs. Passing through the latter town it follows the general course of the old turnpike, but with several rather important new locations, crosses the Greenbrier River to Lewisburg, continues west through Richlands, Clintonville, Rupert, Rainelle, and leaves the county to enter Fayette just west of Rainelle.

It continues west across Fayette, traversing rugged and beautiful scenery along the New River gorge to Charleston, thence on west to Huntington on the Ohio by way of Teays Valley.

This route is now one of the most important east and west highways and because of its scenic grandeur is very popular with tourists.

U. S. No. 219 (formerly State Route No. 24) or what is otherwise generally known as the **Seneca Trail**, is another important highway passing through Greenbrier that crosses the State from north to south. It enters West Virginia three miles south of Red House, Maryland, and continues southwest across West Virginia through Thomas, Parsons, Elkins, Huttonsville, Valley Head, Marlinton, and enters Greenbrier County on the south side of Droop Mountain. In Greenbrier it continues southwest generally paralleling the Greenbrier River through Renick, Frankford, and Maxwelton to Lewisburg where it crosses U. S. Route No. 60. From Lewisburg it

to Russellville. Just north of the town the railroad crosses to the southwest side of Meadow River, following the river to Nallen, Fayette County. The part of the line from Burdette Creek to Nallen was constructed in 1916-17. This railroad between East Rainelle and Nallen was known as the Loop and Lookout Railroad previous to its consolidation with the Nicholas, Fayette, and Greenbrier Railroad. The branch railroads leaving the main line at Burdette Creek, extending up that creek, and along the south side of Meadow River, are logging roads.

The original Nicholas, Fayette, and Greenbrier Railroad connects with the Loop and Lookout at Nallen and follows Meadow River and Gauley River to Swiss, Nicholas County, crossing Gauley River from the south to the north side of the mouth of Peters Creek.

HIGHWAYS.

State Roads.

Road building in West Virginia has progressed rapidly since the legislative enactment of 1921 with the organizing of a State Road Commission, and following a definite plan of construction. In this road building program Greenbrier County has received its proportionate share of new roads. It is true that the county was traversed from both north to south and east to west by two well-established through routes, both of which, however, needed much improvement to meet the needs of modern traffic. These two routes, U. S. 60 and W. Va. 24, (now U. S. Route 219), have both been straightened, widened, and hard surfaced under the new program and are now a part of two of the most widely traveled routes in the State.

From the 1936 edition of the State Road Map, issued by the State Road Commission, and in conjunction with the more detailed topographic maps, the following descriptions of U. S. and State routes in Greenbrier County have been compiled. Their terminals in other counties or at the State line have been indicated.

leaving the general course of the river to the westward, enters Monroe County at Second Creek. From here it continues generally southwestward through Union, Peterstown, Princeton, and finally Bluefield at the Virginia State line. This route is now graded and paved throughout its entire length in West Virginia. It is one of the most scenic and picturesque highways in the State and is fast becoming a favorite with tourists.

State Route No. 54 is a short route connecting State Route No. 3 at Alderson with U. S. Route No. 60 at Alta. It lies entirely within the boundaries of Greenbrier County and is entirely paved.

State Route No. 3 lies within the limits of Greenbrier County for a distance of only about a mile, this being from Alderson to the Summers County line. This route, however, starts near Sweet Springs, Monroe County, coincides with U. S. Route No. 219 between Union and Piekaway, and proceeds westward across that county to Alderson; thence southwestward and westward to Hinton, Beekley, through a number of small towns in Raleigh, Boone, and Lincoln Counties to join State Route No. 10 at West Hamlin. This road is paved throughout except for a short distance between Woodville and Yawkey, Lincoln County.

State Route No. 44 originates on State Route No. 39 at Nettie, Nicholas County and proceeds southward entering Greenbrier County a few miles north of Quinwood, passing through that town. Continuing southward it joins U. S. Route No. 60 at Charleo and coincides with it through East Rainelle to Rainelle where it leaves U. S. Route No. 60. This route leaves Greenbrier about two and one-half miles southwest of Rainelle; thence southward to Hinton, to Athens and ends at its junction with U. S. Route 219 near Princeton, Mercer County. It is paved from Nettie to Rainelle and between Athens and its junction with U. S. Route No. 219 but is only graded or unimproved for the other 52 miles.

State Route No. 63 is a proposed road connecting Alderson and Benmar, wholly in Greenbrier County, that is par-

The bath establishment, instituted about 1860, has since that time been under the management of Dr. George Kahio, after an exhaustive study of the foreign cures, is the most complete and luxurious in its appointments of all institutions of its kind in the United States. Sulphur-water baths are a special feature.

"In general, it may be said that the White Sulphur Springs waters are of the highest value in conditions associated with impaired digestion, disturbed metabolism, and insufficient elimination. Conditions resulting from an accumulation of toxins, such as gout, rheumatism and arthritis, are acted upon favorably by the waters and treatment.

"Contributed to by both environment and tradition, the life at White Sulphur is probably equaled in charms at no other American resort. There are three splendid golf courses and several tennis courts. Several hundred miles of carefully laid-out mountain riding trails and a large stable of Virginia-bred and Kentucky-bred saddle horses lend variety to the sports life. There are good roads for motoring. Beautiful scenery² and clear, stimulating air at an altitude of nearly three thousand feet add to the attractions of the resort.

"The modern Greenbrier Hotel,³ a beautiful Georgian structure standing in parklike grounds of several thousand acres and adjacent to the former site of the Old White Hotel, is charmingly situated in the valley of the Greenbrier River⁴ at the base of the towering ridges of the Alleghenies as they sweep through the West Virginia country toward the Ohio Valley. Built about fifteen years ago, it is on a par with the finest hotelries in the large Eastern and Western cities. A large cottage colony surrounds the Greenbrier.

"The resort is pleasantly and comfortably reached by an over-night trip from nearly all the large Eastern and mid-Western cities."

There are many points of geological interest on and around the grounds of the resort. Every series of the Devonian outcrops on the resort grounds and much of the Mississippian and the Silurian are exposed at near-by points. There are numerous fossils to tempt the collector and jointing often leaves interestingly shaped rocks (see Plate XXXII). There are numerous illustrations of anticlines and synclines in the area, with the most striking example in the Anthony Creek gorge at Alvon. The Oriskany Sand, that is such a prolific producer of gas in Kanawha County, outcrops on Bobs Ridge, Coles Mountain, and on Beaverliek Mountain.

The geology of the springs of the resort is quite interesting, but description of these is reserved for Chapter XIII.

With an ideal climate, Greenbrier County, of course, has a number of summer camps. Camp Greenbrier, a scout camp,

²See Plate III and others in this report.

³See Plates I and II in this report.

and proceeds to Richwood via Nettie and Fenwick. The proposed extension of this route crosses the northern end of Greenbrier County and connects Richwood, Nicholas County, to U. S. Route No. 219 at Mill Point. The extension is only partially graded.

The greater part of the remaining area is covered with a network of dirt roads, several miles of which are now paved. There remain, however, several areas, namely North and South Forks of Cherry River, Little Clear Creek, North Fork of Anthony Creek and Meadow Creek west of Allegheny Mountain, to which access is quite difficult.

AIRPORT.

Airport.—Air travel is just coming into prominence in West Virginia. The only commercial airport in Greenbrier County is located near White Sulphur Springs. The following information is reprinted from the State Road Commission Map:

"White Sulphur Springs—1.5 mi. S. W. on U. S. Highway 60; 1 mi. S. W. of Greenbrier Hotel. Alt. 1,795. 5,000 by 2,000 turf; level. Pole line to N. W., woods to N., trees along creek to S. and W. Service day only."

RESORTS AND SUMMER CAMPS.

What is perhaps the most famous resort in the western hemisphere is located in Greenbrier County. With historie and soeial tradition reaching baek for more than a century and a half, White Sulphur Springs has become one of the institutions of this county and it is visited by thousands each year.

The following quotation is taken from the West Virginia Encyclopedia, West Virginia Publishing Company, Charleston, W. Va., pp. 1005-6, 1929:

"While the general public, perhaps, looks upon White Sulphur as a gathering place for the fashionable society of the country during the spring, summer, and fall seasons, its importance as a health-giving resort is not secondary. A superior thermal and medical equipment which provides for all forms of hydrotherapy, including such special baths as are given at Nauheim, Aix-les Bains, Carlsbad, Vlchy, and other foreign spas, makes it compare favorably with any of the Eu-

Camp Anagnany, Camp Loopmount, and many others are located along the Greenbrier River.

Both the State and Federal governments have established forests and parks in the area adjacent to the Greenbrier-Pocahontas County line and each year the county becomes more and more popular with vacationists and tourists.

CHAPTER II.

PHYSIOGRAPHY.

INTRODUCTION.

In any area the present land surface, or the distribution of land forms, i. e., mountains and valleys, caverns, etc., is the result, or the expression, of the interaction of earth forces with those of the atmosphere, and represents the geologic history of the region during the time it has been a land area. The Appalachian System, of which the local area is a part, constitutes one of the oldest mountain chains of the earth, and still retains certain features that go back to the Tertiary or Cretaceous Age.

Greenbrier County lies near the source of several of the major streams of the eastern United States. In this area, as in any other area, the streams are the oldest surviving remnants, and represent by far the most important factor in the development of the present land outline. The character and position of the strata, upon which the land forms are developed, will influence and in part control their development. The rocks of Greenbrier County—sandstone, limestone, and shale—are all of sedimentary origin; that is, deposited from a transporting medium, generally in water of varying depths and salinity, while the coals represent abundant vegetation spread over a low-lying swamp area, but in sufficient water to prevent decomposition, which would follow if not arrested by the formation of a toxic acid that prevents bacterial decay.

Let us consider then for a moment the important events in the geologic history of the eastern United States that directly concern this area. Suffice it is to say that since all the strata found in this county are of sedimentary origin, the region must have been below the ancient sea-level to permit their formation, the sediments being carried by streams from an

movements acting as a thrust from the east caused the entire area to be strongly folded, with some faulting, and elevated above the level of the sea, then erosive agents went to work to reduce it. After sufficient time or during the early Tertiary¹ Period the entire eastern United States was reduced to a more or less even plain. The region was again elevated to be followed by erosive action with new vigor. This time the planation was not so complete except in the areas of the less resistant strata, but with mature dissection in the areas of the more resistant strata. The time of this leveling is attributed to late Tertiary. The whole has since been again uplifted and further dissection is now in progress.

The result of these respective influences is the development of similar land forms in regions where like factors have been equally effective. These regions have been divided into physiographic provinces or subdivisions that show similar geologic histories. A map (Figure 2) has been prepared showing the position of Greenbrier County in the physiographic provinces of a portion of the eastern United States.

PHYSIOGRAPHIC PROVINCES.

The eastern United States has been divided into Physiographic provinces by Fenneman² from east to west as follows: (1) Continental Shelf, (2) Coastal plain, (3) Piedmont Province, (4) Blue Ridge Province, (5) Valley and Ridge Province, and (6) Appalachian Plateau.

Portions of the latter four of these divisions are shown on Figure 2 and the boundary between the Appalachian Plateau and Valley and Ridge Province, in Greenbrier County is given in more detail on Figure 3. It will be noted that this boundary in Greenbrier County has been shifted some ten miles farther west than the division line given by Fenneman.³

¹The age of this erosion surface is subject to considerable discussion, but the consensus of opinion now seems to favor early Tertiary for planation and late Tertiary for the uplift.

²Fenneman, N. M. Map. Physical Divisions of the United States, 1916.

³Fenneman, N. M., "Physical Divisions of the United States." Map.

PART I.

History and Physiography.

CHAPTER I.

HISTORICAL AND INDUSTRIAL DEVELOPMENT.

LOCATION.

Greenbrier County, the territory comprising this report, is the second largest county in the State, and is one of the counties bordering on Virginia, situated in the southeastern part of the State. It is included between the parallels of $37^{\circ} 41'$ and $38^{\circ} 16'$ north latitude and the meridians of $79^{\circ} 58'$ and $80^{\circ} 50'$ west longitude from Greenwich. Although it is quite irregular in outline it is roughly pentagonal. A line projecting north and south through its greatest extremity, or a distance of 41 miles, will roughly bisect it. Its greatest width from east to west is 51 miles along a line somewhat north of center. It is bounded on the north by Nicholas, Webster, and Pocahontas Counties, West Virginia; on the east by Bath and Alleghany Counties, Virginia; on the south by Monroe and Summers Counties, West Virginia; and on the west by Summers, Fayette, and Nicholas Counties, West Virginia. More than half the county, on the eastern side, is drained by the Greenbrier River and its tributaries, while the western side is drained by Meadow River and tributaries of the Ganley and Cherry Rivers, all of which go into the Kanawha River and ultimately the Gulf of Mexico.

The geographical position of the county is shown on

general acknowledgment is here made. By way of exception, however, special acknowledgment is due to Mr. J. S. MeWhorter, Mr. G. W. Watts, Mr. J. C. Kennedy, Mr. L. G. Swing, Mr. W. W. Coleman, Mr. J. W. Raine, Mr. B. L. Roberts, Mr. R. B. Holt, Mr. H. H. Blackburn, and Mr. F. W. Tuckwiller whose extraordinary interest in mineral matters and whose wide knowledge of many interesting outcrops and exposures have materially added to the value of the report.

PAUL H. PRICE.

E. T. HECK.

Morgantown, W. Va., December 15, 1938.

ERRATA.

- Page 12, line 2 from top, for Division, read District.
- Page 27, line 17 from bottom, for Costal, read Coastal.
- Page 28, Reverse figure. Top is on binding edge.
- Page 36, line 17 from bottom, for channel, read channels.
- Page 39, first line of table, for source to, read source of.
- Page 42, line 11 of table, for Preaser Branch, read Peaser Branch.
- Page 44, line 13 of table, for Bear Run, read Bear Branch.
- Page 45, line 14 of table, for Old Knob Branch, read Job Knob Branch.
- Page 76, transfer heading to part of table above years 1922-23.
- Page 115, line 12 from top, for subferraneus, read subterraneus.
- Page 116, line 13 from bottom, for bettles, read beetles.
- Page 155, line 11 from top, for basel, read basal.
- Page 188, line 26 from bottom, add Section after Renick Station.
- Page 204, lines 11 and 12 from bottom, for Rensselaria, read Rensselaeria.
- Pages 210 and 211, for Renlck, read Renlck Station.
- Pages 210 and 211, for Renick Valley, read Renicks Valley.
- Page 229, line 14 from bottom, for number, read member.

Relief.—The topography of Greenbrier County is for the most part rugged and mountainous, the causes of which will be discussed in detail under the Chapter on Physiography. Greenbrier River and its tributaries flowing in a southward direction have highly dissected the eastern half of the county. Where resistant rocks were encountered steep precipitous cliffs have been formed. This is particularly true along the banks of the Greenbrier as well as Anthony and Howard Creeks where the latter streams have cut channels transverse to the trend of the mountains. The western side of the county is that of a highly dissected plateau with a general westward drainage of the dendritic type. These streams have cut steep precipitous V-shaped gorges through the more nearly horizontal rocks. The surface varies in elevation from 4372 feet at Grassy Knob at the junction of Old Field Mountain and Cold Knob Mountain in the north central part of the county to 1520 feet along Greenbrier River at a point where this stream leaves the county at the common corner of Greenbrier, Monroe, and Summers Counties one mile west of Alderson, making a total relief of 2852 feet. Other points standing above 4200 feet are: Cold Knob, 4345; Job Knob, 4338; Sugartree Bench, 4276; and Mikes Knob, 4243.

Climate.—From the standpoint of climate that of Greenbrier County, for the most part, is excellent. The winters are neither too long nor severe, and the summers are not unduly warm. July, the warmest month in the year, has an average temperature of 71°, while December and January, the two coldest months, average only 31° and 32°F. The popularity of this area as a summer resort attests to the fact that it is ideal for summer vacationing. Numerous camps for both boys and girls are located along Greenbrier River while many summer homes and cottages are to be found in the vicinity of Lewisburg and White Sulphur Springs.

The following statistics concerning temperature, precipitation, snowfall, and frosts were furnished by United States Weather Bureau, Parkersburg, West Virginia:

Formation.—Greenbrier County, the second largest in the State, was established by act of the Virginia General Assembly, passed January 12, 1778, from parts of Montgomery and Boteourt Counties. Greenbrier is the mother of counties of southern West Virginia as was Monongalia in the northern part of the State. From its original territory Cabell, Kanawha, Mason, Monroe, Nicholas, Webster, Jackson, Wayne, Boone, Putnam, and Roane Counties have been taken.

The county is divided into ten magisterial districts as follows: Anthony Creek, Blue Sulphur, Falling Springs, Fort Springs, Frankford, Irish Corner, Lewisburg, Meadow Bluff, White Sulphur, and Williamsburg. The town of Lewisburg maintained an independent school district until the County Unit bill went into effect.

The county takes its name from the river which flows across it, but just how the river secured its name is still in doubt, although it is generally believed it derived its name from the greenbriers which grow in abundance in the river valley. The county was one of the earliest settled and is rich in historic interest. The present boundaries of Greenbrier County, as carefully surveyed by topographers of the United States Geological Survey, are delineated on Maps I and II. accompanying this report in a separate Atlas.

Area.—The area of Greenbrier County, as determined by planimeter from the topographic maps of the United States Geological Survey, surveyed in cooperation with the West Virginia Geological Survey is as follows:

Districts	Square Miles.
Anthony Creek.....	137.22
Blue Sulphur.....	91.71
Falling Springs.....	180.06
Fort Springs.....	34.38
Frankford.....	51.37
Irish Corner.....	45.53
Lewisburg.....	51.35
Meadow Bluff.....	216.20
White Sulphur.....	22.22

Monthly, Annual, and Mean Precipitation in Inches at Lewisburg (El., 2250').

Year	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Annual
1900.....	2.22	3.44	4.86	1.16	2.43	2.96	4.00	1.63	4.56	4.80	6.60	2.34	41.17
1901.....	3.29	0.85	2.27	5.04	0.00	6.23	3.78	8.50	1.52	0.82	1.23	0.80	46.33
1902.....	8.52	3.20	3.77	2.28	2.38	4.33	0.06	1.08	1.02	1.03	3.09	3.94	38.06
1903.....	3.74	5.86	4.58	3.07	3.19	4.44	5.05	4.30	2.43	2.21	1.30	1.01	42.73
1904.....	1.83	2.37	2.83	2.62	5.23	4.95	2.75	2.65	1.37	0.05	1.38	3.27	31.00
1905.....	3.21	1.24	3.00	2.11	5.76	8.36	6.80	2.80	3.23	2.87	1.72	3.18	39.40
1906.....	5.18	1.21	4.70	2.00	3.30	4.40	4.87	3.76	3.73	2.86	3.08	3.18	42.60
1907.....	3.81	2.77	4.72	4.37	2.66	7.17	3.07	3.38	4.48	1.49	4.56	3.58	46.00
1908.....	3.03	2.22	5.62	4.37	5.45	3.20	6.77	5.09	0.51	3.11	1.96	3.35	44.75
1909.....	3.38	8.10	4.85	3.03	3.77	2.47	2.73	1.63	2.75	3.59	0.65	2.35	35.50
1910.....	3.91	2.00	0.62	3.71	2.29	0.05	4.09	3.77	3.01	1.18	1.53	2.18	35.03
1911.....	0.36	2.05	4.64	4.43	1.30	1.88	2.82	5.06	2.05	5.33	3.02	2.15	42.08
1912.....	2.17	1.41	6.37	4.21	3.43	4.21	3.04	1.21	4.03	1.87	1.70	2.78	37.00
1913.....	3.47	2.71	5.18	3.04	4.14	3.77	6.47	2.07	2.26	4.10	3.20	2.46	42.00
1914.....	3.00	4.71	3.55	3.70	0.96	1.44	4.74	3.26	1.40	3.57	0.99	5.30	37.67
1915.....	4.44	2.94	0.84	2.13	2.88	4.07	3.95	3.65	4.01	4.00	2.17	2.92	38.60
1916.....	3.38	3.57	2.71	3.13	2.43	5.55	2.88	4.74	4.01	1.76	1.58	2.30	38.04
1917.....	2.71	4.07	7.26	2.13	3.88	2.00	3.45	2.95	3.83	2.20	0.75	1.53	36.01
1918.....	4.89	2.13	5.08	5.07	1.06	7.41	4.42	5.81	2.87	4.09	1.50	3.72	50.75
1919.....	4.87	2.35	3.12	2.51	5.02	5.44	7.59	3.30	2.17	4.00	3.74	3.20	48.30
1920.....	3.45	2.44	3.35	5.01	2.50	4.88	3.90	5.08	3.30	0.34	4.70	2.07	41.02
1921.....	1.65	2.76	1.31	2.24	2.36	3.10	3.61	3.25	3.03	3.05	4.18	4.42	30.44
1922.....	2.09	3.40	5.51	3.23	4.28	4.32	3.01	4.07	1.78	2.74	0.01	6.04	42.28
1923.....	3.55	2.51	4.40	3.38	2.16	2.68	2.96	5.74	2.03	0.08	3.13	2.00	37.35
1924.....	4.05	3.69	2.58	2.85	5.05	2.54	3.09	6.03	4.49	1.05	2.51	2.96	40.26
1925.....	3.18	2.17	1.45	3.65	2.04	5.83	2.57	2.36	1.32	5.58	2.23	1.10	32.82
1926.....	3.98	2.77	2.48	2.18	1.85	1.27	3.13	0.01	2.00	7.40	2.10	6.26	42.18
1927.....	1.53	5.82	2.75	6.04	2.63	3.61	3.75	4.02	3.26	3.18	3.17	3.71	44.07
1928.....	1.79	1.57	2.56	3.10	2.61	5.45	4.11	5.98	3.74	2.03	2.00	1.35	35.78
1929.....	2.77	3.55	3.91	4.03	4.41	3.07	2.71	3.06	0.70	6.51	4.71	2.37	42.43
1930.....	1.63	2.25	1.04	1.65	1.94	1.17	3.01	1.09	0.30	0.05	1.61	2.18	18.88
1931.....	3.72	1.45	3.66	0.06	6.97
1932.....	1.24	2.25	2.18	4.15	5.53	3.02	1.26	4.63	4.18	2.31
1933.....	3.52	4.10	3.80	2.80	5.72	1.87	7.26	4.16	1.30	0.78	1.35	2.54	39.20
1934.....	2.80	2.03	2.63	1.50	2.02	3.45	4.24	4.06	2.37
1935.....	9.57	5.70
1936.....
Mean.....	3.44	2.83	3.73	3.33	3.38	3.83	4.00	3.92	2.69	2.00	2.47	3.10	39.70

**Monthly, Annual, and Mean Temperature in Degrees
Fahrenheit at White Sulphur Springs (El., 1914').**

Year	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Annual
1895.....	68.7	69.0	72.6	68.7	46.0
1896.....	30.6	33.4	37.8	50.8	66.4	68.0	71.6	71.5	64.2	56.6	47.0	33.8	52.6
1897.....	29.6	36.2	48.1	50.5	64.7	68.2	75.9	72.6	63.0
1898.....	51.2	47.2	63.0	66.3	55.0	43.2
.....
1915.....	64.4	68.4	67.6	65.0	55.4	44.2	31.0
1916.....	33.3	34.2	39.9	49.6	62.6	63.4	69.7	69.6	62.1	54.6	41.3	32.0	51.4
1917.....	36.2	32.7	44.0	52.9	53.6	66.8	70.0	69.5	62.0	49.7	40.5	24.8	50.3
1918.....	21.6	36.9	47.3	50.2	64.8	69.0	73.6	61.1	57.6
1919.....
1920.....	70.4	65.2	55.0	40.0	32.4
1921.....	31.9	36.6	55.4	54.0	60.0	70.5	74.5	69.6	69.6	53.2	46.3	37.6	54.9
1922.....	30.0	37.8	45.6	50.4	63.0	70.2	70.8	67.1	64.6	53.2	42.7	36.4	52.6
1923.....	33.3	31.2	41.9	51.0	66.2	70.0	71.0	70.5	66.1	56.1	40.5	39.2	52.1
1924.....	26.6	28.2	35.8	49.6	55.0	69.4	60.2	71.6	59.2	52.2	42.8	33.8	49.5
1925.....	30.4	46.8	43.0	55.8	58.4	73.0	72.8	76.8	70.2	48.2	39.0	31.8	52.8
1926.....	29.4	35.5	30.4	48.0	62.0	66.2	72.0	73.9	68.8	54.0	38.6	33.0	51.7
1927.....	33.6	42.6	45.5	52.2	63.4	67.7	73.4	68.3	67.8	50.2	46.2	33.8	54.1
1928.....	32.6	34.9	41.6	50.0	59.8	67.5	72.8	73.2	58.5	52.6
1929.....	36.7	30.1	46.4	54.2	60.6	66.0	76.4	67.1	63.1	48.8	40.2	34.4	51.0
1930.....	30.6	36.0	36.2	52.1	61.8	67.1	72.6	68.0	66.3	40.6	37.4	26.8	56.5
1931.....	30.3	33.2	35.0	50.4	59.0	68.6	75.6	69.6	67.6	54.6	47.4	37.4	52.3
1932.....	40.6	39.2	30.2	56.4	60.4	68.0	71.6	68.8	63.0	51.6	38.6	31.1	51.5
1933.....	36.6	31.4	38.6	50.0	64.4	68.2	76.6	68.9	67.0	50.4	38.2	36.3	51.7
1934.....	34.1	27.9	41.8	52.8	66.9	75.2	77.5	72.6	67.2	55.8	48.8	31.6	54.3
1935.....	30.8	32.8	48.4	50.6	60.1	67.6	74.0	72.6	64.4	54.0	45.5	26.9	52.3
1936.....	20.6	30.2	45.0	49.5	62.3	69.4	73.6	73.0	66.2	56.1	40.6	33.0	52.1
Mean.....	31.6	34.4	42.8	51.3	61.0	68.4	72.0	76.5	65.1	52.7	42.3	32.9	52.1

Monthly, Annual, and Mean Temperature in Degrees Fahrenheit at Lewisburg (El., 2250').

Year	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Annual
1900.....	60.8	70.2	72.7	73.8	60.9	58.8	45.2	34.4
1901.....	32.4	26.6	41.5	45.8	59.6	68.5	75.0	70.0	62.4	51.8	35.4	30.8	50.0
1902.....	28.8	20.2	41.3	47.4	63.2	65.4	72.0	60.0	62.0	54.8	48.8	32.9	51.0
1903.....	29.0	34.8	49.2	48.8	61.5	63.0	71.1	70.6	63.1	52.4	36.8	26.4	50.0
1904.....	28.4	28.5	43.2	47.1	60.2	68.2	69.4	66.6	65.0	51.8	37.6	32.2	50.1
1905.....	27.8	25.8	40.6	52.0	64.4	69.3	71.8	68.8	64.2	52.0	41.0	36.2	51.0
1906.....	37.6	34.7	37.8	53.0	60.6	69.2	70.3	74.2	68.2	53.0	43.8	35.2	53.1
1907.....	40.2	30.8	47.3	44.2	58.4	63.8	71.6	60.0	60.0	40.2	40.7	33.0	51.2
1908.....	28.3	28.0	48.1	54.5	61.0	67.6	72.5	69.4	62.6	52.8	43.0	34.2	51.9
1909.....	35.8	30.8	40.4	52.4	60.0	71.0	69.0	69.2	62.2	40.0	47.4	29.0	52.1
1910.....	31.8	31.2	48.8	52.2	56.0	64.8	70.9	67.8	66.8	36.0	26.3
1911.....	36.4	38.8	39.9	47.0	63.4	68.0	70.6	72.0	67.8	56.1	30.0	37.8	53.2
1912.....	23.0	28.0	39.0	54.2	61.2	65.0	71.2	67.7	66.8	54.1	40.3	35.2	50.7
1913.....	30.4	33.8	44.2	50.4	61.8	68.2	72.4	69.0	61.0	53.0	43.5	37.0	52.0
1914.....	34.2	20.6	35.8	51.2	60.5	71.4	70.0	60.8	61.2	55.0	40.8	29.1	50.8
1915.....	30.2	37.7	33.4	52.8	60.0	65.3	60.5	68.0	65.2	55.4	43.0	30.8	51.0
1916.....	38.4	33.2	38.2	49.0	62.7	65.0	72.0	71.0	61.4	52.6	42.0	31.7	51.4
1917.....	34.6	32.2	42.4	51.8	54.2	65.5	70.4	69.2	62.0	47.4	39.1	22.1	49.2
1918.....	19.6	34.0	40.8	48.7	64.4	65.8	67.4	72.2	58.8	56.0	41.4	38.8	51.3
1919.....	31.4	34.0	44.0	50.8	60.4	60.0	71.7	67.8	63.2	61.2	44.1	31.0	52.4
1920.....	20.8	30.4	40.6	48.8	57.3	64.8	67.8	69.2	65.0	54.1	41.2	33.6	50.2
1921.....	33.7	30.8	52.2	53.2	58.4	69.8	73.1	68.0	60.5	52.2	45.0	36.4	54.1
1922.....	29.3	37.4	44.6	53.0	61.6	68.8	71.7	60.8	65.2	54.0	41.0	36.9	52.7
1923.....	34.4	31.8	41.8	49.4	59.0	68.5	69.4	69.8	65.0	50.4	40.6	42.4	51.9
1924.....	29.8	31.4	38.4	40.7	55.0	67.0	68.3	69.5	59.3	52.4	42.8	34.6	40.9
1925.....	31.3	41.4	44.5	54.0	56.0	69.6	70.6	67.8	69.0	49.0	39.8	33.0	52.2
1926.....	30.2	35.8	30.2	47.8	60.0	65.6	71.2	72.6	68.2	53.8	39.2	33.8	51.2
1927.....	32.6	41.7	44.4	51.6	60.3	64.5	69.4	65.0	64.1	54.8	46.0	34.2	52.4
1928.....	33.2	35.2	42.0	48.0	57.4	64.5	70.4	71.3	59.2	54.0	42.2	34.8	51.1
1929.....	32.1	30.2	45.7	54.5	59.4	65.5	69.8	65.0	64.2	51.2	43.0	35.5	51.5
1930.....	34.0	30.4	40.6	52.2	62.9	60.8	72.6	68.0	68.6	50.2	40.8	29.6	52.2
1931.....	69.8	71.1	75.2	54.9	50.8
1932.....	44.6	41.7	36.6	49.6	60.5	67.4	71.4	65.5	53.6	40.8	34.4
1933.....	38.1	33.8	40.3	51.3	64.2	68.9	70.2	69.4	69.3	54.4	41.1	40.0	53.4
1934.....	36.0	26.6	40.0	53.2	63.9	74.2	78.3	71.2	66.4	54.3
1935.....	47.4	62.0
1936.....
Mean.....	32.7	33.1	42.5	50.6	60.3	67.5	71.2	69.5	64.7	53.3	41.9	33.4	51.6

Monthly, Annual, and Mean Precipitation in Inches at White Sulphur Springs (El., 1914').

Year	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Annual
1866.....	1.64	5.04	3.37	1.65	1.05
1896.....	2.76	2.13	5.68	5.36	3.86	5.15	6.60	5.07
1897.....	6.96	7.07	4.91	1.77	5.86	6.55	6.56	5.93
1898.....	3.67	4.56	1.86	5.76	2.71
1916.....	4.92	3.21	2.86	3.86	3.28	1.93	4.43
1916.....	2.73	3.38	2.61	2.76	2.47	4.77	3.83	4.96	2.47	2.64	1.65	2.38	36.06
1917.....	3.67	3.86	7.55	2.55	4.15	1.70	6.66	2.65	4.40	3.85	6.36	3.32	48.39
1918.....	3.60	2.57	5.46	5.66	3.10	7.25	4.86	2.65	2.46	4.55	1.56	3.56	47.62
1919.....
1926.....	8.63	3.32	6.56	3.44	2.15
1921.....	2.88	2.55	0.83	3.56	3.29	2.22	4.09	4.61	7.02	2.58	2.53	2.55	38.66
1922.....	1.64	2.69	3.65	1.53	3.34	4.88	4.16	2.66	1.78	1.76	6.52	6.52	34.46
1923.....	4.82	3.78	3.13	2.69	2.64	2.83	2.96	3.16	4.76	1.23	2.66	1.30	35.66
1924.....	6.62	1.86	2.62	2.69	5.94	4.66	3.45	7.37	4.93	1.25	1.24	3.63	39.66
1925.....	3.76	1.56	1.54	2.76	2.26	5.26	1.61	6.91	1.62	3.78	2.86	3.42	31.54
1926.....	5.52	1.48	2.47	1.67	2.86	6.66	3.26	5.61	1.71	6.59	2.86	5.45	39.28
1927.....	1.72	6.23	1.21	5.55	2.26	3.58	3.44	2.62	2.13	3.70	3.27	3.16	36.22
1928.....	2.12	1.15	2.51	3.67	1.99	4.36	3.46	4.54	6.15	2.65	1.96	1.55	34.76
1929.....	2.66	1.97	3.76	3.85	3.76	3.26	2.56	2.56	6.86	5.26	3.96	1.33	34.75
1930.....	1.37	1.55	1.26	1.76	2.25	6.96	1.36	1.26	6.46	6.86	1.71	1.45	15.88
1931.....	0.66	2.66	2.65	3.15	5.35	1.55	4.89	4.25	3.15	1.16	1.25	4.45	33.55
1932.....	3.20	3.35	4.35	3.69	3.15	4.95	2.76	2.06	2.66	4.10	3.26	2.88	36.87
1933.....	3.16	4.55	4.46	3.17	4.85	1.15	5.86	2.86	1.65	1.65	1.45	2.45	35.82
1934.....	1.85	3.06	6.26	2.45	1.75	2.56	5.15	4.66	5.16	2.16	5.25	2.87	42.82
1935.....	5.62	2.16	8.25	3.46	3.95	4.45	8.46	3.25	4.20	1.25	4.50	2.79	51.61
1936.....	5.55	4.66	5.35	2.75	1.25	2.50	3.65	2.41	1.86	2.86	0.75	2.66	34.81
†Mean....	2.92	2.79	3.63	3.11	3.15	3.31	3.94	3.43	3.68	2.65	2.32	2.99	37.36

*Partly interpolated.

†1916-1918 and 1921-1936, inclusive.

Assessed Value of Real Estate, Personal Property and Public Utility Property in Greenbrier County,
Taxes Levied and the Total Average Rate of Levy, for the Years 1929 to 1936, Inclusive.

Year	Assessed Value of Property				Taxes Levied			Total Average Rate of Levy on the \$100 Value- tion
	Real Estate	Personal Property	Public Utility Property	Total	On Real Estate	On Personal Property	On Public Utility Property	On Total
1929.....	\$10,575,280	\$5,230,300	\$ 7,845,092	\$29,659,672	\$521,816	\$175,435	\$245,895	\$943,145
1930.....	16,546,560	5,004,705	9,732,345	31,283,670	508,031	163,574	290,901	962,506
1931.....	8,031,910	4,110,615	9,718,115	22,466,640	331,290	105,298	326,200	822,854
1932.....	8,514,910	3,260,470	9,701,397	21,476,777	318,417	120,271	305,531	750,210
1933.....	8,522,570	4,132,010	10,121,700	22,777,180	149,804	01,204	203,562	414,720
1934.....	8,513,605	4,010,502	10,134,000	22,664,167	150,550	00,770	202,077	413,409
1935.....	8,531,590	4,133,100	10,189,600	22,854,290	143,488	59,428	196,218	399,134
1936.....	8,320,155	4,330,685	10,147,300	22,798,440	135,500	62,322	190,354	388,245

	Lewisburg (El., 2250').		White Sulphur Springs (El., 1914')	
Year	Last in Spring	First in Autumn	Last in Spring	First in Autumn
1931	Sept. 30	May 9	Oct. 17
1932	May 3	Sept. 25	May 29	Oct. 6
1933	April 27	Oct. 26	April 28	Oct. 19
1934	April 27	Oct. 13	April 28	Oct. 14
1935	April 17	Sept. 30
1936	Oct. 28	May 14	Oct. 28
Average	April 29	Oct. 6	May 6	Oct. 14

Products.—Greenbrier County is fortunate in being able to boast of a diversified list of products of natural wealth. It may justly claim to be a coal mining, a lumbering, an agricultural, or a resort and mineral spring county. Few other counties in the State can offer so great a variety.

A broad limestone belt crossing the center of the county from north to south furnishes excellent agricultural land. The western side of the county produced over 2,000,000 tons of coal during 1930 with a value of over \$3,000,000. Valuable growths of timber, both hard and soft woods, are found throughout the county with large acreages on both the east and west sides. The county is the most popular resort area in the State, both for resort hotels and summer camps in conjunction with several valuable mineral springs. Numerous limestone quarries are found throughout the limestone belt that can furnish limestone for any purpose.

The products will be treated in more detail under their respective headings.

The principal crops in Greenbrier in order of their importance are hay, corn, wheat, oats, Irish potatoes, barley, and buckwheat.

The principal animal products in order of their importance are cattle, sheep, horses, hogs, chickens, and turkeys. Dairy products (included under cattle) are a very important resource in this county.

Property Valuation.—According to the State Tax Com-

Official Postal Guide for 1936, shows the number of post-offices in the county. The figures following the letters R, P, and S after an office indicate the following, as of April 1, 1936: R, boxes on rural routes emanating therefrom; P, post-office boxes at offices not having city letter-carrier service; S, boxes on star routes emanating therefrom.

Alvon S14	Dawson S14	Organ Cave S15
Aujan P72	Duo	Quinwood P109, S78
Anthony	East Ralmonde P138	Ralmonde P159, S78
Asbury S7	Esty S43	Renick P29, S215
Auto S29	Fort Spring P22	Renicks Valley S12
Bellburn P4	Frankford P34, S52	Richlands S23
Bingham S3	Friars Hill S10	Ronceverte R156, P201, S5
Blaker Mills S7	Grassy Meadows S7	Rupert S10
Blue Sulphur Springs	Hughart S26	Russellville
R65, S8	Kletter S14	Smoot S28
Caldwell P15, S110	Law S1	Spring Creek S15
Camp Alleghany	Leonard S18	Sunlight S6
Charmco P7	Leslie P52	Trout S11
Clearco	Lewisburg R175, P335, S97	Unus S100
Climonville S48	Marrance P32	Vargo S35
Cordova S27	Maxwellton P3, S24	Vale
Cornstalk S11	Meadow Bluff S20	White Sulphur Springs
Crag	Neola S41	P213, S54
Crawley S20	Nutterville S22	Williamsburg P30, S19
Crichton P6		
Alderson, R. O. in Monroe County, R125, P250, S100.		

Population.—The following table, taken from the United States Census Returns for 1930, shows the population of Greenbrier County by districts for the last three enumerations:

Anthony Creek Division.....	1,050	1,224	1,164
Blue Sulphur District.....	3,339	3,871	3,382
Alderson town, total.....	1,458	1,401	1,252
In Greenbrier County.....	930	841	677
In Monroe County.....	528	560	575
Falling Springs District.....	2,735	2,752	5,689
Falling Springs town.....	355	263	270
Fort Springs District.....	3,720	3,585	3,443
Ronceverte city.....	2,254	2,319	2,157
Frankford District.....	2,078	1,966
Frankford town.....	140	110	102
Irish Corner District.....	2,420	1,947	1,846
Lewisburg District.....	3,022	2,403	2,558
Lewisburg town.....	1,293	1,202	803
Meadow Bluff District.....	11,540	3,928	2,688
East Rainelle town*.....	1,272
Marfrance town*.....	1,066
Rainelle town.....	920	566
White Sulphur District.....	3,693	2,837	1,609
White Sulphur Springs town.....	1,484	837	338
Williamsburg District.....	2,272	2,229	2,454
Williamsburg town.....	148	161	120
Totals for county.....	35,878	26,242	24,833

*East Rainelle town incorporated in 1921 and Marfrance town in 1926.

TOWNS AND INDUSTRIES.

LEWISBURG.

Lewisburg, the county-seat, stands on the site of old Fort Union, and was named in honor of General Andrew Lewis who was active in military operations in this vicinity in 1774. The town was established in October, 1782, but its settlement dates from the gathering of the frontier army in 1774. The town is located in a topographic depression, or limestone "sink" in the southeastern part of the county at the junction of two old and historic trails, namely, the Midland Trail or James River and Kanawha Turnpike (U. S. Route 60) traversing east and west, and the Seneca Trail (U. S. Route 219) traversing north and south. It is located in the midst of a beautiful and valuable agricultural community and is noted for its schools and fine homes. There is a large General Store, a Hotel, a

College for Women which dates its history back to 1812, and is the successor of Lewisburg Seminary, Lewisburg Female Institute, and Lewisburg Academy. The town is well supplied with churches and is represented by the Methodist, Presbyterian, and Episcopal as well as a church for the colored inhabitants. Because of its institutions of learning and churches Lewisburg had always been known as a town of culture. The United States bench mark at Lewisburg is 2084 feet above sea-level. Its population according to the 1930 Census was 1293.

RONCEVERTE.

Roneeverte was laid out in 1871 by Colonel Ceeil Clay and incorporated in 1882. It was given the name of the French equivalent of Greenbrier (**Ronce**—brier, **-vert**—green), the river along which it is situated. The town was a result of the construction of the Chesapeake and Ohio Railway. Its growth was largely determined by its timber industries, its convenient access to an excellent agricultural region and its location at the junction of the Greenbrier Division of the Chesapeake and Ohio with the main line of the railroad. It is now the largest town in Greenbrier County.

The town has two banks, with capital stock and resources of \$1,500,000; one weekly newspaper; one theater; an armory; a concentration depot for receiving milk and cream from the neighboring dairies; and a large steam generating power unit of the Virginia Public Service Company.

The town is well supplied with elementary and high schools, as well as with nine churches.

The water system and filtration plant are municipally owned and operated.

A United States Government bench mark at Roneeverte is 1665 feet above sea-level. The population of Roneeverte according to the 1930 Census was 2254.

WHITE SULPHUR SPRINGS.

The town of White Sulphur Springs is located in the southeast part of the county in a wide valley cut by Howard

east of Charleston on the Midland Trail (U. S. Route 60) and is served by the main line of the Chesapeake and Ohio Railway. The land upon which it is situated was originally patented to Nathan Carpenter, who built his cabin near the spring and removed his family to it in 1774. It was incorporated in 1910. The town is built largely around the famous White Sulphur Springs resort which furnishes employment for a large number of the inhabitants. Aside from numerous hotels and tourists' houses within the corporate limits numerous excellent summer homes are located within a few miles radius of the town. The Government has established one of its Federal Fish Hatcheries here. It is also local headquarters for the Labor Nursery that does a large business in West Virginia evergreens.

In 1930 the town had a population of 1484. Its elevation is 1917 feet above sea-level.

ALDERSON.

The town of Alderson is located along the Greenbrier River, being partly in Monroe and partly in Greenbrier, near their common corner with Summers County. The town as originally incorporated in 1890 included only that part situated in Monroe County, but in 1902 the charter was amended to include that portion of the town lying in Greenbrier County. The principal business section is on the Monroe side while its main residential section is on the Greenbrier side with the latter county having the largest number of inhabitants, 930 of a total of 1458, according to the 1930 Census.

The town is served by the main line of the Chesapeake and Ohio Railway. It is situated upon the flood-plain and terraces of the Greenbrier River with an elevation of 1555 feet above sea-level.

The Alderson Academy, a Baptist school, is located here but has recently (1932) been consolidated with Broadus College at Philippi and will be removed to that place.

A Federal Industrial Institution for women is maintained on the Monroe side of the river. One of the several excellent summer camps (Camp Greenbrier) that are located along the Greenbrier River is located on the Greenbrier County side.

consists of supplying the needs of the rich farming community that surrounds the town.

RAINELLE.

Rainelle is located on a broad terrace near the junction of Sewell Creek and Meadow River in the western side of Greenbrier County. It is primarily a lumber town and is situated in the midst of one of the finest hardwood lumber tracts in the State. The town was incorporated April 25, 1913, and was named in honor of John and T. W. Raine, pioneer lumber, railroad, and coal mine operators in this area. The town is almost entirely made up of employees of the Meadow River Lumber Company which boasts the finest and largest hardwood lumber mill in the country.

The town is supplied with well-equipped hotels, banks, schools, and churches. The growth of the town was simultaneous and in conjunction with the building of the Sewell Valley Railroad (now owned by the Chesapeake and Ohio) and the Meadow River Lumber Company plant.

The Midland Trail (U. S. Route 60) passes through Rainelle. The population of the town in 1930 was 920. The elevation near the center of the town is 2425 feet above sea-level.

EAST RAINELLE.

East Rainelle, formerly Sewell Valley, and separated from Rainelle proper only by Sewell Creek, was incorporated under its own charter in 1921. The town is made up largely of small business establishments along the Midland Trail, which bisects the town, that serve the immediate town and surrounding area.

In 1930 the town had a population of 1272.

MARFRANCE.

Marfrance, a coal-mining town, is located on the headwaters of Meadow Creek, a tributary of Meadow River on the western side of the county.

General Account and Section, Pottsville Series.....	214-217
General Section, Pottsville Series.....	215-217
Topographic Expression.....	218
Contacts and Unconformities.....	218-219
Fossil Life.....	219
Correlation, Pottsville Series.....	219-220
Description of Members, Kanawha Group.....	220-222
Lower Gilbert Sandstone.....	221
Gilbert "A" Coal.....	221
Gilbert Shale.....	221
Gilbert Coal.....	221
Dotson Sandstone.....	221
Douglas "A" Coal and Douglas Coal.....	221
Lower Dotson Sandstone.....	222
Douglas Shale.....	222
Description of Members, New River Group.....	222-242
Upper Nuttall Sandstone.....	223
Jaeger "B" Coal.....	223-224
Lower Nuttall Sandstone.....	224
Jaeger "A" Coal.....	224
Upper Jaeger Shale.....	224
Hughes Ferry Coal.....	225
Middle Jaeger Sandstone.....	225
Lower Jaeger Coal.....	225
Lower Jaeger Sandstone.....	226
Lower Jaeger Shale.....	226
Harvey (Conglomerate) Sandstone.....	226
Sandy Huff Shale.....	226-227
Castle Coal.....	227
Gnyandot Sandstone.....	227
Skelt Shale.....	227
Sewell "B" Coal.....	228
Sewell "A" Coal.....	228
Lower Guyandot Sandstone.....	229
Hartridge Black Shale.....	229
Sewell Coal.....	229-234
Erratic Boulders in the Sewell Coal.....	230-234
Welch Sandstone.....	234
Welch Coal.....	234
Upper Raleigh Sandstone.....	234-235
Little Raleigh "A" Coal.....	235
Little Raleigh Coal.....	235-236
Lower Raleigh Sandstone.....	236
Beekley "Rider" Coal.....	236
Beekley Coal.....	236-237
Quinnimont Sandstone.....	237-238
Quinnimont Shale.....	238
Fire Creek Coal.....	238
Little Fire Creek Coal.....	239-241
Pineville Sandstone.....	241
No. 9 Pocahontas Coal.....	241
No. 8 Pocahontas Coal.....	241-242
Description of Members, Pocahontas Group.....	242-250
Flattop Mountain Sandstone.....	242-243
Rift Shale, No. 7 Pocahontas Coal, Plerpont Sandstone,.....	242-245

Pocono Series.....	283-294
General Account and Section.....	283
General Section.....	283
Topographic Expression.....	284
Areal Extent.....	284-285
Contacts	286
Fossil Life.....	286-289
Correlation	289
Description of Members.....	290-294
Merrimac Coal.....	290-292
Broad Ford Sandstone.....	292-293
Berca Sandstone.....	293-294
Economic Aspects.....	294
Chapter VIII.—Stratigraphy—Devonian Rocks.....	295-326
General Statement.....	295-296
Upper Devonian Rocks.....	296-309
Catskill Series.....	296-300
General Account.....	296-297
Topographic Expression.....	297
Areal Extent.....	297-298
Contacts	299
Fossil Life.....	300
Correlation	300
Economic Aspects.....	300
Chemung Series.....	300-304
General Account and Section.....	300-301
General Section.....	301
Topographic Expression.....	300-301
Areal Extent.....	302
Contacts	302
Fossil Life.....	303
Correlation	303
Description of Members.....	303
Hendricks Sandstone.....	303
Economic Aspects.....	304
Portage Series.....	304-307
General Account.....	304-305
Topographic Expression.....	305
Areal Extent.....	306
Contacts	306
Fossil Life.....	306
Correlation	306-307
Economic Aspects.....	307
Genesee Series.....	307-309
General Account.....	307-308
Topographic Expression.....	308
Areal Extent.....	308
Contacts	308
Fossil Life.....	309
Correlation	309
Description of Members.....	309
Economic Aspects.....	309
Middle Devonian Rocks.....	310-313
General Statement.....	310
Marcelius Series.....	310-313
General Account.....	310-311

Upper Pocahontas Sandstone.....	248
No. 3 Pocahontas "Rider" Coal.....	248
No. 3 Pocahontas Coal.....	249
Lower Pocahontas Sandstone.....	249
No. 2 "A" Pocahontas Coal.....	249
No. 2 Pocahontas Coal.....	250
No. 1 Pocahontas Coal.....	250
Economic Aspects, Pottsville Series.....	251

Chapter VII.—Stratigraphy—Mississippian Rocks..... 252-294

General Statement.....	252
Correlation, Mississippian Period.....	252-254
Mauch Chunk Series.....	254-255
General Account and Section.....	254-255
General Section.....	255
Topographic Expression.....	256
Areal Extent.....	256-257
Contacts	258
Fossil Life.....	258
Correlation	258-259
Description of Members, Bluestone Group.....	259-260
Description of Members, Princeton Group.....	260
Princeton Sandstone.....	260
Description of Members, Hinton Group.....	260-262
Avis Limestone.....	261
Stony Gap Sandstone.....	261-262
Description of Members, Bluefield Group.....	262-265
Droop Sandstone.....	263
Talcott and Ada Shales.....	263
Reynolds Limestone.....	264
Webster Springs Sandstone.....	264
Glenray Limestone.....	264-265
Lillydale Shale.....	265
Economic Aspects.....	265
Greenbrier Series.....	266-279
General Account and Section.....	266-267
General Section.....	267
Topographic Expression.....	267-268
Areal Extent.....	268-269
Contacts	270
Fossil Life.....	270
Correlation	270
Description of Members.....	271-279
Alderson Limestone.....	271
Greenville Shale.....	271-272
Union Limestone.....	272
Pickaway Limestone.....	272-277
Taggard Limestone.....	277-278
Fulton Limestone.....	278
Sinks Grove Limestone.....	278-279
Hillsdale Limestone.....	279
Economic Aspects.....	279
Maccrady Series.....	280-282
General Account.....	280
Topographic Expression.....	280
Areal Extent.....	281

Contacts	337
Fossil Life.....	337
Correlation	338-339
Description of Members.....	338
Upper Shales.....	338
Keefer Sandstone.....	338
Shales and Thin Limestones.....	338
Fossil Ore Horizon.....	338
Middle Shales.....	339
Iron Sandstone.....	339
Economic Aspects.....	339-341
White Medina Series.....	339
General Account.....	339-340
Topographic Expression.....	340
Areal Extent.....	340
Contacts	340
Fossil Life.....	341
Correlation	341
Economic Aspects.....	341-343
Red Medina Series.....	341-342
General Account.....	342
Topographic Expression.....	342
Areal Extent.....	342
Contacts	342
Fossil Life.....	342-343
Correlation	343
Economic Aspects.....	

PART III. MINERAL RESOURCES.

Chapter X.—Petroleum and Natural Gas.....	344-362
General Statement.....	344-346
Prospective Oil and Gas Areas.....	346-347
Prospective Oil and Gas Horizons.....	347-350
Table Showing the Estimated Depths to Geologic Horizons at Various Points.....	349
Mineralogy of Lower Devonian and Upper Silurian Sandstones	350
Well Records.....	351-362
Summary of Oil and Gas Possibilities.....	362
Chapter XI.—Commercial Coal.....	363-617
Introduction	363-365
Diagram Showing the Position and Thickness of Coal Seams.....	364
Statistics of Coal Production.....	365-384
Records of Coal Test Borings.....	385-472
Summarized Records.....	385-389
Summarized Record of Coal Test Borings (Table)	386-389
Detailed Coal Test Borings:	
Meadow Bluff District.....	390-406
Williamsburg District.....	406-407
Irish Corner District.....	407-409
Nicholas County.....	409-434
Fayette County.....	434-472

	Page.
Topographic Expression.....	311
Areal Extent.....	311
Contacts	311-312
Fossil Life.....	312
Correlation	312
Description of Members.....	312-313
Economic Aspects.....	313
Lower Devonian Rocks.....	313-326
General Statement.....	313-314
Oriskany Series.....	314-321
General Account and Section.....	314-315
General Section.....	315
Topographic Expression.....	315
Areal Extent.....	315-316
Contacts	317
Fossil Life.....	317
Correlation	317-318
Description of Members.....	318-320
Huntersville Chert.....	318-319
Ridgeley Sandstone.....	320
Economic Aspects.....	320-321
Helderberg Series.....	321-326
General Account.....	321
Topographic Expression.....	321-322
Areal Extent.....	322
Contacts	322
Fossil Life.....	322
Correlation	323
Description of Members.....	323-325
Beerft Member.....	323
New Scotland Member.....	324
Coeymans Member.....	325
Keyser Member.....	325
Economic Aspects.....	326
Chapter IX.—Stratigraphy—Silurian Rocks.....	327-343
General Statement.....	327-329
Sallina Series.....	329-332
General Account.....	329
Topographic Expression.....	330
Areal Extent.....	330
Contacts	330
Fossil Life.....	330
Correlation	330-331
Description of Groups.....	331-332
Bossardville Limestone Group.....	331
Rondout Waterlime Group.....	331-332
Economic Aspects.....	332
Niagara Series.....	332-334
General Account.....	332-333
Topographic Expression.....	333
Areal Extent.....	333
Contacts	333
Fossil Life.....	333
Correlation	334
Description of Members.....	334
Economic Aspects.....	334

Mineable Coals, New River Group of Pottsville Series.....	473-573
Sewell Coal.....	473-529
Meadow Bluff District.....	473-520
Williamsburg District.....	520-526
Falling Springs District.....	526-528
Quantity Available.....	529
Little Raleigh Coal.....	530-539
Meadow Bluff District.....	532-537
Williamsburg District.....	537-538
Quantity Available.....	538-539
Beckley Coal.....	539-553
Meadow Bluff District.....	530-550
Williamsburg District.....	551-552
Falling Springs District.....	552-553
Quantity Available.....	553
Fire Creek Coal.....	554-573
Meadow Bluff District.....	556-570
Williamsburg District.....	570-571
Quantity Available.....	572-573
Mineable Coals of the Pocahontas Group of Pottsville Series.....	573-607
No. 7 Pocahontas Coal.....	573-579
Meadow Bluff District.....	573-579
No. 6 Pocahontas Coal.....	580-600
Meadow Bluff District.....	582-599
Williamsburg and Falling Springs Districts.....	599-600
Quantity Available.....	600
No. 3 Pocahontas Coal.....	601-607
Meadow Bluff District.....	603-606
Williamsburg District.....	606
Quantity Available.....	607
Summary of Available Coal.....	607-608
Summary of Available Coal by Districts (Table).....	608
Table of Coal Analyses.....	608-617
Table of Coal Analyses.....	611-617

Chapter XII.—Limestone, Road Material, Clay, Building Stone,

Glass-Sand, Forests, and Soils.....	618-652
Limestone	618-634
General Statement.....	618-619
Limestones of the Mississippian Period.....	619-629
Limestones of the Mauch Chunk Series.....	619-620
Avis Limestone.....	619-620
Reynolds Limestone.....	620
Glenray Limestone.....	620
Limestones of the Greenbrier Series.....	620-629
Alderson Limestone.....	621-622
Union Limestone.....	622-626
Fickaway Limestone.....	626
Taggard Limestone.....	626
Patton Limestone.....	626-627
Sinks Grove Limestone.....	627-628
Hillsdale Limestone.....	628-629
Limestones of the Devonian and Silurian Periods.....	629-630
General Statement.....	629
Helderberg Limestone.....	629
Salina Series.....	629-630
Niagara Series.....	630

Clay	636-640
General Statement.....	636
Available Clay and Shale.....	637-640
Residual Clay.....	637
Transported Clay and Consolidated Clay or Shale	637-640
Fire Clay.....	640
Glass-Sand	640-642
Table of Sandstone Analyses.....	642
Forosts	643-647
Original Forest Conditions.....	643-644
The Lumber Industry.....	644-646
Present Forest Conditions.....	646
Monongahela National Forest.....	646-647
Lumber Mills.....	647
Soils of Greenbrier County (by Anton J. Vessel).....	648-652
Chapter XIII.—Mineral Waters, Water-Power, Iron Ore,	
Manganese, and Precious Metals.....	653-678
Mineral Waters.....	653-660
General Statement.....	653
Mineral Springs.....	654-660
Water-Power	660-664
Present Development.....	660
Greenbrier River.....	661
Anthony Creek.....	661
Howard Creek.....	662
Muddy Creek.....	662
Meadow River.....	662
Indicated Horse-Power of Streams.....	662-664
Iron Ore.....	665-671
General Statement.....	665
Oriskany Iron Ore Prospects and Exposures.....	665-670
Clinton Ores.....	670
Table of Iron Ore Analyses.....	670-671
Manganese	672-677
General Statement.....	672-674
Manganese Prospects.....	674-676
Table of Manganese Ore Analyses.....	676-677
Precious Metals.....	678

PART IV. PALEONTOLOGY.

Chapter XIV.—Notes on Paleontology of Greenbrier County	
(b by John L. Tilton and Dana Wells).....	679-734
Fossil Collections from Greenbrier County, W. Va.	
(Register of Localities by Collection Nos.).....	680-692
Distribution of Collections by Geologic Formations.....	692-693
Pennsylvanian	693-694
Pottsville Series.....	693-694
New River Group.....	693-694
Pocahontas Group.....	694
Mississippian	695-714
Mauch Chunk Series.....	695-703
Bluestone Group.....	695-699

Road Material.....	635-636
Building Stone.....	636-640
Clay	636
General Statement.....	637-640
Available Clay and Shale.....	637
Residual Clay.....	637-640
Transported Clay and Consolidated Clay or Shale	640
Fire Clay.....	640-642
Glass-Sand	642
Table of Sandstone Analyses.....	643-647
Forests	643-644
Original Forest Conditions.....	644-646
The Lumber Industry.....	646
Present Forest Conditions.....	646-647
Monongahela National Forest.....	647
Lumber Mills.....	648-652
Soils of Greenbrier County (by Anton J. Vessel).....	
Chapter XIII.—Mineral Waters, Water-Power, Iron Ore, Manganese, and Precious Metals.....	653-678
Mineral Waters.....	653
General Statement.....	654-660
Mineral Springs.....	660-664
Water-Power	660
Present Development.....	661
Greenbrier River.....	661
Anthony Creek.....	662
Howard Creek.....	662
Muddy Creek.....	662
Meadow River.....	662-664
Indicated Horse-Power of Streams.....	665-671
Iron Ore.....	665
General Statement.....	665-670
Oriskany Iron Ore Prospects and Exposures.....	670
Clinton Ores.....	670-671
Table of Iron Ore Analyses.....	672-677
Manganese	672-674
Manganese Prospects.....	674-676
Table of Manganese Ore Analyses.....	
Precious Metals.....	

PART IV. PALEONTOLOGY.

Chapter XIV.—Notes on Paleontology of Greenbrier County (by John L. Tilton and Dana Wells).....	
Fossil Collections from Greenbrier County, W. Va. (Register of Localities by Collection Nos.).....	
Distribution of Collections by Geologic Formations.....	
Pennsylvanian	
Pottsville Series.....	
New River Group.....	
Pocahontas Group.....	
Mississippian	
Monk Chunk Series.....	
Bluestone Group.....	
Hinton Group.....	
Terry Limestone.....	
Avila Limestone.....	

No.	Plates.	Page.
I.—	View from side of Greenbrier Mountain, featuring the grounds of the White Sulphur Springs, Inc. One of the golf courses is shown in the foreground, the casino and Greenbrier Hotel in the center, and Kates Mountain with typical Devonian topography in the background. Photo. by Cummins.....	Frontispiece
II.—	Front view of Greenbrier Hotel at White Sulphur Springs. Photo by Cummins.....	16-A
III.—	View of Howard Creek Valley from Kates Mountain, looking west toward Caldwell. Water-gap in background cut in rapidly dipping Pocono and Chemung Rocks. Photo. by Cummins.....	16-B
IV.—	View from U. S. Route 60 near Clintonville showing Mauch Chunk and Greenbrier topography. The cone-shaped Wenver Knob can be faintly seen in the background.....	16-C
V.—	View from U. S. Route 219, near Renick, showing Greenbrier Limestone topography with sink-holes.....	16-D
VI.—	Outcrop of Pickaway Limestone and Shale in Monroe County on the east or right limb of a symmetrical anticline. Note the vertical fracture cleavage developed in the shale. Note also that the joints in the overlying limestone are normal to the bedding. This outcrop apparently proves that the formation of the joints in the limestone, described under "Pickaway Limestone" in Chapter VII, was independent of and occurred prior to the major folding of the Appalachians.....	144-A
VII.—	Small local fault in top of the Pocono along the C. & O. R. R. track at Caldwell.....	144-B
VIII.—	Drag folding in interbedded limestones and shales of the Rondout Group, 0.5 mile west of Alvon.....	144-C
IX.—	Gouge in bedding-plane fault in the Niagara Series, north side of Anthony Creek, 0.5 mile west of Alvon.....	144-D
X.—	Pottsville Sandstone (Guyandot) on Little Rocky Rim of South Fork of Cherry River.....	224-A
XI.—	View of operations in the Sewell Coal just east of Quinwood	224-B
XII.—	Erratic boulders from the Sewell Coal, Greenbrier County. No. 28 is a granite that was broken by the miners. Photo. by Paul H. Price.....	224-C
XIII.—	Fossil plant showing attached rootlets. Pottsville Sandstone (Upper Raleigh) at Duo.....	224-D
XIV.—	Gullying in the red shale of the Hinton Group of the Mauch Chunk Series, near Rupert. Photo. taken in 1930 See Plate XV for picture in 1931.....	256-A
XV.—	Gullying in the red shale of the Hinton Group of the Mauch Chunk Series, near Rupert. Gullies have deepened two to three feet in one year. Photo taken in 1931	

Bluefield Group.....	697-701
Ada Shale.....	698
Reynolds Limestone.....	698-699
Glenray Limestone.....	699-700
Lillydale Shale.....	700-701
Greenbrler Series.....	701-714
Alderson Limestone.....	701-705
Greenville Shale.....	705
Union Limestone.....	705-706
Plekaway Limestone.....	706
Taggard Limestone.....	706
Patton Limestone.....	707
Sinks Grove Limestone.....	707
Hillsdale Limestone.....	707-714
Discussion of Relation Between Maerady and Hillsdale (St. Louis).....	713-714
Pocono Series	714-717
Broad Ford Sandstone.....	715-717
Devonian	717-733
Chemung Series.....	717-725
Portage Series.....	725-728
Genesee Series.....	728-729
Hamilton Series.....	729-730
Maerellus Series.....	730
Oriskany Series.....	731
Helderberg Series.....	731-733
Beerast Limestone.....	731-733
New Scotland Member.....	733
Silurian	733-734
Sallna Series.....	733-734
Bossardville Member.....	733 734
Niagara Series.....	734
Clinton Series.....	734
Appendix.—Levels Above Mean Tide.....	735-769
Railroad Levels.....	735-737
Chesapeake & Ohio Railroad—Main Line.....	735
Chesapeake & Ohio Railroad—Greenbrler Branch.....	736
Nicholas, Fayette, and Greenbrler Railroad, Formerly Sewell Valley Railroad.....	737
United States Geological Survey Levels.....	737-769
Alderson Quadrangle.....	739-740
Callaghan Quadrangle.....	740-742
Clintonville Quadrangle.....	743-748
Lobelia Quadrangle.....	748-753
Marlinton Quadrangle.....	753-755
Meadow Creek Quadrangle.....	755-756
Richwood Quadrangle.....	756-759
Roneeverte Quadrangle.....	759-760
White Sulphur Springs Quadrangle.....	761-769
Winona Quadrangle.....	769
Index	770-846

XLII.—Hearing Springs Sandstone exposed in stream bed where Howard Creek cuts through Bobs Ridge.....	320-D
XLII.—Apparent unconformity at or near the base of Helderberg Limestone. At spring-house of Alvon Springs Nos. 1 and 2 southwest of Alvon.....	336-A
XLIII.—Niagara Limestone showing numerous veins of calcite, on the north side of Anthony Creek, 0.5 mile west of Alvon	336-B
XLIV.—Anthony Cave, formed by a fold in the Keefer Sandstone. On north side of Anthony Creek, 0.45 mile west of Alvon	336-C
XLV.—La Barr Evergreen Nursery north of White Sulphur Springs	336-D
XLVI.—Crushing and screening plant of the Acme Limestone Company west of Fort Spring. Photo. by courtesy of Acme Limestone Company.....	624-A
XLVII.—Putting off a large shot at the Acme Limestone Company quarry (No. 5 on Map II) 0.9 mile west of Fort Spring. Photo. by courtesy of Acme Limestone Company	624-B
XLVIII.—Acme Limestone Company quarry face (No. 5 on Map II). Note drilling machines and stripping the limestone of soil by hydraulic "gun".....	624-C
IL.—Small shot in the Frazier Limestone quarry (No. 4 on Map II), 1.4 miles southwest of Fort Spring.....	624-D
L.—The Black Sulphur Spring at White Sulphur Springs. This lovely pavilion is situated just in the rear of the Greenbrier Hotel. The spring rises in the center, and the waters from the other springs are brought to the dispensing fountain where all who wish may drink of them freely. This is said to be the oldest spring pavilion in the United States, being erected about 1818. Photo. by Cummins	656-A
LI.—Alvon Springs, spring-house, pump-house, and bottling-house, owned by White Sulphur Springs, Inc. This is the source of the water used in the town of White Sulphur Springs. View looking northeast.....	656-B
LII.—Blue Sulphur Springs. The pavilion shown above is over one hundred years old, mute reminder of times when this spring was the site of a famous resort. The hotel and bath-houses were destroyed during the Civil War and never rebuilt.....	656-C
LIII.—Airplane photograph of Rainelle and the lumber yards of Meadow River Lumber Company.....	656-D

Figures.

1.—Map of West Virginia showing Greenbrier County area and area covered by Detailed Reports.....	xxiii
2.—Map showing the Physiographic Provinces in Greenbrier County and surrounding territory as modified after N. M. Fennerman.....	28

XVI.—Terry Limestone just below the Princeton Conglomerate on the South Fork of Cherry River.....	256-C
XVII.—Subaerial scouring in the Hinton Group of the Mauch Chunk along Midland Trail (U. S. Route 60), 1.2 miles southeast of Crawley.....	256-D
XVIII.—Cross-bedding in the Webster Springs Sandstone (Mauch Chunk) one mile southwest of Modoe P. O., on Falling Spring Mountain.....	272-A
XIX.—Greenbrier Limestone topography near Lewisburg. Mauch Chunk hills in background.....	272-B
XX.—Stylolites in basal Greenbrier Limestone along Midland Trail (U. S. Route 60) west of Lewisburg.....	272-C
XXI.—Mud-cracks in Taggard Limestone along Midland Trail (U. S. Route 60) one mile west of Alta.....	272-D
XXII.—Greenbrier Limestone stripped of cover for quarrying at the Acme Limestone Company quarry near Fort Spring. Note typical Pickaway joints.....	272-E
XXIII.—Typical joints in Pickaway Limestone near Union, Monroe County. Cross-section view of bed.....	272-F
XXIV.—Typical joints in Pickaway Limestone near Union, Monroe County. Top of bed shown.....	272-G
XXV.—Typical joints in Pickaway Limestone near Union, Monroe County.....	272-H
XXVI.—Chert nodules weathered in relief in Hillsdale Limestone on Mill Creek, 1.6 miles south of Asbury.....	288-A
XXVII.—Quarrying road material from the Broad Ford Sandstone in a road cut east of Caldwell.....	288-B
XXVIII.—Broad Ford Sandstone in C. & O. Railroad cut at Caldwell	288-C
XXIX.—Basal Pocono sandstone conglomerate (Berea?) along Midland Trail (U. S. Route 60) 0.9 mile east of Caldwell	288-D
XXX.—Sandstone conglomerate (Berea?) at or near the base of the Pocono near Anthony.....	304-A
XXXI.—Giant ripple-marks in basal Pocono conglomerate on Meadow Creek, 2.6 miles southeast of Neola.....	304-B
XXXII.—Natural whetstones formed by jointing in Chemung sandstone on Kates Mountain.....	304-C
XXXIII.—Close folding in Portage strata along Anthony Creek, north of Neola.....	304-D
XXXIV.—Close folding in Portage strata along Anthony Creek, north of Neola.....	304-E
XXXV.—Interfingering folds in Portage strata, north of Neola....	304-F
XXXVI.—Portage strata in bed of Anthony Creek, 2.3 miles southwest of Neola.....	304-G
XXXVII.—Portage shale and flaggy sandstone along the C. & O. Railroad at Tuckahoe.....	304-H
XXXVIII.—Mareellus Shale showing calcareous (Onondaga?) beds, near the mouth of Slash Lick Run.....	320-A
XXXIX.—Contorted Mareellus Shale, one mile northwest of	320-B

This book is a general geologic and economic report on Greenbrier County, West Virginia. As shown in the Table of Contents, it contains a chapter on Historical and Industrial Development, a chapter on Physiography, seven chapters on Geologic History, Structure, and Stratigraphy, four chapters on Mineral Resources, and one chapter on Paleontology, as well as an Appendix giving all available spirit-level bench marks and railroad levels for the county.

In a separate Atlas, Maps I and II, respectively, show the topography and geology of the county. For these maps the topographic base was assembled and photolithographed from the standard topographic quadrangles as surveyed and published by the United States Geological Survey in cooperation with the West Virginia Geological Survey, with certain cultural corrections added by the authors. On this corrected base the geologic map was drawn.

The field work for this report was begun by Price in June, 1929, and continued by him during the summer months of 1930 and 1931. Price was assisted in the field during the summers of 1929 and 1930 by John P. Nolting, Jr. During the first half of the three summer months of 1931, Price was assisted by Charles W. Furbee, Jr. His assistant during the latter half was E. T. Heek. Lack of available funds caused virtual suspension of the work on this report during 1932, 1933, and 1934. The appointment of Price as State Geologist prevented his resuming field work on the report and the task of completing it was assigned to Heek in July, 1935. The field work was brought up to date, as of 1937, and completed by Heek under the direction of Price during the years of 1935, 1936, and 1937. Heek was assisted for short periods by Charles E. Hare and S. S. Galpin. The manuscript was completed in December, 1938. The chapter on Paleontology is the work of the late Dr. John L. Tilton and Professor Dana Wells, present cooperating Paleontologist. The chemical tests, except as otherwise specified in the text, were made by B. B. Kaplan and Homer A. Hoskins, Survey Chemists.

Including a portion of both the plateau and folded Appalachians regions, Greenbrier County offers a most interesting area for geologic study. The outcropping rocks, including those from the lower Silurian to the Kanawha Group of the Pennsylvanian, embrace a total of about 14,385 feet of strata and contain large quantities of coal, limestone, building stone, clays, iron ore, and some manganese ore. In addition, the

5.—Same as Figure 4, with superimposed projected profiles.....	31
6.—Map of Greenbrier County showing drainage.....	38
7.—General Columnar Section of rocks exposed in Greenbrier County	131-3
8.—Map of Greenbrier County showing outcrops of Pottsville Series	213
9.—Map of Greenbrier County showing outcrops of Mauch Chunk Series.....	257
10.—Map of Greenbrier County showing outcrops of Greenbrier Series	269
11.—Map showing localities where the direction of Pickaway joints was measured, and near-by major structural features	275
12.—Map showing outcrops of Maecrady and Pocono Series.....	285
13.—Map showing outcrops of Upper and Middle Devonian Rocks	298
14.—Map showing outcrops of Lower Devonian Rocks.....	316
15.—Map showing outcrops of Silurian Rocks.....	328
16.—Diagram showing the position and thickness of coal seams in Greenbrier County.....	364
17.—Map showing probable area of minable Sewell Coal.....	474
18.—"Fault", Imperial Smokeless Coal Co. mine.....	482
19.—Map showing probable area of minable Little Raleigh Coal and Beekley Coal.....	531
20.—Map showing probable area of minable Fire Creek Coal....	555
21.—Map showing probable area of minable No. 6 Pocahontas Coal	581
22.—Map showing the location of recent prospect openings in the No. 6 Pocahontas Coal.....	590
23.—Map showing probable area of minable No. 3 Pocahontas Coal	602

Whorter, Mr. G. W. Watts, Mr. J. C. Kennedy, Mr. Swing, Mr. W. W. Coleman, Mr. J. W. Raine, Mr. B. L. erts, Mr. R. B. Holt, Mr. H. H. Blackburn, and Mr. Tuckwiller whose extraordinary interest in mineral m and whose wide knowledge of many interesting outcrop exposures have materially added to the value of the r

PAUL H. PRICE

E. T. HECK.

Morgantown, W. Va., December 15, 1938.

ERRATA.

- Page 12, line 2 from top, for Division, read District.
Page 27, line 17 from bottom, for Costal, read Coastal.
Page 28, Reverse figure. Top is on binding edge.
Page 36, line 17 from bottom, for channel, read channels.
Page 39, first line of table, for source to, read source of.
Page 42, line 11 of table, for Preaser Branch, read Peaser Branch.
Page 44, line 13 of table, for Bear Run, read Bear Branch.
Page 45, line 14 of table, for Old Knob Branch, read Job Knob Branch.
Page 76, transfer heading to part of table above years 1922-23.
Page 115, line 12 from top, for subferraneus, read subterraneus.
Page 116, line 13 from bottom, for bettles, read beetles.
Page 155, line 11 from top, for basel, read basal.
Page 188, line 26 from bottom, add Section after Renick Station.
Page 204, lines 11 and 12 from bottom, for Rensselaria, read Rensselaeria.
Pages 210 and 211, for Renick, read Renick Station.
Pages 210 and 211, for Renick Valley, read Renicks Valley.
Page 229, line 14 from bottom, for number, read member.

LETTER OF TRANSMITTAL

To His Excellency, Honorable Homer A. Holt, Governor of West Virginia, and President of the Geological Survey Commission.

SIR:

I have the honor and pleasure to transmit herewith for publication the Detailed Geologic Report and accompanying topographic and geologic maps covering Greenbrier County prepared by myself and E. T. Heck.

The county contains 1022.8 square miles of territory and is, therefore, the second largest in point of size in the State. Greenbrier County is rich in many ways including minerals, soils, timber, waters, climate, and especially her fine people. While other counties in the State may have fared better at the hands of Providence in some mineral resources certainly none has been more favorably blessed from the point of view of diversity of resources. The entire county was, before the coming of the white man, entirely covered with a fine growth of timber with hardwoods predominating. It is interesting to note that the forests of the county can be roughly divided into three districts just as can the geology, and is, of course, a reflection of the latter. The mountainous sections of the west and northwest are characterized by spruce, hemlock, and yellow birch and others that thrive at high altitudes, with hardwoods predominating below 3,000 feet. The main limestone section lying between the mountainous area and Greenbrier River produced excellent timber, most of which was hardwoods such as white oak, red oak, poplar, black walnut, hickory, and wild cherry. East of the Greenbrier River to the State line and especially along Anthony and Howard Creeks the predominating species was white pine.

In the western side of the county there is a wide zone of Carboniferous or Pennsylvanian rocks containing large reserves of New River and Pocahontas coals now in active development. West of the Greenbrier River and extending the entire length of the county is a wide belt of Mississippian rocks composed of thick limestones of the Greenbrier Series with overlying beds of red shale and shaly limestones of the Mauch Chunk Series. These rocks when weathered form certain soil types used most successfully for grazing and for cultivation of crops. The Greenbrier Series also affords an inexhaustible supply of limestone suitable for practically all purposes for which limestone may be used, i. e., industrial, chemical, and agricultural. Numerous quarries now operating attest their worth.

In the eastern part of the county, the rocks of Devonian and Silurian are not suitable for agriculture because of their generally siliceous nature but offer possibilities for iron ore and manganese both of which need further study. Sandstones suit-

able for building purposes and clays and shales adaptable to brick and tile manufacture are found throughout the county.

The mineral springs of the county represent one of its valuable resources and these together with its pure streams and agreeable climate have led to the development of the magnificent White Sulphur Springs resort area; the several boys and girls schools and summer camps; and the many fine summer homes which make Greenbrier County the most attractive vacation land in the country.

The field studies of the agricultural soils have been completed by a soil specialist of the Bureau of Chemistry and Soils of the U. S. Department of Agriculture in cooperation with the West Virginia Geological Survey, and a report together with a soil map will be published in the near future. With its completion we will have a geologic, a topographic, and a soils map as well as a geologic and soils report on each county of the State.

It is especially fitting that this report is released under your administration as Governor since Greenbrier is your native county.

Respectfully submitted,

PAUL H. PRICE,
State Geologist.

Morgantown, W. Va., June 30, 1939.

CONTENTS

	Page.
Geological Survey Commission and State Board of Control.....	iii
Geological Survey Staff.....	v
Letter of Transmittal.....	vi-vii
Table of Contents.....	viii-xvii
List of Illustrations.....	xviii-xxi
Author's Preface.....	xxii-xxiv
Errata.....	xxiv

PART I. HISTORY AND PHYSIOGRAPHY.

Chapter I.—Historical and Industrial Development.....	1-25
Location.....	1
General Description.....	2-12
Miscellaneous Items.....	2-12
Formation.....	2
Area.....	2
Relief.....	3
Climate.....	3-9
Products.....	9
Property Valuation.....	9-10
Postal Service.....	11
Population.....	11-12
Towns and Industries.....	12-16
Transportation.....	16-23
Railroads.....	17-20
Highways.....	20-23
State Roads.....	20-23
Airport.....	23
Resorts and Summer Camps.....	23-25
Chapter II.—Physiography.....	26-117
Introduction.....	26-27
Physiographic Provinces.....	27-30
The Earliest Restored Surface.....	31-33
An Intermediate Surface.....	34
Stream Terraces.....	34
Flood-Plains.....	35-36
Present Topographic Features.....	36-37
Drainage Basins.....	37-110
Table of Stream Data.....	39-42
Drainage Areas of Greenbrier County.....	43-46
Areas of Drainage Basins.....	43-46
Description of Drainage Basins.....	46-110
Greenbrier River.....	46-51
Meadow River.....	52-56
Cherry River.....	57-110
Sinks, Caverns, and Subsurface Drainage.....	111-117
Solution in Carbonate Rocks.....	111
Sinks.....	111-112
Caverns.....	112-114
Present Fauna in the Caves.....	114-117
Subsurface Drainage.....	117

PART II. GEOLOGY.

Chapter III.—Geologic Processes: Erosion and Deposition.....	118-133
Hills and Valleys are Temporary Features.....	118

CONTENTS.

ix

	Page.
Weathering.....	118-119
Effects of Changes of Temperature.....	119
The Processes of Erosion and Deposition Never Cease.....	119-121
Classification of Rocks.....	121
How Sediments Change to Stone.....	121-122
The Sedimentary Rocks.....	122-124
Derivation of Sediments and Implied Environment.....	124-128
Bibliography.....	128-129
Nomenclature and Correlation.....	129-130
Classification of Outcropping Rocks.....	130-133
General Columnar Section of Rocks Exposed in Greenbrier County.....	131-133

Chapter IV.—Structural Geology.....	134-153
Introduction.....	134-135
Methods of Geologic Work and Representation of Structure.....	135-138
Intervals Above and Below Sowell Coal.....	137
Detailed Structure.....	138-153
Anticlines and Synclines.....	138-149
Webster Springs Anticline.....	138-139
Koyan Syncline.....	139-140
Boggs Knob Anticline.....	140
Springdale Syncline.....	141
Alderson Anticline.....	141
Creamery Syncline.....	141
Williamsburg (Mount Pleasant) Anticline.....	141-143
Muddy Creek Mountain Syncline.....	143
Sinks Grove Anticline.....	143-144
Caldwell (Fulton) Syncline.....	144-145
Maple Grove Anticline.....	145
Hurricane Ridge Syncline.....	145
Browns Mountain Anticline.....	145-147
Neola Anticline.....	147-148
Meadow Creek Syncline.....	148
Kates Mountain Syncline.....	148
Glaze Anticline.....	149
Tuckahoe Syncline.....	149
Cross-Sections.....	149-150
Unconformities.....	150-151
Faults.....	151-153
Burr Funn.....	153

Chapter V.—Measured Sections.....	154-211
Introduction.....	154-156
Measured Sections, Meadow Bluff District.....	156-178
Measured Sections, Williamsburg District.....	179-182
Measured Sections, Falling Springs District.....	182-189
Measured Sections, Blue Sulphur District.....	189-193
Measured Sections, Irish Corner District.....	193-198
Measured Sections, Fort Springs District.....	198-199
Measured Sections, Lewisburg District.....	199-200
Measured Sections, Frankford District.....	200-201
Measured Sections, White Sulphur District.....	201-205
Measured Sections, Anthony Creek District.....	205-208
Summary of Measured Sections.....	209-211
Summary of Measured Sections (Table).....	210-211

Chapter VI.—Stratigraphy—Pennsylvanian Rocks.....	212-251
Introduction.....	212-213

	Page.
General Account and Section, Pottsville Series.....	214-217
General Section, Pottsville Series.....	215-217
Topographic Expression.....	218
Contacts and Unconformities.....	218-219
Fossil Life.....	219
Correlation, Pottsville Series.....	219-220
Description of Members, Kanawha Group.....	220-222
Lower Gilbert Sandstone.....	221
Gilbert "A" Coal.....	221
Gilbert Shale.....	221
Gilbert Coal.....	221
Dotson Sandstone.....	221
Douglas "A" Coal and Douglas Coal.....	221
Lower Dotson Sandstone.....	222
Douglas Shale.....	222
Description of Members, New River Group.....	222-242
Upper Nuttall Sandstone.....	223
Jaeger "B" Coal.....	223-224
Lower Nuttall Sandstone.....	224
Jaeger "A" Coal.....	224
Upper Jaeger Shale.....	224
Hughes Ferry Coal.....	225
Middle Jaeger Sandstone.....	225
Lower Jaeger Coal.....	225
Lower Jaeger Sandstone.....	225
Lower Jaeger Shale.....	225
Harvey (Conglomerate) Sandstone.....	226
Sandy Huff Shale.....	226-227
Castle Coal.....	227
Guyandot Sandstone.....	227
Skelt Shale.....	227
Sewell "B" Coal.....	227
Sewell "A" Coal.....	227
Lower Guyandot Sandstone.....	227
Harridge Black Shale.....	227
Sewell Coal.....	229-230
Erratic Boulders in the Sewell Coal.....	230-231
Welch Sandstone.....	231
Welch Coal.....	231
Upper Raleigh Sandstone.....	234-235
Little Raleigh "A" Coal.....	235
Little Raleigh Coal.....	235-236
Lower Raleigh Sandstone.....	236
Beckley "Rider" Coal.....	236
Beckley Coal.....	236-237
Quinnlont Sandstone.....	237-238
Quinnlont Shale.....	238
Fire Creek Coal.....	238
Little Fire Creek Coal.....	239-240
Pineville Sandstone.....	240
No. 9 Pocahontas Coal.....	241
No. 8 Pocahontas Coal.....	241-242
Description of Members, Pocahontas Group.....	242-243
Flattop Mountain Sandstone.....	242-243
Rift Shale, No. 7 Pocahontas Coal, Pierpont Sandstone, Royal Shale, No. 6 Pocahontas Coal.....	243-244
Eckman Sandstone.....	244
No. 5 Pocahontas Coal and No. 4 Pocahontas Coal.....	246-247

	Page.
Upper Pocahontas Sandstone.....	248
No. 3 Pocahontas "Rider" Coal.....	248
No. 3 Pocahontas Coal.....	249
Lower Pocahontas Sandstone.....	249
No. 2 "A" Pocahontas Coal.....	249
No. 2 Pocahontas Coal.....	250
No. 1 Pocahontas Coal.....	250
Economic Aspects, Pottsville Series.....	251
Chapter VII.—Stratigraphy—Mississippian Rocks.....	252-284
General Statement.....	252
Correlation, Mississippian Period.....	252-254
Mauch Chunk Series.....	254-255
General Account and Section.....	254-255
General Section.....	255
Topographic Expression.....	256
Areal Extent.....	256-257
Contacts.....	258
Fossil Life.....	258
Correlation.....	258-259
Description of Members, Bluestone Group.....	259-260
Description of Members, Princeton Group.....	260
Princeton Sandstone.....	260
Description of Members, Hinton Group.....	260-262
Avis Limestone.....	261
Stony Gap Sandstone.....	261-262
Description of Members, Bluefield Group.....	262-265
Droop Sandstone.....	263
Talcott and Ada Shales.....	263
Reynolds Limestone.....	264
Webster Springs Sandstone.....	264
Glenray Limestone.....	264-265
Lillydale Shale.....	265
Economic Aspects.....	265
Greenbrier Series.....	266-279
General Account and Section.....	266-267
General Section.....	267
Topographic Expression.....	267-268
Areal Extent.....	268-269
Contacts.....	270
Fossil Life.....	270
Correlation.....	270
Description of Members.....	271-279
Alderson Limestone.....	271
Greenville Shale.....	271-272
Union Limestone.....	272
Pickaway Limestone.....	272-277
Taggard Limestone.....	277-278
Fenton Limestone.....	278
Sinks Grove Limestone.....	278-279
Hillsdale Limestone.....	279
Economic Aspects.....	279
Maccrady Series.....	280-282
General Account.....	280
Topographic Expression.....	280
Areal Extent.....	281
Contacts.....	281
Fossil Life.....	281

	Page.
Correlation	281-282
Economic Aspects.....	282
Pocono Series.....	283-294
General Account and Section.....	283
General Section.....	283
Topographic Expression.....	284
Areal Extent.....	284-285
Contacts	286
Fossil Life.....	286-289
Correlation	289
Description of Members.....	290-294
Merrimac Coal.....	290-292
Broad Ford Sandstone.....	292-293
Berea Sandstone.....	293-294
Economic Aspects.....	294
Chapter VIII.—Stratigraphy—Devonian Rocks.....	295-326
General Statement.....	295-296
Upper Devonian Rocks.....	296-309
Catskill Series.....	296-300
General Account.....	296-297
Topographic Expression.....	297
Areal Extent.....	297-298
Contacts	299
Fossil Life.....	300
Correlation	300
Economic Aspects.....	300
Chemung Series.....	300-304
General Account and Section.....	300-301
General Section.....	301
Topographic Expression.....	300-301
Areal Extent.....	302
Contacts	302
Fossil Life.....	303
Correlation	303
Description of Members.....	303
Hendricks Sandstone.....	303
Economic Aspects.....	304
Portage Series.....	304-307
General Account.....	304-305
Topographic Expression.....	305
Areal Extent.....	306
Contacts	306
Fossil Life.....	306
Correlation	306-307
Economic Aspects.....	307
Genesee Series.....	307-309
General Account.....	307-308
Topographic Expression.....	308
Areal Extent.....	308
Contacts	308
Fossil Life.....	309
Correlation	309
Description of Members.....	309
Economic Aspects.....	309
Middle Devonian Rocks.....	310-313
General Statement.....	310
Marcellus Series.....	310-313
General Account.....	310-311

GEOLOGICAL SURVEY STAFF

PAUL H. PRICE.....*State Geologist*
RIETZ C. TUCKER.....*Assistant State Geologist*
CHARLES E. HARE.....*Assistant Geologist*
E. T. HECK.....*Assistant Geologist*
JOHN P. NOLTING, JR.....*Assistant Geologist*
J. B. McCUE.....*Chief Chemist*
A. J. W. HEADLEE.....*Chemist and Gas Analyst*
HOMER A. HOSKINS.....*Chemist*
CHARLES B. WAGNER.....*Assistant Chemist*
JAMES H. C. MARTENS.....*Mineralogist*
JOHN L. TILTON (deceased, November 17, 1930) *Paleontologist*
DANA WELLS.....*Cooperating Paleontologist*
MARIE S. AZURE.....*Secretary-Bookkeeper*
W. J. HESSLER.....*Clerk*
IRENE SPEICHER.....*Stenographer-Typist*

GEOLOGICAL SURVEY STAFF

PAUL H. PRICE.....*State Geologist*

RIETZ C. TUCKER.....*Assistant State Geologist*

CHARLES E. HARE.....*Assistant Geologist*

E. T. HECK.....*Assistant Geologist*

JOHN P. NOLTING, JR.....*Assistant Geologist*

J. B. McCUE.....*Chief Chemist*

A. J. W. HEADLEE.....*Chemist and Gas Analyst*

HOMER A. HOSKINS.....*Chemist*

CHARLES B. WAGNER.....*Assistant Chemist*

JAMES H. C. MARTENS.....*Mineralogist*

JOHN L. TILTON (deceased, November 17, 1930) *Paleontologist*

DANA WELLS.....*Cooperating Paleontologist*

MARIE S. AZURE.....*Secretary-Bookkeeper*

W. J. HESSLER.....*Clerk*

IRENE SPEICHER.....*Stenographer-Typist*

able for building purposes and clays and shales adaptable to brick and tile manufacture are found throughout the county.

The mineral springs of the county represent one of its valuable resources and these together with its pure streams and agreeable climate have led to the development of the magnificent White Sulphur Springs resort area; the several boys and girls schools and summer camps; and the many fine summer homes which make Greenbrier County the most attractive vacation land in the country.

The field studies of the agricultural soils have been completed by a soil specialist of the Bureau of Chemistry and Soils of the U. S. Department of Agriculture in cooperation with the West Virginia Geological Survey, and a report together with a soil map will be published in the near future. With its completion we will have a geologic, a topographic, and a soils map as well as a geologic and soils report on each county of the State.

It is especially fitting that this report is released under your administration as Governor since Greenbrier is your native county.

Respectfully submitted,

PAUL H. PRICE,
State Geologist.

Morgantown, W. Va., June 30, 1939.

SIR:

I have the honor and pleasure to transmit herewith for publication the Detailed Geologic Report and accompanying topographic and geologic maps covering Greenbrier County prepared by myself and E. T. Heck.

The county contains 1022.8 square miles of territory and is, therefore, the second largest in point of size in the State. Greenbrier County is rich in many ways including minerals, soils, timber, waters, climate, and especially her fine people. While other counties in the State may have fared better at the hands of Providence in some mineral resources certainly none has been more favorably blessed from the point of view of diversity of resources. The entire county was, before the coming of the white man, entirely covered with a fine growth of timber with hardwoods predominating. It is interesting to note that the forests of the county can be roughly divided into three districts just as can the geology, and is, of course, a reflection of the latter. The mountainous sections of the west and northwest are characterized by spruce, hemlock, and yellow birch and others that thrive at high altitudes, with hardwoods predominating below 3,000 feet. The main limestone section lying between the mountainous area and Greenbrier River produced excellent timber, most of which was hardwoods such as white oak, red oak, poplar, black walnut, hickory, and wild cherry. East of the Greenbrier River to the State line and especially along Anthony and Howard Creeks the predominating species was white pine.

In the western side of the county there is a wide zone of Carboniferous or Pennsylvanian rocks containing large reserves of New River and Poahontas coals now in active development. West of the Greenbrier River and extending the entire length of the county is a wide belt of Mississippian rocks composed of thick limestones of the Greenbrier Series with overlying beds of red shale and shaly limestones of the Mauch Chunk Series. These rocks when weathered form certain soil types used most successfully for grazing and for cultivation of crops. The Greenbrier Series also affords an inexhaustible supply of limestone suitable for practically all purposes for which limestone may be used, i. e., industrial, chemical, and agricultural. Numerous quarries now operating attest their worth.

In the eastern part of the county, the rocks of Devonian and Silurian are not suitable for agriculture because of their generally siliceous nature but offer possibilities for iron ore and manganese both of which need further study. Sandstones suit-

Weathering	118-119
Effects of Changes of Temperature.....	119
The Processes of Erosion and Deposition Never Cease.....	119-121
Classification of Rocks.....	121
How Sediments Change to Stone.....	121-122
The Sedimentary Rocks.....	122-124
Derivation of Sediments and Implied Environment.....	124-128
Bibliography	128-129
Nomenclature and Correlation.....	129-130
Classification of Outcropping Rocks.....	130-133
General Columnar Section of Rocks Exposed in Greenbrier County.....	131-133
Chapter IV.—Structural Geology.....	134-153
Introduction	134-135
Methods of Geologic Work and Representation of Structure	135-138
Intervals Above and Below Sewell Coal.....	137
Detailed Structure.....	138-153
Anticlines and Synclines.....	138-149
Webster Springs Anticline.....	138-139
Kovan Syncline.....	139-140
Boggs Knob Anticline.....	140
Springdale Syncline.....	141
Alderson Anticline.....	141
Creamery Syncline.....	141
Williamsburg (Mount Pleasant) Anticline.....	141-143
Muddy Creek Mountain Syncline.....	143
Sinks Grove Anticline.....	143-144
Caldwell (Patton) Syncline.....	144-145
Maple Grove Anticline.....	145
Hurricane Ridge Syncline.....	145
Browns Mountain Anticline.....	145-147
Neola Anticline.....	147-148
Meadow Creek Syncline.....	148
Kates Mountain Syncline.....	148
Glaze Anticline.....	149
Tuckahoe Syncline.....	149
Cross-Sections	149-150
Unconformities	150-151
Faults	151-153
Burr Fault.....	153
Chapter V.—Measured Sections.....	154-211
Introduction	154-156
Measured Sections, Meadow Bluff District.....	156-178
Measured Sections, Williamsburg District.....	179-182
Measured Sections, Falling Springs District.....	182-189
Measured Sections, Blue Sulphur District.....	189-195
Measured Sections, Irish Corner District.....	196-198
Measured Sections, Fort Springs District.....	198-199
Measured Sections, Lewisburg District.....	199-200
Measured Sections, Frankford District.....	200-201
Measured Sections, White Sulphur District.....	201-205
Measured Sections, Anthony Creek District.....	205-208
Summary of Measured Sections.....	209-211
Summary of Measured Sections (Table).....	210-211

	Page.
Geological Survey Commission and State Board of Control.....	iii
Geological Survey Staff.....	v
Letter of Transmittal.....	vi-vii
Table of Contents.....	viii-xvii
List of Illustrations.....	xviii-xxi
Author's Preface.....	xxii-xxlv
Errata	xxlv

PART I. HISTORY AND PHYSIOGRAPHY.

Chapter I.—Historical and Industrial Development.....	1-25
Location	1
General Description.....	2-12
Miscellaneous Items.....	2-12
Formation	2
Area	2
Relief	3
Climate	3-9
Products	9
Property Valuation.....	9-10
Postal Service.....	11
Population	11-12
Towns and Industries.....	12-16
Transportation	16-23
Railroads	17-20
Highways	20-23
State Roads.....	20-23
Airport	23
Resorts and Summer Camps.....	23-25
Chapter II.—Physiography.....	26-117
Introduction	26-27
Physiographic Provinces.....	27-30
The Earliest Restored Surface.....	31-33
An Intermediate Surface.....	34
Stream Terraces.....	34
Flood-Plains	35-36
Present Topographic Features.....	36-37
Drainage Basins.....	37-110
Table of Stream Data.....	39-42
Drainage Areas of Greenbrier County.....	43-46
Areas of Drainage Basins.....	43-46
Description of Drainage Basins.....	46-110
Greenbrier River.....	46-81
Meadow River.....	82-96
Cherry River.....	97-110
Sinks, Caverns, and Subsurface Drainage.....	111-117
Solution in Carbonate Rocks.....	111
Sinks	111-112
Caverns	112-114
Present Fauna in the Caves.....	114-117
Subsurface Drainage.....	117

PART II. GEOLOGY.



GEOLOGICAL SURVEY STAFF

PAUL H. PRICE.....*State Geologist*
RIETZ C. TUCKER.....*Assistant State Geologist*
CHARLES E. HARE.....*Assistant Geologist*
E. T. HECK.....*Assistant Geologist*
JOHN P. NOLTING, JR.....*Assistant Geologist*
J. B. McCUE.....*Chief Chemist*
A. J. W. HEADLEE.....*Chemist and Gas Analyst*
HOMER A. HOSKINS.....*Chemist*
CHARLES B. WAGNER.....*Assistant Chemist*
JAMES H. C. MARTENS.....*Mineralogist*
JOHN L. TILTON (deceased, November 17, 1930) *Paleontologist*
DANA WELLS.....*Cooperating Paleontologist*
MARIE S. AZURE.....*Secretary-Bookkeeper*
W. J. HESSLER.....*Clerk*
IRENE SPEICHER.....*Stenographer-Typist*